OPEN

American Association for the Surgery of Trauma/American College of Surgeons Committee on Trauma clinical protocol for management of acute respiratory distress syndrome and severe hypoxemia

Jason A. Fawley, MD, FACS, Christopher J. Tignanelli, MD, MS, FACS, Nicole L. Werner, MD, MS, FACS, George Kasotakis, MD, MPH, FACS, Samuel P. Mandell, MD, MPH, FACS, Nina E. Glass, MD, FACS, David J. Dries, MD, FACS, MCCM, Todd W. Costantini, MD, FACS, *and* Lena M. Napolitano, MD, FACS, FCCP, MCCM, *Ann Arbor, Michigan*

CONTINUING MEDICAL EDUCATION CREDIT INFORMATION

Accreditation

In support of improving patient care, this activity has been planned and implemented by CineMed and the American Association for the Surgery of Trauma. CineMed is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

AMA PRA Category 1 Credits™

CineMed designates this enduing material for a maximum of 1 AMA PRA Category 1 Credit(s)TM. Physicians should claim only the credit commensurate with the extent of their participation in the activity.



JOINTLY ACCREDITED PROVIDER"

Objectives

After reading the featured articles published in the *Journal of Trauma and Acute Care Surgery*, participants should be able to demonstrate increased understanding of the material specific to the article. Objectives for each article are featured at the beginning of each article and online. Test questions are at the end of the article, with a critique and specific location in the article referencing the question topic.

Disclosure Information

In accordance with the ACCME Accreditation Criteria, CineMed must ensure that anyone in a position to control the content of the educational activity (planners and speakers/authors/discussants/moderators) has disclosed all financial relationships with any commercial interest (termed by the ACCME as "ineligible companies", defined below) held in the last 36 months (see below for definitions). Please note that first authors were required to collect and submit disclosure information on behalf all other authors/contributors, if applicable.

Ineligible Company: The ACCME defines an "ineligible company" as any entity producing, marketing, selling, re-selling, or distributing health care goods or services used on or consumed by patients. Providers of clinical services directly to patients are NOT included in this definition.

Financial Relationships: Relationships in which the individual benefits by receiving a salary, royalty, intellectual property rights, consulting fee, honoraria, ownership interest (e.g., stocks, stock options or other ownership interest, excluding diversified mutual funds), or other financial benefit. Financial benefits are usually associated with roles such as employment, management position, independent contractor (including contracted research), consulting, speaking and teaching, membership on advisory committees or review panels, board membership, and other activities from which remuneration is received, or expected.

Conflict of Interest: Circumstances create a conflict of interest when an individual has an opportunity to affect CME content about products or services of a commercial interest with which he/she has a financial relationship.

The ACCME also requires that CineMed manage any reported conflict and eliminate the potential for bias during the session. Any conflicts noted below have been managed to our satisfaction. The disclosure information is intended to identify any commercial relationships and allow learners to form their own judgments. However, if you perceive a bias during the educational activity, please report it on the evaluation. All relevant financial relationships have been mitigated.

AUTHORS/CONTRIBUTORS

George Kasotakis, Alcresta Inc., Grant, Research Investigator. Samuel P. Mandell, UptoDate, Royalty, Author. Jason A. Fawley, Christopher J. Tignanelli, Nicole L. Werner, Nina E. Glass, David J. Dries, Todd W. Costantini, and Lena M. Napolitano have nothing to disclose.

EDITOR-IN-CHIEF/DEPUTY EDITORS/ ASSOCIATE EDITORS

Conflict of Interest forms for all Journal of Trauma and Acute Care Editors have been supplied and are provided as Supplemental Digital Content (http://links.lww.com/TA/D55).

Claiming Credit

To claim credit, please visit the AAST website at http://www.aast.org/ and click on the "e-Learning/MOC" tab. You must read the article, successfully complete the post-test and evaluation. Your CME certificate will be available immediately upon receiving a passing score of 75% or higher on the post-test. Post-tests receiving a score of below 75% will require a retake of the test to receive credit.

Credits can only be claimed online

Cost

For AAST members and Journal of Trauma and Acute Care Surgery subscribers there is no charge to participate in this activity. For those who are not a member or subscriber, the cost for each credit is \$25.

Questions

If you have any questions, please contact AAST at 800-789-4006. Paper test and evaluations will not be accepted.

LEVEL OF EVIDENCE:	Therapeutic/Care Management: Level V. (J Trauma Acute Care Surg. 2023;95: 592–602. Copyright © 2023 The Author(s). Published
	by Wolters Kluwer Health, Inc.)
KEY WORDS:	Acute respiratory distress syndrome; severe hypoxemia; low tidal volume ventilation; positive end-expiratory pressure; driving
	pressure; plateau pressure; recruitment maneuver; prone position; inhaled nitric oxide; neuromuscular blockade; extracorporeal
	membrane oxygenation.

A cute respiratory distress syndrome (ARDS) is an acute, diffuse lung injury characterized by hypoxemia (Table 1).^{1,2} It was first described in 1967 in 12 patients as acute hypoxic lung injury with markedly reduced pulmonary compliance related to trauma in seven patients, and aspiration/viral pneumonia in five patients.³

Globally, ARDS affects approximately 3 million patients annually, with over 200,000 in the United States. ARDS was present in 10.4% of critically ill patients in a multicenter large international study including nearly 30,000 intensive care unit (ICU) patients, with a hospital mortality of 46.1% in those with most severe hypoxemia.⁴ Acute respiratory distress syndrome accounted for 24% of patients receiving mechanical ventilation in the ICU. Common etiologies of ARDS include pneumonia (bacterial, viral, fungal), aspiration, sepsis, trauma, blood transfusion, burns and inhalation injury, and pancreatitis. A US multicenter study of 2466 moderate/severe ARDS patients reported 28-day mortality rate of 41% (range, 16.7–73.3%), and confirmed substantial variability in ARDS management, with initial adherence to lung protective ventilation of only 31.4%.⁵

Acute respiratory distress syndrome is still associated with high morbidity and mortality in trauma patients. A systematic review of 43 studies reported an overall median ARDS incidence of 8.4% in trauma.⁶ Patients with thoracic injuries, including pulmonary contusions, are particularly at high risk for the development of ARDS, occurring in 10% to 25% of these patients.⁷ Other trauma patients at high risk for ARDS include patients requiring massive transfusion with aggressive fluid and blood product resuscitation, and those with traumatic brain injury. In a meta-analysis of 20 studies (n = 2830), approximately one in five patients had ARDS after traumatic brain injury within a median time of 3 days and was associated with increased mortality and worse neurologic outcome.⁸ ARDS can increase the risk of in-hospital death by three-fold in traumatic brain injury patients.⁹

A systematic review and meta-analysis of prognostic factors for development of ARDS following traumatic injury included 39 studies involving over 5.3 million patients and identified the

DOI: 10.1097/TA.000000000004046

amount of crystalloid resuscitation with the first 24 hours of injury as a modifiable risk factor (adjusted odds ratio, 1.19; 95% confidence interval [CI], 1.15-1.24 for each additional liter of crystalloid administered within the first 6 hours after injury; high certainty). Nonmodifiable prognostic factors with a moderate or high certainty of association with posttraumatic ARDS included increasing age, non-Hispanic White race, blunt mechanism of injury, presence of head injury, pulmonary contusion or rib fracture, and increasing chest injury severity. $^{10}\ {\rm Early}\ {\rm ARDS}$ after trauma due to overresuscitation has decreased due to improved trauma resuscitation strategies, but ARDS related to TBI and pneumonia/sepsis persists. A recent systematic review (20 studies, n = 2830) reported that 19% of patients had ARDS after TBI, with a median time of 3 days. Acute respiratory distress syndrome was associated with higher mortality and worse neurologic outcomes.8 Other studies have confirmed a second peak of ARDS later in the ICU course related to pneumonia and sepsis.11

While the incidence of trauma-induced ARDS has decreased, mortality in trauma patients with ARDS has increased. In the largest study of almost 3 million patients from the National Trauma Data Bank, 1% developed ARDS, but decreased from 1.4% to 0.5% over the 4-year study period. Hospital mortality was 22% in ARDS trauma patients, but increased from 21% to 28% over the study period. Acute respiratory distress syndrome risk factors included direct thoracic injury, increased age, male sex, higher ISS, lower GCS.¹² In another study of over 800,000 Trauma Quality Improvement Program patients between 2010 and 2014, the incidence of ARDS decreased over the study years (3% to 1.1%) but ARDS-related mortality increased (18 to 21%, p = 0.001). Acute respiratory distress syndrome risk factors included older age, male sex, and African American race. Mortality risk factors included older age, male sex, lower Glasgow Coma Scale and higher ISS.¹³

Because many of the ARDS risk factors are nonmodifiable, this protocol focuses on the treatment of ARDS in trauma

TABLE 1. Definition of ARDS

The Berlin criteria definition of ARDS includes:

- Onset of hypoxemia within 7 d of a known clinical insult or worsening respiratory symptoms
- Chest imaging with bilateral opacities not fully explained by effusions, lobar/lung collapse, or nodules
- Pulmonary edema not fully explained by cardiac failure or fluid overload
- Hypoxemia defined as a PaO_2/FIO_2 ratio ${\leq}300$ mmHg with PEEP or CPAP ${\geq}5$ cm H_2O
 - $^{\circ}$ Mild ARDS: PaO_2/FIO_2 ratio 201–300 mm Hg
 - ° Moderate ARDS: PaO2/FIO2 ratio 101-200 mm Hg
 - ° Severe ARDS: PaO₂/FIO₂ ratio ≤100 mm Hg

From: The ARDS Definition Task Force. Acute respiratory distress syndrome: The Berlin Definition. JAMA 2012;307:2526–2533.

Submitted: March 12, 2023, Revised: April 27, 2023, Accepted: April 28, 2023, Published online: June 12, 2023.

From the Departments of Surgery of Scripps Health (J.F.), La Jolla, California; University of Minnesota (C.J.T.), Minneapolis, Minnesota; University of Wisconsin-Madison School of Medicine (N.W.), Madison, Wisconsin; Duke University School of Medicine (G.K.), Durham, North Carolina; University of Texas Southwestern (S.P.M.), Dallas, Texas; Rutgers-New Jersey Medical School (N.E.G.), Newark, New Jersey; Regions Hospital (D.J.D.), St. Paul, Minnesota; University of California (T.W.C.), San Diego, California; and University of Michigan, Ann Arbor (LMN), Michigan.

Address for correspondence: Lena M. Napolitano MD, FACS, FCCP, MCCM, University of Michigan, Ann Arbor, MI; email: lenan@umich.edu.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

patients. Providing ARDS patients with optimal, evidence-based care improves outcomes. Protocolized management ensures that ICU teams at the bedside have guidance regarding what constitutes the best evidence-based care.¹⁴

PROTOCOL RATIONALE AND GOALS

The goals of this protocol are to ensure that: 1) ARDS and severe hypoxemia are recognized in the management of the trauma patient and 2) treated with evidence-based strategies to improve outcomes. Importantly, there are no randomized controlled trials specific to the trauma population for ARDS treatment strategies, and posttraumatic ARDS represented only 8% to 13% of the study cohorts enrolled in existing trials. Thus, these treatment recommendations are mostly extrapolated from the results of randomized controlled trials in medical and nontrauma surgical patients.

While there are many published algorithms and protocols for the management of ARDS and severe hypoxemia, the lack of trauma-specific trials leaves questions as to how to apply them in trauma patients. Given the gaps in the data, there is a need for a consensus clinical ARDS algorithm and protocol that addresses the specific needs of trauma patients for implementation across all trauma centers with a brief evidence summary to guide management. Stakeholders from the American Association for the Surgery of Trauma and the American College of Surgeons—Committee on Trauma established a work group to create a clinical protocol. The work group conducted a literature review to identify prospective and retrospective studies related to ARDS and severe hypoxemia in trauma patients. These studies were reviewed by members of the group, and consensus guidelines were generated based on current literature and expert opinion.

The clinical protocol and evidence-based algorithm (Fig. 1) presented here is based on best available evidence from national and international ARDS guidelines (Table 2) and the consensus of experts on this panel. However, treatment decisions regarding ARDS management in trauma patients should be individualized for each patient and do not exclude other treatment strategies as being within the standard of care. Ultimately, the responsibility to implement treatment decisions rest with the treating physician at bedside in the intensive care unit.

EVIDENCE BASE: BRIEF SUMMARY

Diagnosis

Diagnosis of ARDS (Table 1) is delayed or missed in many patients. In the LUNG-SAFE study (largest international ARDS



Figure 1. Algorithm of treatment strategies for patients with ARDS.¹⁵

TARIF 2	National	and	International		Guidelines
IADLL Z.	INALIONAL	anu	international	AIVD 3	Guidennes

Organization	Guidelines	Reference
Scandinavian Society of Anaesthesiology and Intensive Care Medicine	Scandinavian clinical practice guideline on mechanical ventilation in adults with the acute respiratory distress syndrome 2015	Claesson J, Freundlich M, Gunnarsson I, Laake JH, Vandvik PO, Varpula T, Aasmundstad TA; Scandinavian Society of Anaesthesiology and Intensive Care Medicine. Scandinavian clinical practice guideline on mechanical ventilation in adults with the acute respiratory distress syndrome. Acta Anaesthesiol Scand. 2015 Mar;59(3):286–97. doi: 10.1111/aas.12449 PMID: 25524779. https://onlinelibrary.wiley.com/doi/epdf/10.1111/aas.12449
American Thoracic Society European Society of Intensive Care Medicine Society of Critical Care Medicine	An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice Guideline: Mechanical Ventilation in Adult Patients with Acute Respiratory Distress Syndrome. 2017	 Fan E, Del Sorbo L, Goligher EC et al. An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice Guideline: Mechanical Ventilation in Adult Patients with Acute Respiratory Distress Syndrome. Am J Respir Crit Care Med Vol 2017;195(9): 1253–1263 https://pubmed.ncbi.nlm.nih.gov/28459336/
Faculty of Intensive Care Medicine	Formal guidelines: Management of acute respiratory distress syndrome. 2019	Papazian L, Aubron C, Brochard L, et al.
Intensive Care Society		Formal guidelines: Management of acute respiratory Distress syndrome Ann Intensive Care 2019;9:69 https://pubmed.ncbi.nlm.nih.gov/31197492/
Japanese Society of Intensive Care Medicine	ARDS Clinical Practice Guideline 2021	Tasaka S, Ohshimo S, Takeuchi M et al.
Japanese Respiratory Society		ARDS Clinical Practice Guideline 2021.
Japanese Society of Respiratory Care Medicine		J Intensive Care 10, 32 (2022). https://doi.org/10.1186/s40560-022-00615-6

study), ARDS diagnosis was missed in 40% and delayed/missed in two thirds of patients. Early diagnosis of ARDS facilitates early appropriate treatment including protective lung ventilation and other treatment strategies, which may improve survival.^{4,16}

Treatment

Treatment strategies for ARDS include the three broad categories of mechanical ventilation, pharmacologic treatments, and adjuncts. We provide a brief discussion of the evidence regarding the treatment strategies in each of these categories, with review of appropriate indication and whether the treatment has an impact on mortality or oxygenation (Table 3).

Mechanical Ventilation

Low Tidal Volume, Low Plateau Pressure

Protective lung ventilation using low tidal volume (LTV) (4–8 mL/kg predicted body weight) and low plateau pressure ($<30 \text{ cm H}_2\text{O}$) is the current standard of care for ARDS management, associated with decreased mortality and prevention of ventilator-induced lung injury.^{17,18} A recent systematic review and meta-analysis of 7 RCTs with 1481 patients concluded "The trend toward lower mortality with LTV ventilation in the primary analysis and the significant relationship between the degree of tidal volume reduction and the mortality effect together suggest, but do not prove, that LTV ventilation improves mortality among critically ill adults with ARDS." Importantly, high PEEP co-intervention with LTV showed a greater mortality benefit (relative risk, 0.58; 95% CI, 0.41–0.82) than LTV alone.¹⁹

In trauma patients with ARDS, LTV ventilation will likely result in hypercapnia, and may have potential adverse effects in patients with traumatic brain injury. In patients with both ARDS

© 2023 The Author(s). Published by Wolters Kluwer Health, Inc.

and traumatic brain injury, optimal $PaCO_2$ targets with LTV ventilation should be adjusted to maintain adequate cerebral perfusion pressure and cerebral oxygen delivery using a personalized approach based on intracranial pressure and multimodal brain and cerebral monitoring.^{20–22}

Optimal and High Versus Low PEEP

Current evidence supports that patients with moderate or severe ARDS ($PaO_2/FIO_2 \le 200$) should receive higher levels of PEEP, based on a meta-analysis of individual patient data from 3 large RCTs of higher vs. lower PEEP.^{23,24} Patients with moderate/severe ARDS randomized to higher PEEP had significantly lower mortality (adjusted RR, 0.90; 95% CI, 0.81-1.00) and better oxygenation (higher P/F ratio). In contrast, higher PEEP had no significant effect in the patients with mild ARDS (adjusted RR, 1.29; 95% CI, 0.91–1.83; p = 0.02 vs. moderate/ severe ARDS group).^{25,26} These data support the use of the "HIGH" PEEP Table in moderate/severe ARDS patients in whom it improves oxygenation without reducing pulmonary compliance or worsening hemodynamic status. A recent Bayesian network meta-analysis of 4646 moderate-severe ARDS patients confirmed that the use of higher PEEP was superior to that of lower PEEP for 28-day mortality, and use of a brief recruitment maneuver (<60 seconds) was also associated with a mortality benefit as compared with a lower PEEP strategy.²⁷

The use of "optimal PEEP," adjusted to each patient's potential for lung recruitment, has been advocated to achieve maximum lung static compliance, reduce atelectasis, avoid hyperinflation, and improved oxygenation.^{28,29} Studies are now investigating personalized PEEP levels to optimize lung recruitment

Treatment	Summary	Mortality Benefit
Mechanical ventilation		
Lung protective tidal volume ventilation	Recommend for all ARDS patients Target 4–8 mL/kg predicted body weight (PBW) and low plateau pressure ($<30 \text{ cm H}_2\text{O}$)	Yes
High PEEP in moderate/severe ARDS	Recommend for patients with moderate/severe ARDS Possible contraindication in TBI and shock states, individual PEEP titration in these states	Yes
Lung recruitment maneuvers	Consider in patients with moderate/severe (PaO ₂ /FIO ₂ \leq 200) ARDS to improve oxygenation	Uncertain
Esophageal pressure-guided therapy	Current data do not support the use of esophageal pressure-guided PEEP titration in routine ARDS management, but may be effective as a rescue therapy in obese patients and those with stiff chest walls	No
Adjuncts		
Prone positioning (16 hours/d)	Recommend for patients with moderate/severe ARDS Maintained daily until the PaO ₂ /FIO ₂ ratio remains >150 mm Hg, with PEEP ≤ 10 cm H ₂ O, FIO ₂ $\leq 60\%$ for at least 4 hours after the end of the most recent proning session	Yes
VV-ECMO	Consider for refractory hypoxemia despite maximal therapy with mechanical ventilation, pharmacologic adjuncts, and prone position, in patients without contraindications	Yes
Pharmacologic treatment		
Diuretics for conservative fluid strategy	Consider in ARDS patients without contraindications.	No
Neuromuscular blockade	Consider in ARDS patients with severe refractory hypoxemia and ventilator dysynchrony that cannot be managed with sedation alone	No
Inhaled pulmonary vasodilators	Consider as rescue strategy in severe hypoxemia/ARDS with P/F ratio <100 when other strategies discussed above have not resulted in improved oxygenation, and before consideration of VV-ECMO	No
Steroids	Recommended for patients with COVID-related ARDS	Yes
	Consider in patient with non-COVID-related ARDS	Equivocal
	No clear data for trauma-related ARDS	No

TABLE 3.	Current Manac	ement Strategies	s of ARDS and	Associated Mort	ality Benefit
	0 411 0110 1100				

selected by electrical impedance tomography, transpulmonary pressure with esophageal manometry, and the recently described recruitment-to-inflation ratio.^{30–32}

In trauma patients with traumatic brain injury and ARDS, high PEEP may be detrimental as it can cause increased intracranial pressure by impaired jugular venous outflow, impeding cerebral venous return to the right atrium, decreased mean arterial pressure, and result in decreased cerebral perfusion pressure.³³ Similarly, high PEEP may be contraindicated in those with hemorrhagic shock requiring ongoing blood product resuscitation or with distributive/neurogenic/septic shock due to vasodilation and hypotension.^{34–36} Therefore, PEEP should be determined individually in trauma patients with ARDS, reviewing the risk/benefit ratio of the potential positive impact of higher PEEP in severe hypoxemia vs. the negative impact of higher PEEP on hemodynamics and intracranial pressure in specific injury patterns. For instance, in trauma patients with severe bilateral pulmonary contusions, higher PEEP may be beneficial, but in those with focal areas of pulmonary contusion PEEP can be detrimental.^{31,37–39}

Driving Pressure

Driving pressure is calculated as the difference between plateau pressure and PEEP. A secondary analysis of data from 3562 ARDS patients from nine randomized trials confirmed that reducing driving pressure was strongly associated with increased survival.⁴⁰ A recent, large registry analysis of over 13,000 patients receiving mechanical ventilation in nine ICUs found that increased driving pressure, or mechanical power, was associated with increased mortality and the association was stronger with worse hypoxemia.⁴¹ Increased driving pressure is an indicator of severity of the ARDS, is related to tidal volume in volume control modes, and is associated with more complications and increased mortality. Based on these associations, some authors recommend targeting a driving pressure below 14 cm H_2O .^{42–44} However, at present, there is no randomized trial evidence that a ventilator strategy focused on limiting driving pressure reduces mortality in ARDS.⁴⁵ Interestingly, driving pressure was not predictive of ARDS outcome (mortality or mechanical ventilation duration) in severe trauma patients with chest injuries and ARDS.⁴⁶

Lung Recruitment Maneuver

Lung recruitment maneuvers (LRMs) involve transient increase in airway pressures to open/recruit collapsed lung.⁴⁷ Lung recruitment maneuver strategies include sustained inflation, intermittent sighs, stepwise increase of PEEP or airway pressure and high driving pressure.^{48,49} Lung recruitment maneuvers are commonly associated with improved oxygenation, reduced intrapulmonary shunt and increased pulmonary compliance, but may also cause hemodynamic compromise and/or barotrauma.⁵⁰ The optimal LRM method is not clear.

Current guidelines recommend that patients with moderate or severe ARDS (PaO₂/FIO₂ \leq 200) should be considered for LRM to improve oxygenation. A systematic review and meta-analysis of 5 RCTs suggested that LRMs in combination with higher PEEP was associated with reduced mortality.⁵¹ A more recent systematic review including 14 RCTs reported that LRMs were not associated with reduced 28-day mortality (RR, 0.92; 95% CI, 0.82-1.04, p = 0.21), but did improve oxygenation (PaO₂/FIO₂ ratio was significantly higher (mean difference, 47.6 mm Hg; 95% CI, 33.4–61.8; p < 0.001). Lung recruitment maneuvers were associated with a decreased rate of rescue therapy (RR, 0.69; 95% CI, 0.56-0.84; p < 0.001), and an increased rate of hemodynamic compromise (RR, 1.19; 95% CI, 1.06–1.33; p = 0.002), compared with no-LRM group.⁵² Another review of 10 RCTs showed a significant benefit of LRMs for decreased length of hospital stay (mean difference [MD], -1.75; 95% CI, -3.40 to -0.09; p = 0.04; p for

heterogeneity = 0.3; I2 = 18%) and improved PaO₂/FIO₂ ratio on the third day (MD, 52.72; 95% CI, 18.77–86.67; p = 0.002).⁵³

Given the association of LRM with hemodynamic compromise, caution should be used in trauma patients with ARDS. Lung recruitment maneuver may be contraindicated in patients with traumatic brain injury and shock states, just as in the discussion regarding high PEEP strategies, a personalized approach to LRM is required in these patients.

Esophageal Pressure-Guided Therapy

Transpulmonary pressure is the pressure distending the lung and is measured via esophageal manometry. In the EPVent-2 Study (n = 200), among patients with moderate to severe ARDS, esophageal pressure-guided PEEP, compared with empirical high PEEP-FIO₂, resulted in no significant difference in death and ventilator free days.54 These findings do not support the use of esophageal pressure-guided PEEP titration in routine ARDS management. Esophageal manometry-guided PEEP titration can be helpful in severe hypoxemia patients as a rescue strategy, particularly in obese patients and those with stiff chest walls. A recent single-institution report in obese ARDS patients confirmed that use of an individualized protocol by a lung rescue team using LRM, esophageal manometry, and hemodynamic monitoring was associated with an almost 50% reduction in mortality at 28 days, 3 and 6 months, and 1 year.⁵⁵ A post hoc analysis of the EPVent-2 trial reported that that PEEP titrated to end-expiratory transpulmonary pressure closer to 0 cm H₂O was associated with greater survival.⁵⁶ To our knowledge, a post hoc secondary analysis of obese patients in the EPVent-2 trial has not been reported, and a future randomized trial in obese patients is warranted. In a recent expert review, the authors concluded: "Transpulmonary pressure represents a physiologically sound safety limit for mechanical ventilation that should be measured and targeted at least in the most severe ARDS patients. Targeting transpulmonary pressure means 'personalizing' the ventilatory settings."57

Airway Pressure Release Ventilation

In two recent systematic review/meta-analyses of RCTs, airway pressure release ventilation (APRV) was associated with improved oxygenation, mortality benefit, and increased number of ventilator-free days when compared with conventional ventilation, with no increased risk of barotrauma or worse hemodynamics.^{58,59} These data are limited, however, to inclusion of five RCTs with small sample size (n = 330), consistent with low quality evidence and moderate heterogeneity. A more recent review of seven RCTs with 405 patients reported similar findings.⁶⁰ A recent adaptation of APRV is time-controlled adaptive ventilation (TCAV) which was developed to minimize dynamic alveolar strain, and gradually "nudge" alveoli open and prevent alveolar collapse using a simple strategy of open-valve CPAP with brief, intermittent releases guided by changes in lung mechanics. This personalized and adaptive approach to mechanical ventilation configures each breath guided by the previous one, but still requires confirmation of efficacy in clinical trials.^{61,62} Studies have confirmed that APRV/TCAV is safe in severe TBI and does not increase intracranial pressure.⁶³

High-Frequency Oscillatory Ventilation

Two large multicenter RCTs confirmed no benefit of high-frequency oscillatory ventilation (HFOV), including The

Oscillation in ARDS trial done in United Kingdom⁶⁴ or possible harm (higher mortality: RR, 1.41; 95% CI, 1.21–1.79) in the US Oscillation for Acute Respiratory Distress Syndrome Treated Early trial.⁶⁵ High-frequency oscillatory ventilation should not routinely be used in patients with moderate or severe ARDS. The potential use of HFOV as a rescue strategy for severe hypoxemia has been successful in some studies and warrants further investigation.

PHARMACOLOGIC TREATMENT

Conservative Fluid Management

Hypervolemia can cause pulmonary edema with resultant hypoxemia. Randomized trials comparing liberal to conservative fluid management strategies in ARDS have confirmed better outcomes with conservative fluid management, including shortened duration of mechanical ventilation, and decreased ICU and hospital length of stay.^{66–68} The Fluids and Catheters Treatment Trial (FACTT) confirmed that a conservative fluid management strategy for ARDS reduced net fluid balance and improved oxygenation and ventilator-free days.⁶⁷ A subsequent study with less strict fluid protocol (FACTT Lite) also confirmed improved outcomes.⁶⁹ Additional studies confirm that early diuretic use and conservative fluid management are associated with decreased mortality in ARDS.^{70–72} Diuresis should be considered in ARDS patients without contraindications.

Neuromuscular Blockade

As with other rescue strategies for severe hypoxemia, neuromuscular blockade is associated with improved oxygenation, but no improvement in survival based on the most recent systematic review and meta-analysis of 5 RCTs.⁷³ The French multicenter ACURASYS trial over a decade ago reported significantly lower mortality with a 48-hour infusion of cisatracurium in addition to deep sedation.⁷⁴ The multicenter Reevaluation of Systemic Early Neuromuscular Blockade (ROSE) trial was stopped early for futility after enrollment of 1006 moderate/severe ARDS patients and reported no difference in 90-day mortality (42.5% vs. 42.8%).¹⁵ Based on these results, neuromuscular blockade can be considered in ARDS patients with severe refractory hypoxemia or ventilator dyssynchrony that cannot be managed with sedation alone (Fig. 2). Neuromuscular blockade is not required for prone positioning.

Inhaled Pulmonary Vasodilators (iNO, Epoprostenol, Etc)

Inhaled nitric oxide (iNO) can be considered as a rescue strategy in cases of severe hypoxemia/ARDS with P/F ratio <100 when other strategies discussed above have not resulted in improved oxygenation, and before consideration of veno-venous extracorporeal membrane oxygenation (VV-ECMO). Systematic reviews of all RCTs documented significantly improved oxygenation with iNO treatment, but no mortality benefit.^{75–77} We do not recommend the *routine* use of iNO in ARDS given no mortality benefit, high expense and potential risk for AKI, but it can be considered as a rescue strategy for patients with severe hypoxemia (Table 3). In a report examining ARDS management over a 12-year period of two randomized clinical trials, iNO use increased (24.9–65.8%) with no change in prone positioning



Figure 2. Algorithm for use of NMBA in ARDS. From: Alhazzani W, Belley-Cote E, Møller MH, et al. Neuromuscular blockade in patients with ARDS: a rapid practice guideline. *Intensive Care Med.* 2020;46(11):1977–1986. doi:10.1007/s00134-020-06227-8. NMBA, neuromuscular blocking agents.

(16.2–18.9%).⁷⁸ The increasing use of iNO in severe hypoxemia patients may result in significantly improved oxygenation, which can provide time for treatment of underlying factors contributing to ARDS.

Steroids for ARDS

The DEXA-ARDS multicenter randomized trial (n = 277) documented that dexamethasone treatment (20 mg IV daily for 5 days, reduced to 10 mg once daily from Day 6 to Day 10) was significantly associated with increased ventilator-free days at 28 days (12.3 days vs. 7.5 days, p < 0.0001) and an absolute 15% (36% vs. 21%) reduction in all-cause mortality at 60 days in moderate-to-severe ARDS enrolled 24 hours after onset.⁷⁹ However only 21 trauma patients were included in the DEXA-ARDS study, significantly limiting the generalizability of these conclusions regarding steroids for use in trauma ARDS.

A meta-analysis of 18 RCTs (n = 2826) confirmed that the use of corticosteroids reduced mortality in patients with ARDS of any etiology (RR, 0.82; 95% CI, 0.72–0.95; ARR, 8.0%; 95% CI, 2.2–12.5%, moderate certainty). Patients who received a longer course of corticosteroids (over 7 days) had higher rates of survival compared with a shorter course. This effect was consistent for all ARDS patients, and all steroid types and dosages studied.⁸⁰ But many of these studies did not include trauma-related ARDS patients.

Systemic corticosteroids are recommended for patients with COVID-related ARDS.^{81,82}

ADJUNCTS TO MECHANICAL VENTILATION

Prone Position

The PROSEVA multicenter trial (27 ICUs) confirmed a significant mortality reduction in ARDS patients enrolled after a 12-hour to 24-hour stabilization period with a PaO_2/FIO_2 ratio <150 mm Hg, with PEEP at least 5 cm H_2O , FIO_2 at least 60%,

and tidal volume 6 mL/kg PBW.⁸³ Patients had, on average, four prone sessions of at least 16 hours duration. Prone position was continued even in the absence of improved oxygenation. This trial validated the results of a previous patient-level meta-analysis from 4 earlier randomized trials which demonstrated significant mortality reduction with prone position in severe ARDS patients.⁸⁴ Prone position should be considered in all appropriate patients and maintained daily until the PaO₂/FIO₂ ratio remains >150 mm Hg, with PEEP ≤10 cm H₂O, FIO₂ ≤ 60% for at least 4 hours after the end of the most recent proning session. Importantly, ICU-specific prone position protocols and appropriate ICU staff training are required for patient safety.

In trauma patients with severe ARDS, specific considerations which require discussion prior to prone position therapy include need for spine stabilization, traumatic brain injury management, pelvic fracture external fixation, and open abdomen. In most cases, these are not absolute contraindications to prone positioning for treatment of severe ARDS given the high ARDS-associated mortality rate.

Veno-venous Extracorporeal Carbon Dioxide Removal

Low tidal volume is associated with hypercapnia and respiratory acidosis. If the hypercapnia becomes severe, the resulting acute respiratory acidosis may cause adverse effects, including pulmonary hypertension and altered cardiac function. Extracorporeal carbon dioxide removal (ECCO₂R) provides CO₂ clearance using a small dual-lumen central venous catheter and a pump at low speed with systemic anticoagulation. Previous studies have demonstrated the feasibility of ECCO₂R to facilitate LTV by limiting hypercapnia and the resulting respiratory acidosis. The REST trial⁸⁵ evaluated if use of ECCO₂R in adults receiving mechanical ventilation for acute hypoxemic respiratory failure (n = 405, P/F < 150) to further reduce LTV (≤3 mL/kg PBW) compared with conventional LTV (6 ml/kg PBW) would improve 90-day all-cause mortality. The study was stopped due to futility; 90-day mortality 41.5% ECCO₂R versus 39.5% standard care. ECCO₂R was associated with increased serious adverse events including intracranial hemorrhage, and higher need for neuromuscular blockade. The prior Xtravent study used the same trial design as the REST trial and showed decreased ventilator days and no hospital mortality difference, but the mortality rate was low at 16.5%.⁸⁶ At present, VV-ECCO₂R is not indicated for use in ARDS patients.

Veno-venous Extracorporeal Membrane Oxygenation

Veno-venous extracorporeal membrane oxygenation should be considered in severe ARDS patients if the severe hypoxemia rescue strategies described above have failed to improve oxygenation and the patient is a potential ECMO candidate with potentially reversible respiratory failure (Fig. 3).

Two randomized trials examined the efficacy of VV-ECMO in adults with severe ARDS. The CESAR trial (n = 180) reported an ECMO survival benefit in the intention-to-treat analysis (RR, 0.69; 95% CI, 0.05–0.97, p = 0.03), but the study conclusion was limited since 25% of subjects assigned to the ECMO group never received ECMO therapy.⁸⁸ The subsequent EOLIA trial randomized 249 patients with severe ARDS to early ECMO (immediate VV-ECMO if P/F < 50 for >3 hours; P/F < 80 for >6 hours; or arterial blood pH < 7.25 with paCO₂ 60 for >6 hours) versus protocolized mechanical ventilation, and 60-day mortality was not statistically different (RR, 0.76; 95% CI, 0.55 to 1.04; 35% vs. 46%; p = 0.09). The risk of 90-day treatment failure (death in ECMO cohort, death or crossover to ECMO in control cohort) was significantly higher in the control group. A high percentage of control patients required "rescue" ECMO for refractory hypoxemia (n = 35, 28%) with a higher 60-day mortality (57%) and 7 patients required VA-ECMO.⁸⁹ A post hoc Bayesian analysis of EOLIA data reported a high likelihood of survival benefit with ECMO.⁹⁰

In a systematic review and individual patient data meta-analysis of the two ECMO-ARDS RCTs (CESAR and EOLIA) with 429 patients, 90-day mortality was significantly lower with ECMO compared with conventional management (RR 0.75, 95% confidence interval (CI) 0.6–0.94; p = 0.013; $I^2 = 0\%$). The RR of 90-day treatment failure (death in ECMO cohort, death or crossover to ECMO in control cohort) was 0.65 (95% CI, 0.52–0.8; $I^2 = 0\%$).⁹¹ A systematic review and meta-analysis of 5 studies (2 RCTs and 3 observational studies with matching techniques, n = 773) reported lower 60-day mortality in the VV-ECMO group (RR, 0.73; 95% CI, 0.58–0.92; p = 0.008, $I^2 0\%$), but higher risk (19%) of major hemorrhage.⁹²

A Trauma Quality Improvement Program database study (2013–2016) compared trauma patients with ARDS who received ECMO (97) to a propensity-matched cohort of patients who underwent conventional management. Extracorporeal membrane oxygenation was associated with significant lower overall hospital mortality (23 vs. 50%, p < 0.001) but higher rates of complications and prolonged hospital length of stay.⁹³

In centers without ECMO capability, planning for early transfer to an ECMO-capable center should be discussed.



Figure 3. Algorithm for ECMO in acute respiratory distress syndrome. From: Abrams D, Ferguson N, Brochard L, et al. ECMO for ARDS: from salvage to standard of care? *Lancet Respir Med.* 2019 Feb;7(2):108–110. doi:10.1016/S2213-2600(18)30506-X. Relative contraindications to VV-ECMO for ARDS include the following⁸⁷: central nervous system hemorrhage; significant central nervous system injury; irreversible and incapacitating central nervous system or pulmonary pathology; systemic bleeding, diffuse alveolar hemorrhage; contraindications to anticoagulation (VV-ECMO can be managed with low-dose or without continuous systemic anticoagulation if flow rates are high, but anticoagulation bolus is strongly recommended with ECMO cannulae placement); inability to receive blood product transfusion; immunosuppression; older age (increasing risk of death with increasing age, but no age threshold is established); mechanical ventilation for more than 7 days with Pplat > 30cm H2O and FiO2 > 90%; obesity with a BMI greater than 40 kg/m2 (This is a historical relative contraindication due to difficulty in ECMO cannulation and achieving adequate ECMO circuit flow, as well as failure to wean. Increasing data are emerging supporting the use of ECMO in the obese population).

Common adult ECMO transfer criteria include P/F ratio $\leq 100 \text{ mmHg}$, no contraindications to ECMO, single-system organ failure, and intubation $\leq 7 \text{ days.}^{87}$

FUTURE RESEARCH

As we have reviewed above, most studies regarding the specific efficacy of treatment strategies in adult ARDS patients are in medical and surgical nontrauma patients. Future ARDS studies should aim to include a sub-population of trauma-related ARDS patients to aim to provide better evidence for effective treatments in trauma-related ARDS.

Precision Medicine and ARDS

We recognize that ARDS is a clinically heterogeneous syndrome, and that different patient phenotypes exist which may have quite variable responses to specific ARDS treatment strategies.^{94–96} In the future, precision medicine may be able to incorporate variability in specific factors and phenotypes to better individualize appropriate therapies for to improve care for trauma patients with ARDS.⁹⁷ Even with current limitations, it is important to tailor aspects of ARDS management such as LRM and PEEP to the specific needs of TBI and hemorrhaging patients.

Limitations

While an extensive literature review was conducted and current studies were evaluated and discussed by workgroup members, a formal evaluation of the level of evidence reviewed and the strength of recommendations provided are not included as part of this clinical protocol. We have provided references to formal ARDS guidelines to provide this information (Table 2).

CONCLUSION

Acute respiratory distress syndrome is life-threatening in trauma patients, and clinicians must have knowledge of appropriate evidence-based strategies for optimal management. This clinical trauma protocol outlines effective ARDS treatment strategies with specific reference to their use and potential impact in trauma patients with ARDS. Acute respiratory distress syndrome is a clinically heterogeneous syndrome, and recent studies have reported different phenotypes which have variable clinical responses to specific ARDS treatments. Randomized treatment trials in trauma patients with ARDS are warranted to further determine optimal treatment strategies in this patient cohort.

AUTHORSHIP

All authors participated in the literature review, development of the clinical protocol and algorithm, article writing and critical review of all article revisions.

DISCLOSURE

The authors declare no funding or conflicts of interest.

REFERENCES

- ARDS Definition Task Force, Ranieri VM, Rubenfeld GD, et al. Acute respiratory distress syndrome: the Berlin definition. JAMA. 2012;307:2526–2533.
- Fan E, Brodie D, Slutsky AS. Acute respiratory distress syndrome: advances in diagnosis and treatment. JAMA. 2018;319(7):698–710.
- Ashbaugh DG, Bigelow DB, Petty TL, Levine BE. Acute respiratory distress in adults. *Lancet*. 1967;2:319–323.

- Bellani G, Laffey JG, Pham T, Fan E, Brochard L, Esteban A, et al. Epidemiology, patterns of care, and mortality for patients with acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA*. 2016;315:788–800.
 - Qadir N, Bartz RR, Cooter ML, Hough CL, Lanspa ML, Banner-Goodspeed VM, et al, Severe ARDS: Generating Evidence (SAGE) Study Investigators; Society of Critical Care Medicine's Discovery Network. Variation in early management practices in moderate-to-severe ARDS in the United States: the severe ARDS: generating evidence study. *Chest.* 2021;160(4):1304–1315.
 - Pfeifer R, Heussen N, Michalewicz E, Hilgers RD, Pape HC. Incidence of adult respiratory distress syndrome in trauma patients. *J Trauma Acute Care Surg.* 2017;83(3):496–506.
 - Ramin S, Charbit J, Jaber S, Capdevila X. Acute respiratory distress syndrome after chest trauma: epidemiology, specific physiopathology and ventilation strategies. *Anaesth Crit Care Pain Med.* 2019;38(3):265–276.
 - Fan TH, Huang M, Gedansky A, Price C, Robba C, Hernandez AV, et al. Prevalence and outcome of acute respiratory distress syndrome in traumatic brain injury: a systematic review and meta-analysis. *Lung.* 2021;199(6):603–610.
 - Rincon F, Ghosh S, Dey S, Maltenfort M, Vibbert M, Urtecho J, et al. Impact of acute lung injury and acute respiratory distress syndrome after traumatic brain injury in the United States. *Neurosurgery*. 2012;71(4):795–803.
 - Tran A, Fernando SM, Brochard LJ, Fan E, Inaba K, Ferguson ND, et al. Prognostic factors for development of acute respiratory distress syndrome following traumatic injury: a systematic review and meta-analysis. *Eur Respir J.* 2022;59(4):2100857.
 - Humayun M, Premraj L, Shah V, Cho SM. Mechanical ventilation in acute brain injury patients with acute respiratory distress syndrome. Review. Front Med (Lausanne) 2022;9:999885.
 - Tignanelli CJ, Hemmila MR, Rogers MAM, Raghavendran K. Nationwide cohort study of independent risk factors for acute respiratory distress syndrome after trauma. *Trauma Surg Acute Care Open.* 2019;4(1):e000249.
 - Kasotakis G, Stanfield B, Haines K, Vatsaas C, Alger A, Vaslef SN, et al. Acute respiratory distress syndrome (ARDS) after trauma: improving incidence, but increasing mortality. *J Crit Care*. 2021;64:213–218.
 - Menk M, Estenssoro E, Sahetya SK, Neto AS, Sinha P, Slutsky AS, et al. Current and evolving standards of care for patients with ARDS. *Intensive Care Med.* 2020;46(12):2157–2167.
 - The National Heart, Lung, and Blood Institute PETAL Clinical Trials Network. Early neuromuscular blockade in the acute respiratory distress syndrome. N Engl J Med. 2019;380:1997–2008.
 - Bellani G, Pham T, Laffey JG. Missed or delayed diagnosis of ARDS: a common and serious problem. *Intensive Care Med.* 2020;46:1180–1183.
 - Liaqat A, Mason M, Foster BJ, Kulkarni S, Barlas A, Farooq AM, et al. Evidence-based mechanical ventilatory strategies in ARDS. *J Clin Med.* 2022;11(2):319.
 - Acute Respiratory Distress Syndrome Network, Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, Wheeler A. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med.* 2000; 342(18):1301–1308.
 - Walkey AJ, Goligher EC, Del Sorbo L, Hodgson CL, Adhikari NKJ, Wunsch H, et al. Low tidal volume versus non-volume-limited strategies for patients with acute respiratory distress syndrome. A systematic review and metaanalysis. *Ann Am Thorac Soc.* 2017;14(Supplement_4):S271–S279.
 - Kim JA, Wahlster S, LaBuzetta JN, Nobleza COS, Johnson NJ, Rubinos C, et al. Focused management of patients with severe acute brain injury and ARDS. *Chest.* 2022;161(1):140–151.
 - Chacon-Aponte AA, Duran-Vargas EA, Arevalo-Carrillo JA, Lozada-Martinez ID, Bolano-Romero MP, Moscote-Salazar LR, et al. Brain-lung interaction: a vicious cycle in traumatic brain injury. *Acute Crit Care*. 2022;37(1):35–44.
 - Taran S, Wahlster S, Robba C. Ventilatory targets following brain injury. *Curr Opin Crit Care*. 2023;29(2):41–49.
 - Briel M, Meade M, Mercat A, Brower RG, Talmor D, Walter SD, et al. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome: systematic review and metaanalysis. *JAMA*. 2010;303:865–873.
 - Guo L, Xie J, Huang Y, Pan C, Yang Y, Qiu H, et al. Higher PEEP improves outcomes in ARDS patients with clinically objective positive oxygenation response to PEEP: a systematic review and meta-analysis. *BMC Anesthesiol*. 2018;18(1):172.
 - 25. Fan E, Del Sorbo L, Goligher EC, Hodgson CL, Munshi L, Walkey AJ, et al. An official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine clinical practice guideline:

Fawley et al.

mechanical ventilation in adult patients with acute respiratory distress syndrome. Am J Respir Crit Care Med. 2017;195(9):1253–1263.

- Walkey AJ, Del Sorbo L, Hodgson CL, Adhikari NKJ, Wunsch H, O'Meade MO, et al. Higher PEEP vs. lower PEEP strategies for patients with acute respiratory distress syndrome: a systematic review and meta-analysis. *Ann Am Thorac Soc.* 2017;14(Supplement 4):S297–S303.
- Dianti J, Tisminetzky M, Ferreyro BL, Englesakis M, Del Sorbo L, Sud S, et al. Association of positive end-expiratory pressure and lung recruitment selection strategies with mortality in acute respiratory distress syndrome: a systematic review and network meta-analysis. *Am J Respir Crit Care Med.* 2022;205:1300–1310.
- Pelosi P, Ball L, Barbas CSV, Bellomo R, Burns KEA, Einav S, et al. Personalized mechanical ventilation in acute respiratory distress syndrome. *Crit Care*. 2021;25(1):250.
- Wendel-Garcia PD, Roche-Campo F, Buehler PK. High positive end-expiratory pressure and lung recruitment in moderate to severe acute respiratory distress syndrome: does one size really fit all? *Am J Respir Crit Care Med.* 2022; 205(11):1260–1262.
- Scaramuzzo G, Spadaro S, Dalla Corte F, Waldmann AD, Böhm SH, Ragazzi R, et al. Personalized positive end-expiratory pressure in acute respiratory distress syndrome: comparison between optimal distribution of regional ventilation and positive transpulmonary pressure. *Crit Care Med.* 2020;48(8):1148–1156.
- Mauri T. Personalized positive end-expiratory pressure and tidal volume in acute respiratory distress syndrome: bedside physiology-based approach. *Crit Care Explor.* 2021;3(7):e0486.
- 32. Chen L, Del Sorbo L, Grieco DL, Junhasavasdikul D, Rittayamai N, Soliman I, et al. Potential for lung recruitment estimated by the recruitment-to-inflation ratio in acute respiratory distress syndrome: a clinical trial. Am J Respir Crit Care Med. 2020;201:178–187.
- 33. Della Torre V, Badenes R, Corradi F, Racca F, Lavinio A, Matta B, et al. Acute respiratory distress syndrome in traumatic brain injury: how do we manage it? *J Thorac Dis.* 2017;9(12):5368–5381.
- Coruh B, Luks AM. Positive end-expiratory pressure: when more may not be better. Ann Am Thorac Soc. 2014;11(8):1327–1331.
- Fougères E, Teboul JL, Richard C, Osman D, Chemla D, Monnet X. Hemodynamic impact of a positive end-expiratory pressure setting in acute respiratory distress syndrome: importance of the volume status. *Crit Care Med.* 2010;38:802–807.
- Herff H, Paal P, von Goedecke A, Lindner KH, Severing AC, Wenzel V. Influence of ventilation strategies on survival in severe controlled hemorrhagic shock. *Crit Care Med.* 2008;36(9):2613–2620.
- Santa Cruz R, Villarejo F, Irrazabal C, Ciapponi A. High versus low positive end-expiratory pressure (PEEP) levels for mechanically ventilated adult patients with acute lung injury and acute respiratory distress syndrome. *Cochrane Database Syst Rev.* 2021;3(3):CD009098.
- 38. Constantin JM, Jabaudon M, Lefrant JY, Jaber S, Quenot JP, Langeron O, et al. For the AZUREA network. Personalised mechanical ventilation tailored to lung morphology versus low positive end-expiratory pressure for patients with acute respiratory distress syndrome in France (the LIVE study): a multicenter, single-blind, randomised controlled trial. *Lancet Respir Med*. 2019;7(10):870–880.
- Dhar SM, Breite MD, Barnes SL, Quick JA. Pulmonary contusion in mechanically ventilated subjects after severe trauma. *Respir Care*. 2018;63(8):950–954.
- Amato MBP, Meade MO, Slutsky AS, Brochard L, Costa ELV, Schoenfeld DA, et al. Driving pressure and survival in the acute respiratory distress syndrome. N Engl J Med. 2015;372:747–755.
- Urner M, Jüni P, Hansen B, Wettstein MS, Ferguson ND, Fan E. Time-varying intensity of mechanical ventilation and mortality in patients with acute respiratory failure: a registry-based, prospective cohort study. *Lancet Respir Med.* 2020;8(9):905–913.
- Aoyama H, Pettenuzzo T, Aoyama K, Pinto R, Englesakis M, Fan E. Association of driving pressure with mortality among ventilated patients with acute respiratory distress syndrome. *Crit Care Med.* 2018;46(2):300–306.
- Bugedo G, Retamal J, Bruhn A. Driving pressure: a marker of severity, a safety limit, or a goal for mechanical ventilation? *Crit Care*. 2017;21(1):199.
- Meier A, Sell RE, Malhotra A. Driving pressure for ventilation of patients with acute respiratory distress syndrome. *Anesthesiology*. 2020;132(6):1569–1576.
- Pelosi P, Ball L. Should we titrate ventilation based on driving pressure? Maybe not in the way we would expect. *Ann Transl Med.* 2018;6(19):389.
- Ramin S, Arcelli M, Bouchdoug K, Laumon T, Duflos C, De Jong A, et al. Driving pressure is not predictive of ARDS outcome in chest trauma patients under mechanical ventilation. *Anaesth Crit Care Pain Med.* 2022;41(4):101095.

- Borges JB, Okamoto VN, Matos GF, Caramez MP, Arantes PR, Barros F, et al. Reversibility of lung collapse and hypoxemia in early acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2006;174:268–278.
- 48. Hodgson CL, Tuxen DV, Davies AR, Bailey MJ, Higgins AM, Holland AE, et al. A randomised controlled trial of an open lung strategy with staircase recruitment, titrated PEEP and targeted low airway pressures in patients with acute respiratory distress syndrome. *Crit Care*. 2011;15:R133.
- Lim SC, Adams AB, Simonson DA, Dries DJ, Broccard AF, Hotchkiss JR, et al. Inter-comparison of recruitment maneuver efficacy in three models of acute lung injury. *Crit Care Med.* 2004;32:2371–2377.
- Fan E, Wilcox ME, Brower RG, Stewart TE, Mehta S, Lapinsky SE, et al. Recruitment maneuvers for acute lung injury: a systematic review. *Am J Respir Crit Care Med.* 2008;178:1156–1163.
- Goligher EC, Hodgson CL, Adhikari NKJ, Meade MO, Wunsch H, Uleryk E, et al. Lung recruitment maneuvers for adult patients with acute respiratory distress syndrome. A systematic review and meta-analysis. *Ann Am Thorac Soc.* 2017;14(Supplement_4):S304–S311.
- 52. Pensier J, de Jong A, Hajjej Z, Molinari N, Carr J, Belafia F, et al. Effect of lung recruitment maneuver on oxygenation, physiological parameters and mortality in acute respiratory distress syndrome patients: a systematic review and meta-analysis. *Intensive Care Med.* 2019;45(12):1691–1702.
- Cui Y, Cao R, Wang Y, Li G. Lung recruitment maneuvers for ARDS patients: a systematic review and meta-analysis. *Respiration*. 2020;99(3):264–276.
- 54. Beitler JR, Sarge T, Banner-Goodspeed VM, Gong MN, Cook D, Novack V, et al, EPVent-2 Study Group. Effect of titrating positive end-expiratory pressure (PEEP) with an esophageal pressure-guided strategy vs an empirical high PEEP-Fio2 strategy on death and days free from mechanical ventilation among patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA*. 2019;321(9):846–857.
- 55. Florio G, Ferrari M, Bittner EA, De Santis Santiago R, Pirrone M, et al, Investigators of the Lung Rescue Team. A lung rescue team improves survival in obesity with acute respiratory distress syndrome. *Crit Care*. 2020;24(1):4.
- 56. Sarge T, Baedorf-Kassis E, Banner-Goodspeed V, Novack V, Loring SH, Gong MN, et al, EPVent-2 Study Group. Effect of esophageal pressure-guided positive end-expiratory pressure on survival from acute respiratory distress syndrome: a risk-based and mechanistic reanalysis of the EPVent-2 trial. *Am J Respir Crit Care Med.* 2021;204:1153–1163.
- Gattinoni L, Giosa L, Bonifazi M, Pasticci I, Busana M, Macri M, et al. Targeting transpulmonary pressure to prevent ventilator-induced lung injury. *Expert Rev Respir Med.* 2019;13(8):737–746.
- Lim J, Litton E. Airway pressure release ventilation in adult patients with acute hypoxemic respiratory failure: a systematic review and meta-analysis. *Crit Care Med.* 2019;47:1794–1799.
- Carsetti A, Damiani E, Domizi R, Scorcella C, Pantanetti S, Falcetta S, et al. Airway pressure release ventilation during acute hypoxemic respiratory failure: a systematic review and meta-analysis of randomized controlled trials. *Ann Intensive Care*. 2019;9(1):44.
- 60. Zhong X, Wu Q, Yang H, Dong W, Wang B, Zhang Z, et al. Airway pressure release ventilation versus low tidal volume ventilation for patients with acute respiratory distress syndrome/acute lung injury: a meta-analysis of randomized clinical trials. *Ann Transl Med.* 2020;8(24):1641.
- Nieman GF, Andrews P, Satalin J, Wilcox K, Kollisch-Singule M, Madden M, et al. Acute lung injury: how to stabilize a broken lung. *Crit Care*. 2018;22(1):136.
- 62. Nieman GF, Gatto LA, Andrews P, Satalin J, Camporota L, Daxon B, et al. Prevention and treatment of acute lung injury with time-controlled adaptive ventilation: physiologically informed modification of airway pressure release ventilation. *Ann Intensive Care.* 2020;10:3.
- Fletcher JJ, Wilson TJ, Rajajee V, Davidson SB, Walsh JC. Changes in therapeutic intensity level following airway pressure release ventilation in severe traumatic brain injury. *J Intensive Care Med.* 2018;33(3):196–202.
- Young D, Lamb SE, Shah S, MacKenzie I, Tunnicliffe W, Lall R, et al, OSCAR Study Group. High-frequency oscillation for acute respiratory distress syndrome. *N Engl J Med.* 2013;368:806–813.
- Ferguson ND, Cook DJ, Guyatt GH, Mehta S, Hand L, Austin P, et al, OS-CILLATE Trial Investigators; Canadian Critical Care Trials Group. Highfrequency oscillation in early acute respiratory distress syndrome. N Engl J Med. 2013;368:795–805.
- Keddissi JI, Youness HA, Jones KR, Kinasewitz GT. Fluid management in acute respiratory distress syndrome: a narrative review. *Can J Respir Ther*. 2019;55:1–8.
- National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network, Wiedemann HP, Wheeler AP,

Bernard GR, Thompson BT, Hayden D, deBoisblanc B, et al. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med.* 2006; 354:2564–2575.

- Humphrey H, Hall J, Sznajder I, Silverstein M, Wood L. Improved survival in ARDS patients associated with a reduction in pulmonary capillary wedge pressure. *Chest.* 1990;97:1176–1180.
- Grissom CK, Hirshberg EL, Dickerson JB, Brown SM, Lanspa MJ, Liu KD, et al. Fluid management with a simplified conservative protocol for the acute respiratory distress syndrome. *Crit Care Med.* 2015;43(2):288–295.
- Seitz KP, Caldwell ES, Hough CL. Fluid management in ARDS: an evaluation of current practice and the association between early diuretic use and hospital mortality. *J Intensive Care.* 2020;8:78.
- Zhang R, Chen H, Gao Z, Liang M, Qiu H, Yang Y, et al. The effect of loop diuretics on 28-day mortality in patients with acute respiratory distress syndrome. *Front Med (Lausanne)*. 2021;8:740675.
- Malbrain MLNG, Marik PE, Witters I, Cordemans C, Kirkpatrick AW, Roberts DJ, et al. Fluid overload, de-resuscitation, and outcomes in critically ill or injured patients: a systematic review with suggestions for clinical practice. *Anaesthesiol Intensive Ther.* 2014;46(5):361–380.
- Ho ATN, Patolia S, Guervilly C. Neuromuscular blockade in acute respiratory distress syndrome: a systematic review and meta-analysis of randomized controlled trials. *J Intensive Care*. 2020;8:12.
- Papazian L, Forel J-M, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med.* 2010;363:1107–1116.
- Gebistorf F, Karam O, Wetterslev J, Afshari A. Inhaled nitric oxide for acute respiratory distress syndrome (ARDS) in children and adults. *Cochrane Database Syst Rev.* 2016;6:CD002787.
- Karam O, Gebistorf F, Wetterslev J, Afshari A. The effect of inhaled nitric oxide in acute respiratory distress syndrome in children and adults: a Cochrane Systematic Review with trial sequential analysis. *Anaesthesia*. 2017;72(1):106–117.
- Adhikari NKJ, Dellinger RP, Lundin S, Payen D, Vallet B, Gerlach H, et al. Inhaled nitric oxide does not reduce mortality in patients with acute respiratory distress syndrome regardless of severity: systematic review and metaanalysis. *Crit Care Med.* 2014;42(2):404–412.
- Tatham KC, Ferguson ND, Zhou Q, Hand L, Austin P, Taneja R, et al. Evolution of practice patterns in the management of acute respiratory distress syndrome: a secondary analysis of two successive randomized controlled trials. J Crit Care. 2021;65:274–281.
- Villar J, Ferrando C, Martínez D, Ambros A, Munoz T, Soler JA, et al. Dexamethasone treatment for the acute respiratory distress syndrome: a multicentre, randomised controlled trial. *Lancet Respir Med.* 2020;8(3): 267–276.
- Chaudhuri D, Sasaki K, Karkar A, Sharif S, Lewis K, Mammen MJ, et al. Corticosteroids in COVID-19 and non-COVID-19 ARDS: a systematic review and meta-analysis. *Intensive Care Med.* 2021;47(5):521–537.
- Wagner C, Griesel M, Mikolajewska A, Mueller A, Nothacker M, Kley K, et al. Systemic corticosteroids for the treatment of COVID-19. *Cochrane Database Syst Rev.* 2021;8(8):CD014963.
- Villar J, Confalonieri M, Pastores SM, Meduri GU. Rationale for prolonged corticosteroid treatment in the acute respiratory distress syndrome caused by coronavirus disease 2019. *Crit Care Explor*. 2020;2:e0111.

- Guerin C, Reignier J, Richard JC, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. N Engl J Med. 2013;368(23):2159–2168.
- Gattinoni L, Carlesso E, Taccone P, Polli F, Guerin C, Mancebo J. Prone positioning improves survival in severe ARDS: a pathophysiologic review and individual patient meta-analysis. *Minerva Anestesiol*. 2010;76(6):448–454.
- 85. McNamee JJ, Gillies MA, Barrett NA, Perkins GD, Tunnicliffe W, Young D, et al. Effect of lower tidal volume ventilation facilitated by extracorporeal carbon dioxide removal vs standard care ventilation on 90-day mortality in patients with acute hypoxemic respiratory failure: the REST randomized clinical trial. *JAMA*. 2021;326(11):1013–1023.
- 86. Bein T, Weber-Carstens S, Goldmann A, Müller T, Staudinger T, Brederlau J, et al. Lower tidal volume strategy (≈3 ml/kg) combined with extracorporeal CO2 removal versus 'conventional' protective ventilation (6 ml/kg) in severe ARDS: the prospective randomized Xtravent-study. *Intensive Care Med.* 2013;39(5):847–856.
- Tonna JE, Abrams D, Brodie D, Greenwood JC, Rubio Mateo-Sidron JA, Usman A, et al. Management of adult patients supported with venovenous extracorporeal membrane oxygenation (VV ECMO): guideline from the Extracorporeal Life Support Organization (ELSO). ASAIO J. 2021;67(6):601–610.
- Peek GJ, Mugford M, Tiruvoipati R, Wilson A, Allen E, Thalanany MM, et al. Efficacy and economic assessment of conventional ventilatory support versus extracorporeal membrane oxygenation for severe adult respiratory failure (CESAR): a multicentre randomised controlled trial. *Lancet.* 2009; 374:1351–1363.
- Combes A, Hajage D, Capellier G, Demoule A, Lavoue S, Guervilly C, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. N Engl J Med. 2018;378(21):1965–1975.
- Goligher EC, Tomlinson G, Hajage D, Wijeysundera DN, Fan E, Juni P, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome and posterior probability of mortality benefit in a post hoc Bayesian analysis of a randomized clinical trial. JAMA. 2018;320(21):2251–2259.
- Combes A, Peek GJ, Hajage D, Hardy P, Abrams D, Schmidt M, et al. ECMO for severe ARDS: systematic review and individual patient data meta-analysis. *Intensive Care Med.* 2020;46(11):2048–2057.
- Munshi L, Walkey A, Goligher E, Pham T, Uleryk E, Fan E. Venovenous extracorporeal membrane oxygenation for acute respiratory distress syndrome: a systematic review and meta-analysis. *Lancet Respir Med*. 2019;7(2):163–172.
- Henry R, Ghafil C, Piccinini A, Liasidis PK, Matsushima K, Golden A, et al. Extracorporeal support for trauma: a trauma quality improvement project (TQIP) analysis in patients with acute respiratory distress syndrome. *Am J Emerg Med*. 2021;48:170–176.
- Khan YA, Fan E, Ferguson ND. Precision medicine and heterogeneity of treatment effect in therapies for ARDS. *Chest*. 2021;160(5):1729–1738.
- Beitler JR, Thompson BT, Baron RM, Bastarache JA, Denlinger LC, Esserman L, et al. Advancing precision medicine for acute respiratory distress syndrome. *Lancet Respir Med.* 2022;10(1):107–120.
- 96. Shah FA, Meyer NJ, Angus DC, Awdish R, Azoulay É, Calfee CS, et al. A research agenda for precision medicine in sepsis and acute respiratory distress syndrome: an official American Thoracic Society Research statement. Am J Respir Crit Care Med. 2021;204(8):891–901.
- Wick KD, McAuley DF, Levitt JE, Beitler JR, Annane D, Riviello ED, et al. Promises and challenges of personalized medicine to guide ARDS therapy. *Crit Care*. 2021;25(1):404.