

The Dubai Definition and Diagnostic Criteria of Laryngopharyngeal Reflux: The IFOS Consensus

Jerome R. Lechien, MD, PhD, MSc ; Michael F. Vaezi, MD, PhD, MS; Walter W. Chan, MD ;
Jacqueline E. Allen, MD ; Petros D. Karkos, MD, PhD, MPhil ; Sven Saussez, MD, PhD;
Kenneth W. Altman, MD, PhD ; Milan R. Amin, MD; Tareck Ayad, MD; Maria R. Barillari, MD, PhD, MS;
Peter C. Belafsky, MD, PhD, MPH ; Joel H. Blumin, MD, FACS ; Nikki Johnston, PhD;
Francois Bobin, MD; Matthew Broadhurst, MD; Fabio P. Ceccon, MD, PhD ;
Christian Calvo-Henriquez, MD ; Young-Gyu Eun, MD, PhD ;
Carlos M. Chiesa-Estomba, MD, PhD, MS ; Lise Crevier-Buchman, MD, PhD, MS; John O. Clarke, MD;
Giovanni Dapri, MD, PhD, FACS; Claudia A. Eckley, MD, PhD; Camille Finck, MD, PhD;
P. Marco Fisichella, MD, FACS; Abdul-Latif Hamdan, MD, MPH, FACS ; Stephane Hans, MD, PhD, MS;
Kathy Huet, PhD ; Rui Imamura, MD, PhD; Blair A. Jobe, MD, FACS; Toshitaka Hoppo, MD, PhD;
Lance P. Maron, MD; Vinciane Muls, MD; Ashli K. O'Rourke, MD ; Paulo S. Perazzo, MD, PhD;
Gregory Postma, MD; Vyas M.N. Prasad, MD, MSc, FRCS ; Marc Remacle, MD, PhD;
Geraldo D. Sant'Anna, MD, PhD; Robert T. Sataloff, MD, FACS ; Edoardo V. Savarino, MD, PhD;
Antonio Schindler, MD, PhD; Nora Siupsinskiene, MD, PhD; Ping-Huei Tseng, MD, PhD;
Craig H. Zalvan, MD ; Karol Zelenik, MD, PhD ; Bernard Frayssse, MD, PhD;
Jonathan M. Bock, MD, FACS ; Lee M. Akst, MD ; Thomas L. Carroll, MD

Objective: The objective of this work was to gather an international consensus group to propose a global definition and diagnostic approach of laryngopharyngeal reflux (LPR) to guide primary care and specialist physicians in the management of LPR.

Methods: Forty-eight international experts (otolaryngologists, gastroenterologists, surgeons, and physiologists) were included in a modified Delphi process to revise 48 statements about definition, clinical presentation, and diagnostic approaches to LPR. Three voting rounds determined a consensus statement to be acceptable when 80% of experts agreed with a rating of at least 8/10. Votes were anonymous and the analyses of voting rounds were performed by an independent statistician.

From the Department of Otorhinolaryngology and Head and Neck Surgery (J.R.L., S.S.), CHU de Bruxelles, CHU Saint-Pierre, School of Medicine, Brussels, Belgium; Department of Laryngology and Broncho-Esophagology (Anatomy Department), EpiCURA Hospital, UMONS Research Institute for Health Sciences and Technology (J.R.L., S.S.), University of Mons (UMons), Mons, Belgium; Department of Otolaryngology-Head and Neck Surgery (J.R.L., L.C.-B., S.H.), Foch Hospital, Paris Saclay University, Research Committee of Young-Otolaryngologists of the International Federations of Oto-Rhino-Laryngological Societies (YO-IFOS), Paris, France; Department of Otorhinolaryngology and Head and Neck Surgery (J.R.L., F.B.), Polyclinic of Poitiers, Poitiers, France; Division of Gastroenterology, Hepatology and Nutrition (M.F.V.), Vanderbilt University Medical Center, Nashville, Tennessee, USA; Division of Gastroenterology, Hepatology and Endoscopy, Department of Medicine (W.W.C.), Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; Department of Otolaryngology, Head and Neck Surgery (J.E.A.), University of Auckland, Auckland, New Zealand; Department of Otorhinolaryngology and Head and Neck Surgery (P.D.K.), Thessaloniki Medical School, Thessaloniki, Greece; Department of Otolaryngology-Head and Neck Surgery (K.W.A.), Geisinger Medical Center, Danville, Pennsylvania, USA; Department of Otolaryngology-Head and Neck Surgery (M.R.A.), New York University School of Medicine, New York, New York, USA; Department of Otolaryngology-Head Neck Surgery (T.A.), Montreal Hospital, Montreal, Canada; Division of Phoniatrics and Audiology, Department of Mental and Physical Health and Preventive Medicine (M.R.B.), L. Vanvitelli University, Naples, Italy; Department of Otolaryngology (P.C.B.), UC Davis Medical Center, Sacramento, California, USA; Department of Otolaryngology and Communication Sciences (J.H.B., N.J., J.M.B.), Medical College of Wisconsin, Milwaukee, Wisconsin, USA; Queensland Voice Centre (M.B.), Brisbane, Australia; Department of Otolaryngology-Head and Neck Surgery (F.P.C.), Hospital Israelita Albert Einstein, São Paulo, Brazil; Department of Otorhinolaryngology and Head and Neck Surgery (C.C.-H.), Hospital Complex of Santiago de Compostela, Santiago de Compostela, Spain; Department of Otolaryngology Head & Neck Surgery, Department of Biomedical Science and Technology, Graduate School (Y.-G.E.), Kyung Hee University, Kyung Hee University School of Medicine, Kyung Hee University Medical Center, Seoul, Republic of Korea; Department of Otorhinolaryngology-Head and Neck Surgery (C.M.C.-E.), Hospital Universitario Donostia, San Sebastian, Spain; Division of Gastroenterology and Hepatology, Department of Medicine (J.O.C.), Stanford University, Redwood City, California, USA; International School Reduced Scar Laparoscopy (G.D.), Minimally Invasive General and Oncologic Surgery Center, Humanitas Gavazzeni University Hospital, Bergamo, Italy; Divisão de Otorrinolaringologia (C.A.E.), Fleury Medicina e Saúde Laboratórios de Diagnóstico, São Paulo, Brazil; Department of Otorhinolaryngology-Head and Neck Surgery (C.F.), CHU de Liege, Liege, Belgium; Abbvie Inc. (P.M.F.), Chicago, Illinois, USA; Department of Otorhinolaryngology-Head and Neck Surgery (A.-L.H.), American University of Beirut Medical Center, Beirut, Lebanon; Department of Metrology and Langage Science (K.H.), University of Mons, Mons, Belgium; Department of Otorhinolaryngology, Clinical Hospital (R.I.), University of São Paulo School of Medicine, São Paulo, Brazil; Department of Surgery (B.A.J., T.H.), Drexel University, Philadelphia, Pennsylvania, USA; Netcare Park Lane Hospital (L.P.M.), Johannesburg, South Africa; Department of gastroenterology (V.M.), CHU Saint-Pierre, Brussels; Department of Otolaryngology, Head and Neck Surgery (A.K.O.), Medical University of South Carolina, Charleston, South Carolina, USA; Department of Otolaryngology, School of Medicine (P.S.P.), São Paulo

Results: After the third round, 79.2% of statements ($N = 38/48$) were approved. LPR was defined as a disease of the upper aerodigestive tract resulting from the direct and/or indirect effects of gastroduodenal content reflux, inducing morphological and/or neurological changes in the upper aerodigestive tract. LPR is associated with recognized non-specific laryngeal and extra-laryngeal symptoms and signs that can be evaluated with validated patient-reported outcome questionnaires and clinical instruments. The hypopharyngeal-esophageal multichannel intraluminal impedance-pH testing can suggest the diagnosis of LPR when there is >1 acid, weakly acid or nonacid hypopharyngeal reflux event in 24 h.

Conclusion: A global consensus definition for LPR is presented to improve detection and diagnosis of the disease for otolaryngologists, pulmonologists, gastroenterologists, surgeons, and primary care practitioners. The approved statements are offered to improve collaborative research by adopting common and validated diagnostic approaches to LPR.

Key Words: consensus, definition, diagnostic, gastroesophageal, guidelines, laryngitis, laryngopharyngeal, reflux.

Level of Evidence: 5

Laryngoscope, 00:1–11, 2023

INTRODUCTION

Laryngopharyngeal reflux (LPR) was initially defined as the backflow of stomach contents into the laryngopharynx.¹ LPR is a disease with different pathophysiological mechanisms and clinical picture than classic gastroesophageal reflux disease (GERD).^{2–4} Various definitions and diagnostic approaches have been published but a universally accepted LPR definition remains lacking.^{5,6} LPR symptoms and findings are prevalent in primary care, otolaryngological, and gastroenterological consultations.^{2,7–10} The non-specificity of symptoms and findings attributed to LPR and the poor effectiveness of empiric proton pump inhibitor (PPI)-therapy make the diagnosis challenging, resulting in over- and under-diagnosis of the condition.^{11–13} Finally, the role of additional testing such as impedance pH-monitoring, remains poorly defined.^{14–17} In this current climate of uncertainty, an international group of experts from the five continents was convened to develop consensus

statements regarding definition and diagnosis of LPR. The evidence used to create the statements of this modified Delphi consensus study was based on expert opinion and was not always based on other guidelines developed from systematic reviews or randomized trials. Because of this methodology, this work should not be considered a clinical guideline. Rather, the IFOS consensus is offered to disseminate expert knowledge from thought leaders who treat and study LPR, and is presented to aid primary care and specialist physicians in the diagnosis of LPR.

METHODS

Development of the consensus definition and diagnostic approach to LPR combined principles of evidence-based medicine and a modified Delphi approach.¹⁸ Experts were invited to vote anonymously on a series of proposed statements through SurveyMonkey® (San Mateo, California, USA), allowing each

Federal University, Sao Paulo, Brazil; Department of Otolaryngology, Head & Neck Surgery (G.P.), Medical College of Georgia at Augusta University, Augusta, Georgia, USA; ENT Centre and Singapore Medical Specialist Centre (V.M.N.P.), Singapore, Singapore; Department of Otolaryngology (M.R.), Centre Hospitalier du Luxembourg, Luxembourg, Luxembourg; Disciplina de Otorrinolaringologia (G.D.S.), Universidade Federal de Ciências da Saúde de Porto Alegre (UFCSPA), Porto Alegre, Brazil; Department of Otolaryngology – Head and Neck Surgery (R.T.S.), Drexel University College of Medicine, Philadelphia, Pennsylvania, USA; Department of Surgery, Oncology and Gastroenterology (E.V.S.), University of Padua, Padua, Italy; Department of Biomedical and Clinical Sciences “L. Sacco” (A.S.), University of Milan, Milan, Italy; Department of Otolaryngology (N.S.), Academy of Medicine, Lithuanian University of Health Sciences, Kaunas, Lithuania; Faculty of Health Sciences (N.S.), Klaipeda university, Klaipeda, Lithuania; Department of Internal Medicine (P.-H.T.), National Taiwan University Hospital, Taipei, Taiwan; Department of Otolaryngology-New York Medical College, School of Medicine, Valhalla, NY; Institute for Voice and Swallowing Disorders, Sleepy Hollow (C.H.Z.), NY, USA; Department of Otorhinolaryngology and Head and Neck Surgery (K.Z.), University Hospital Ostrava, Ostrava, Czech Republic; IFOS Board (B.F.), Paris, France; Department of Otolaryngology-Head and Neck Surgery (L.M.A.), Johns Hopkins Hospital, Johns Hopkins School of Medicine, Baltimore, Maryland, USA; and the Division of Otolaryngology-Head and Neck Surgery, Department of Surgery, Brigham and Women’s Hospital and Department of Otolaryngology-Head and Neck Surgery (T.L.C.), Harvard Medical School, Boston, Massachusetts, USA.

Additional supporting information may be found in the online version of this article.

Editor’s Note: This Manuscript was accepted for publication on October 10, 2023.

J.R.L., W.W.C., and M.F.V. contributed equally to the paper and are joined as co-first authors.

T.L.C., L.M.A., and J.M.B. contributed equally to the paper and are joined as co-senior authors.

The authors whose names are listed immediately below report the following details of affiliation or involvement in an organization or entity with a financial or non-financial interest in the subject matter or materials discussed in this manuscript. The remaining authors have nothing to disclose. Edoardo Vincenzo Savarino has served as speaker for Abbvie, Agave, AGPharma, Alfasigma, Aurora Pharma, CaDiGroup, Celltrion, Dr Falk, EG Stada Group, Fenix Pharma, Fresenius Kabi, Galapagos, Janssen, JB Pharmaceuticals, Innovamedica/Adaclyte, Malesci, Mayoly Biohealth, Omega Pharma, Pfizer, Reckitt Benckiser, Sandoz, SILA, Sofar, Takeda, Tillots, Unifarco; has served as consultant for Abbvie, Agave, Alfasigma, Biogen, Bristol-Myers Squibb, Celltrion, Diadema Farmaceutici, Dr. Falk, Fenix Pharma, Fresenius Kabi, Janssen, JB Pharmaceuticals, Merck & Co, Reckitt Benckiser, Regeneron, Sanofi, SILA, Sofar, Synformulas GmbH, Takeda, Unifarco; he received research support from Pfizer, Reckitt Benckiser, SILA, Sofar, Unifarco, Zeta Farmaceutici. Thomas L. Carroll serves as consultant for Merck and Co, GSK, and N-Zyme Medical. He has Royalties for the book *Chronic Cough* (Plural Publishing). Nikki Johnston is Co-founder, Chief Scientific Officer and an investor in N-Zyme Biomedical. Inventor on: International Patent Application PCT/US2021/027758, Aerosolized formulations of HIV protease inhibitors for the treatment of airway reflux, filed April 16, 2021, and U.S. Patent Application 63/392,929, Sustained-release oral fosamprenavir for the treatment of reflux, filed July 28, 2022. Jerome R. Lechien: consultant for N-Zyme Medical. Peter C. Belafsky: Recipient of California Institute of Regenerative Medicine—Grant Support Innovio Inc—Research Support, Hope Medical (Co-founder), Reflux Gourmet (Co-founder). Kenneth Altman: Merck (consultant, speaker); AXDEV (in a marketing study sponsored by Merck); Vindico (in CME activities sponsored by Merck); Bellus pharmaceuticals (site-PI in a phase 3 clinical trial); Lyra pharmaceuticals (site-PI in a phase 3 clinical trial).

The authors have no funding, financial relationships, or conflicts of interest to disclose.

Send correspondence to Jerome R. Lechien, Department of Laryngology and Bronchoesophagology, Faculty of Medicine, University of Mons (UMONS), Avenue du Champ de mars, 6, B7000 Mons, Belgium. Email: jerome.lechien@umons.ac.be

DOI: 10.1002/lary.31134

participant to complete the survey only once. The statements were written and proposed by a Consensus Committee (CC) of 8 LPR-experienced board-certified physicians selected on their expertise in LPR. The results of the voting rounds were analyzed by a non-voting chair who performed the statistical analyses. The CC provided systematic or state-of-the-art literature reviews to the panel of experts to steer the consensus process away from clinical opinion and towards methodologically sound evidence.¹⁹ The CC agreed to organize the Delphi process through a maximum of 4 voting rounds, each separated by discussion and revision of statements that did not reach validation on prior voting. In iterative fashion, statements that fell short of validation were either improved to the point of reaching consensus or were discarded, with the goal of the statements reaching consensus where possible. The establishment of present consensus was conducted through five steps:

1. The selection of the members of the CC and the expert panel.
2. The draft statement development by the CC.
3. The systematic literature review to support the proposed statements, which included the grading of the evidence.
4. Repeated voting rounds and panel discussion from the results of the second to the third voting round.
5. Writing the present paper and obtaining endorsement by the International Federation of Otorhinolaryngological Societies (IFOS).

Consensus Committee and Expert Panel

The CC was composed of eight experts who come from the five continents. These experts are all members of their respective regional premier scientific societies in otolaryngology-head and neck surgery. The CC selected the relevant papers in the literature review process and developed the initial statements, which were then submitted to the panel of experts. The panel of experts and the CC were organized by the first author and the two senior authors.

The voting panel assembled was comprised of 48 experts from 18 countries. The scientific backgrounds of the experts, including individual numbers of publications on LPR, and inclusion criteria of being chosen an expert are available in Appendix S1. Briefly, experts published a mean (SD) of 34.6 (41.0) peer-reviewed papers on reflux diseases according to PubMed. With the exception of the one voting panel expert chosen in an effort to be inclusive of the African continent, experts were required to have at least one publication specifically on LPR, but most experts, from the required five continents (North America, South America, Europe, Oceania, Asia, Africa), had more than one publication on LPR. The expert voting panel members were required to be currently active, clinically or through research (not retired), and the panel required at least 20% of experts to be outside the field of Otolaryngology but with an expertise in reflux. The CC developed an initial list of 48 statements, which covered a range of important topics related to diagnosis of LPR, including definition of LPR, pathophysiology and differences compared to GERD ($N = 6$); prevalence and incidence of LPR ($N = 4$); contributing factors and associations with other otolaryngological conditions ($N = 6$); symptoms ($N = 4$); findings ($N = 4$); and additional examinations ($N = 24$).

Literature Search

The PubMed, Cochrane Library, and Scopus database literature search were conducted by three authors (J.R.L., S.S., M.R.B.) for relevant peer-reviewed publications in English-language using relevant keywords (Reflux/pH metry/Impedance/Diagnosis/Definition/Symptoms/Signs) to identify and grade the available evidence that was included to support the development of statements. The literature search was performed according to the PRISMA Statements for systematic review. The following

databases were used for the search: PubMed, Scopus and Cochrane Library from 1990 to 2022. Relevant publications were identified, and references of the included papers were further screened for additional research. Several search strings were used that are too numerous to list in the present paper. The complete list of the search strings may be obtained from the first author. The reviewers reviewed each of the abstracts and selected articles for further review. The full texts of selected papers were available to the expert panel for the Delphi process.

Voting Rounds and Discussion

The Delphi process lasted 24 months. Ultimately, there were three voting rounds, separated by periods of time for revision and discussion between voting rounds. In each voting round, a 10-point Likert scale was used to rate each proposed statement from 1 (strongly disagree) to 10 (strongly agree). Consensus acceptance was defined as a Likert rating of $\geq 8/10$ by at least 80% of experts, thus statements reaching this goal were accepted. The analyses of the results of the voting round were performed by an independent statistician (K.H.). At the end of each round, the level of agreement was communicated to the panel as the percentage of experts who voted $\geq 8/10$ for each proposed statement. Statements that returned with only 60%–80% of scores $\geq 8/10$ were discussed by the CC and revised, based on feedback and comments provided by the voting panel—these revised statements were then subjected to additional voting. Statements that did not reach at least 60% agreement of $\geq 8/10$ by the experts were discarded and were not subjected to additional revision or further voting. After the second voting round, a ZOOM® meeting (Zoom Video Communication Inc, San Jose, CA) was organized with committee members and the experts of the panel to further improve those statements that remained in the 60%–80% of scores $\geq 8/10$ and which had not yet been either accepted or discarded.

Grades of Evidence

The assignments of the grade of evidence to the statements was performed by the CC with the GRADE system as is recommended for Position/Consensus papers.²⁰ The grade of the statement aimed to give a practical indication of the likely impact of further research on confidence in the estimate effect. Based on similar work conducted in gut disease,^{19,21} the CC members assigned one of the following grades to each statement:

- High (A): future investigations are unlikely to change our confidence in the estimate effect.
- Moderate (B): future investigations are likely to have an important impact on our confidence in the estimate effect and may change the estimate effect.
- Low (C): future investigations are likely to have an important impact on our confidence in the estimate effect and are very likely to change the estimate effect.
- Very low (D): any estimate of effect is uncertain.

The final assessment of grade was performed by the CC through a consensus discussion.

Endorsement by the International Federation of Otolaryngology Societies

The results of the Delphi process and the present publication were endorsed by the YO-IFOS and the IFOS as the “first world consensus for the definition and the diagnosis of laryngopharyngeal reflux.” Because the first IFOS congress to be held after the voting rounds was complete was held in Dubai (January 2023) and the results first presented there, the name




Fig. 1. Percentage of unvalidated and validated statements throughout the three voting rounds. From the second round to the third round, four items were merged into two items. Thus, seven statements were submitted to the third voting round in place of nine items. [Color figure can be viewed in the online issue, which is available at www.laryngoscope.com.]

for this LPR consensus is the Dubai Definition and Diagnostic Criteria of LPR.

RESULTS AND DISCUSSION

Of the 48 statements initially presented, 38 statements (79.2%) reached agreement considering definition, pathophysiology, and differences with GERD ($N = 3$); prevalence/incidence of LPR ($N = 1$); contributing factors ($N = 2$) and associations with other otolaryngological conditions ($N = 3$); symptoms ($N = 4$); findings ($N = 4$); and additional examinations ($N = 21$). Among them, 23 statements reached consensus after the first round of voting, with 8 and 7 additional statements subsequently reaching consensus after the second and third voting rounds, respectively. The level of consensus increased through the successive voting rounds, with a higher level of consensus in the third round (Fig. 1). All statements proposed in the third round reached consensus. The final validated statements and the related grades of evidence are available in Table I. Several papers dedicated to definition, clinical picture, associations, and diagnostic approaches were reviewed and considered for the discussion of statements (Appendix S2). Figure 2 summarizes LPR-recognized symptoms, findings, and indications of additional examinations as put forth in agreement by our panel of experts. An executive summary of key position statements is reported in Table II.

Definition, Pathophysiology, and Differences with GERD (Statements 1–5)

The alternative names are related to the clinical characteristics of LPR. Because GERD-symptoms and endoscopic findings are absent in many patients,^{22,23} some authors use the terms “silent,” “atypical,” “pharyngeal,”

“proximal,” or “full column,” reflux.²⁴ They highlight the findings specifically identified on objective reflux monitoring such as HEMII-pH.²⁴ The terms “extraesophageal reflux,” “reflux laryngitis,” and “pharyngolaryngeal reflux” may be assigned based on either features on reflux monitoring or the clinical observation of symptoms and signs caused by reflux that has reached the larynx/pharynx.²⁴

From a pathophysiological standpoint, pepsin is involved in the development of inflammatory reaction of the laryngopharyngeal mucosa and related LPR-symptoms and findings.²⁵ The pathophysiological mechanisms of pepsin and its cell toxicity were demonstrated in laboratory studies,^{25,26} and the morphological changes of tissues were observed in vocal folds, laryngeal, and pharyngeal cells.²⁷ Bile salts were found in higher concentration in the saliva of LPR patients compared to controls²⁸ and may be associated with tissue changes at the microscopic level.²⁹ The neurological impact of LPR may manifest as laryngeal sensitivity³⁰ and as abnormalities of the upper esophageal sphincter (UES).²⁴

The pathophysiology of GERD has been extensively studied and primarily involves transient and inappropriate lower esophageal sphincter (LES) relaxation or LES hypotonia with the backflow of gastric content into the esophagus and subsequent troublesome symptoms and/or complications.¹⁹ LPR has the additional process of retrograde flow above the UES into the laryngopharynx.¹⁵ In that way, GERD and LPR share some common pathophysiological mechanisms, although with notable differences in physiology (UES relaxation and deposit of gastric content into the pharynx) and clinical picture (absence of heartburn and GERD-related esophageal symptoms in some patients). The presence of GERD should be a factor when considering the likelihood of LPR.^{31,32} Indeed, Groome et al. demonstrated in a cohort of 1383 GERD patients that there is a

TABLE I.
Definition, Epidemiological, and Association Statements (1 to 3), Clinical Statements (4 to 5) and Diagnostic Statements (6).

| N | Statements | % (R) | Grade |
|---|--|--------|-------|
| 1. Definition, pathophysiology and differences with GERD | | | |
| 1. | LPR has many alternative names including but not limited to silent reflux, pharyngolaryngeal reflux, extraesophageal reflux, atypical GERD, reflux laryngitis, full column reflux, pharyngeal reflux, and proximal reflux. | 90 (1) | NA |
| 2. | LPR is a disease of the upper aerodigestive tract resulting from the direct and/or indirect effects of gastroduodenal content reflux, inducing morphological and/or neurological changes in the upper aerodigestive tract. | 93 (1) | A |
| 3. | LPR and GERD share some common pathophysiological mechanisms but may present with different clinical pictures. | 91 (2) | A |
| 4. | The following may be factors that impair gastroesophageal function and be associated with the development of LPR: tobacco use; obesity; primary esophageal dysmotility; diets high in fat, salt, sugar, mint or acidic content; diets low in protein. | 92 (3) | B |
| 5. | The consumption of alcohol may be a factor that may impair gastroesophageal function and may be associated with the development of LPR. | 83 (1) | B |
| 2. Prevalence and incidence | | | |
| 6. | The prevalence and incidence of LPR are unknown because no objective gold standard parameters are agreed upon for confirming diagnosis. | 83 (1) | C |
| 3. Associations between LPR and otolaryngologic diseases | | | |
| 7. | Although future studies are needed to confirm this association, LPR may be associated with vocal process granuloma. | 83 (1) | B |
| 8. | LPR may impact the clinical course and therapeutic response of patients with asthma, but future prospective studies are needed to confirm the mechanisms underlying the association. | 80 (2) | C |
| 9. | The relationship between LPR and the following ear, nose and throat diseases requires clarification through future studies: paradoxical vocal fold movement; laryngotracheal stenosis; spasmodic dysphonia; laryngeal infections; obstructive sleep apnea; recurrent respiratory papillomatosis; chronic nasal obstruction; olfactory dysfunction; laryngeal cancer; pharyngeal cancer; oral cancer. | 90 (3) | C |
| 4. Symptoms of LPR | | | |
| 10. | LPR may be associated with the following nonspecific ear, nose, and throat symptoms: dysphonia, dysphagia, throat pain, globus sensation, throat clearing, postnasal drip or throat sticky mucus, troublesome cough, cough after lying down/eating, heartburn, and regurgitations. | 90 (2) | B |
| 11. | Typical esophageal symptoms of GERD such as heartburn and digestive symptoms may be present in some LPR patients. | 91 (2) | A |
| 12. | The association between LPR and the following symptoms requires future study: odynophagia, ear pain, burning tongue, halitosis, dyspnea, chest pain, nausea, acid brash, belching, and dyspepsia. | 93 (3) | B |
| 13. | Given the diverse and non-specific nature of symptoms, validated patient-reported outcome questionnaires would ideally be employed in systematic fashion for the diagnosis and follow-up of LPR. | 81 (1) | A |
| 5. Findings of LPR | | | |
| 14. | No specific laryngeal signs are diagnostic of LPR. However, commonly observed signs associated with LPR symptoms may include arytenoid erythema, posterior commissure hypertrophy and erythema, interarytenoid granulation, diffuse laryngeal erythema, vocal fold edema/erythema, pharyngeal erythema, ventricular band erythema/edema, subglottic edema/erythema, post-cricoid erythema/edema, endolaryngeal mucus, and pharyngeal sticky mucus. | 89 (3) | B |
| 15. | Vocal process granuloma can be a sign of LPR, and their presence should prompt LPR evaluation or treatment in the setting of suggestive clinical symptoms. | 80 (1) | B |
| 16. | Regarding the non-specific nature of examination findings, use of a validated instrument would be ideal when assessing LPR-associated findings for both the diagnosis and the follow-up of LPR. | 81 (1) | A |
| 17. | LPR-related laryngeal findings have weak correlations with reported symptoms, esophageal findings on endoscopy, and results of reflux-monitoring studies. Therefore, diagnosis of LPR should not be made based on laryngeal findings alone. | 80 (1) | A |
| 6. Additional examinations | | | |
| 6.1. Upper gastrointestinal endoscopy or transnasal esophagoscopy | | | |
| 18. | The findings of upper gastrointestinal (GI) endoscopy or transnasal esophagoscopy (TNE) may be normal in LPR patients. | 91 (1) | A |
| 19. | Upper GI endoscopy should be performed for all patients with LPR-related symptoms who do not respond to empirical therapeutic trial to evaluate the presence of esophagitis, hiatal hernia, inlet patch of gastric mucosa in the esophagus, gastritis, Helicobacter pylori infection and if HEMII-pH testing demonstrates reflux that was unresponsive to empiric medication trials. | 81 (1) | B |
| 20. | Upper GI endoscopy should be performed for all patients with LPR-related symptoms and concomitant "alarm" features, such as severe dysphagia, hematemesis, unexplained weight loss, or a family history of upper GI tract cancer. | 96 (1) | A |
| 6.2. Esophageal manometry | | | |
| 21. | High-resolution manometry is indicated to evaluate for esophageal dysmotility and/or to assist in the placement of reflux monitoring probe in patients who do not respond to an empiric treatment for suspected LPR. | 81 (1) | C |
| 6.3. Impedance/pH-monitoring | | | |
| 22. | Single-channel (esophageal) or dual-channel (esophageal-esophageal) pH probes are useful for diagnosing GERD but not adequate for diagnosing LPR because of lack pharyngeal sensor and lack of consideration of non-acid event. | 85 (3) | A |
| 23. | If HEMII-pH is unavailable, an empirical treatment covering acid, weakly acid and nonacid LPR may be prescribed and evaluated at 3 months. Treatment success of LPR should be based on improvement of the patient's LPR-symptoms. | 81 (3) | B |
| 24. | The HEMII-pH results may provide guidance as to appropriate nature, dosing, and timing of medications for the specific patient according to the type of LPR (acid, weakly acid, and nonacid) and time of occurrence (upright and daytime and/or nighttime) | 85 (3) | B |

(Continues)

TABLE I.
Continued

| N | Statements | % (R) | Grade |
|-------------------------------|--|--------|-------|
| 25. | Triple-channel (dual esophageal and pharyngeal) pH-only studies may detect acid pharyngeal reflux events but they are not sufficient to rule out LPR disease as they may miss weakly acidic and nonacidic pharyngeal events. | 83 (1) | A |
| 26. | HEMII-pH monitoring has to respect the following placement characteristics: 1. Proximal pH sensor in the hypopharyngeal cavity at 0.5 cm to 1 cm above upper esophageal sphincter; or within the sphincter. 2. Distal pH sensor in the esophagus as close to 5 cm above lower esophageal sphincter as possible. 3. At least 2 impedance sensor pairs in the esophagus. 4. At least 1 impedance sensor pair in the pharyngeal cavity. It is recommended to control the placement of the upper pH sensor by flexible laryngoscopic or manometric guidance. The recommended duration of the examination is 24-h. During the 24-h testing, the patient should continue their normal diet and activities. | 83 (1) | B |
| 27. | On HEMII-pH, hypopharyngeal acid event consists of an event with a pH <4.0. A hypopharyngeal weakly acid reflux event consists of an event with a pH between 4.0 and 7.0. A hypopharyngeal alkaline reflux event consists of an event with pH > 7.0. | 87 (1) | A |
| 28. | The analysis of the 24-h recording must respect the following: 1. Exclusion of reflux events during meals. 2. Pharyngeal reflux event diagnosed only when a reflux event originating from the distal most impedance channel reaches the pharyngeal channels in a retrograde fashion. 3. Manual analysis to identify reflux events that the computer may have reported incorrectly. | 81 (1) | B |
| 29. | The severity of LPR seen on HEMII-pH or oropharyngeal pH monitoring is not necessarily correlated with the severity of symptoms and findings. | 80 (1) | A |
| 30. | While HEMII-pH is promising as an objective tool for diagnosing LPR, the correlation between its findings and treatment outcomes remains limited. Controlled studies are needed to validate the value of this technology in predicting treatment outcomes. | 91 (1) | A |
| 31. | Reflux monitoring for LPR, whether using HEMII-pH, MII-pH, or pH-metry, should be performed off acid suppression medications, beginning at least 7 days prior to the study. | 81 (1) | B |
| 32. | The LPR diagnosis may not be confirmed with esophageal catheters that are configured with two esophageal pH sensors and without impedance or pH sensors in the pharynx because (1) the proximal esophageal reflux events may not reach the hypopharynx and (2) the presence of reflux events near the UES may be altered by swallowing saliva. | 83 (2) | A |
| 33. | Hypopharyngeal-esophageal multichannel intraluminal impedance pH monitoring (HEMII-pH) is an objective tool to identify esophago-pharyngeal reflux events (acid, weakly acidic, or nonacid) and can suggest the diagnosis of LPR when there is >1 hypopharyngeal reflux event in 24 hours. | 90 (2) | B |
| 6.4. Pepsin saliva detection | | | |
| 34. | Pepsin saliva detection is a noninvasive approach that may demonstrate reflux in the pharynx but not confirm the patient's symptoms are due to LPR. | 83 (1) | B |
| 35. | Pepsin may not be detected in saliva of an LPR-suspect patient, although HEMII-pH may confirm LPR. | 81 (1) | A |
| 36. | The pepsin saliva concentration may be influenced by (1) the patient eating before the saliva sample is produced and (2) the time of day the sample is acquired. | 87 (1) | B |
| 37. | Pepsin saliva detection may prove to be a useful adjunctive diagnostic test and/or screening test for LPR but requires further understanding of how diet and sampling frequency influence results. | 85 (2) | B |
| 6.5. Oropharyngeal pH-testing | | | |
| 38. | The criteria for establishing an LPR diagnosis as determined by HEMII-pH are not transferrable to an oropharyngeal-only pH study. | 80 (2) | A |

Abbreviations: GERD = gastroesophageal reflux disease; HEMII-pH = hypopharyngeal-esophageal multichannel intraluminal impedance-pH monitoring; LPR = laryngopharyngeal reflux; N = number; NA = not available; % (R) = percentage of agreement (R = round 1 vs. 2 vs. 3).

correlation between the severity of GERD and the development of LPR.³² This observation was supported in a recent study in which patients with both GERD and LPR demonstrated higher numbers of pharyngeal reflux events on HEMII-pH testing than those with LPR but without GERD.³¹

The influence of tobacco component on esophageal motility and sphincter functioning was supported in a few studies.³³ Obesity is a well-known predisposing factor for GERD, and overweight or obese patients with GERD may demonstrate LPR more often on HEMII-pH testing.³¹ As the peristaltic function of the esophagus is an important defense mechanisms against proximal migration of refluxate, the presence of a primary esophageal dysmotility may be associated with the development of LPR in some patients.⁸ The importance of diet in the development and management of reflux is supported in many studies.^{34,35} High-fat foods are known to be associated with a higher risk of reflux due to their association with a slower gastric emptying time, a decrease of LES

pressure, and an increase in esophageal acid exposure.³³ The impact of mint on sphincter tonicity was supported in some GERD studies,³³ while high-protein foods may increase in LES tonicity.^{33,34}

Regarding coffee, authors did not reach agreement due to conflicting evidence in the literature. Indeed, some studies suggested that coffee (caffeine) ingestion may lead to heartburn in GERD patients³⁶ and decrease the LES pressure,³⁷ while others did not find negative impact of coffee on esophageal function.³⁸ Similar discussion occurred about tea, highlighting the recent meta-analysis investigating the association between tea consumption and the development of GERD in which authors reported that due to the existence of several subgroups of tea drinkers, some develop reflux disease while others do not.³⁹

Prevalence and Incidence of LPR (Statement 6)

Because consensus definition and diagnostic method of LPR did not exist prior to the current attempt,




Fig. 2. Symptoms, signs and additional examinations of laryngopharyngeal reflux (LPR). GI/TN = gastrointestinal/transnasal; HEMII-pH = hypopharyngeal-esophageal multichannel intraluminal impedance-pH monitoring.

TABLE II.
Executive Summary of Key Position Statements.

Executive summary of key position statements

GRADE A: High-evidence findings

LPR results from direct and indirect effects of gastroduodenal content reflux, which induce morphological or neurological changes in mucosa.

LPR and GERD share some common pathophysiological mechanisms but may present with different clinical pictures. Heartburn and other typical digestive symptoms may be absent.

The LPR

Future diagnostic technologies, such as pepsin test or oropharyngeal pH-testing, are not yet evidence-based and require future studies to establish their reliability and potential cutoff for suggesting the diagnostic. The criteria established for HEMII-pH are not transferrable to oropharyngeal pH-testing.

Upper GI endoscopy may be normal in LPR patients and should be performed in patients with “alarm” features, such as severe dysphagia, hematemesis, unexplained weight loss, or family history of upper GI tract cancer.

GRADE B = Moderate-evidence Findings (state-of-the art, requiring future investigations)

The esophageal sphincter relaxations underlying pharyngeal reflux event may be due to tobacco, alcohol, obesity, primary esophageal dysmotility; diets high in fat, salt, sugar, mint or acidic content; and diets low in protein.

LPR is commonly associated with non-specific symptoms and laryngopharyngeal or oral findings for whom the use of clinical instruments is recommended to improve the baseline and posttreatment assessments and changes.

The LPR diagnosis can be based on symptoms or finding only but requires the identification of >1 acid (pH < 4.0), weakly acid (pH = 4.0–7.0) or non-acid (pH > 7.0) pharyngeal reflux event at the 24-hour hypopharyngeal-esophageal multichannel intraluminal impedance-pH monitoring.

The place and specific indications of esophageal manometry

GRADE C-D: Low-evidence findings requiring future investigations

The associations between LPR and the following conditions are plausible but not yet confirmed: Paradoxical vocal fold movement; Laryngotracheal stenosis; Spasmodic dysphonia; Laryngeal infections; Obstructive sleep apnea; Recurrent respiratory papillomatosis; Chronic nasal obstruction; Olfactory dysfunction; Laryngeal cancer; Pharyngeal cancer; Oral cancer, and Asthma.

Abbreviations: GERD = gastroesophageal reflux disease; GI = gastrointestinal; HEMII-pH = hypopharyngeal-esophageal multichannel intraluminal impedance-pH monitoring; LPR = laryngopharyngeal reflux.

estimates on the prevalence and incidence of LPR in the population vary widely. Prior studies have reported a 1%–30% prevalence of LPR-symptoms in otolaryngologic consultation,⁴⁰ or the general population,⁷ but these symptoms are non-specific and may be encountered in many common otolaryngologic conditions.⁴⁰

Associations between LPR and Otolaryngologic Diseases (Statements 7–9)

LPR has been implicated in the development of some ear, nose, and throat conditions based on the detection of pepsin in laryngeal, oral, sinonasal or middle ear mucosa or secretions,²⁵ for example, paradoxical vocal fold movement⁴¹; laryngotracheal stenosis^{41,42}; spasmodic dysphonia⁴¹; obstructive sleep apnea⁴³; recurrent respiratory papillomatosis⁴⁴; chronic nasal obstruction⁴⁵; olfactory dysfunction⁴⁶; and laryngopharyngeal cancer.⁴⁷ In the same vein, several studies reported the occurrence of vocal process granuloma in patients with LPR based on pH-only testing results or clinical observations.⁴⁸

The association between reflux, asthma severity, and therapeutic response has been extensively investigated over the past decades.^{49,50} Despite methodological discrepancies in reflux diagnostic criteria across studies, LPR may be involved in the development of asthma, and antireflux therapy may improve asthma control when both conditions co-exist.^{49,50} Because most pharyngeal reflux events are weakly acid,^{51,52} HEMII-pH needs to be considered more regularly when considering the diagnosis of LPR in patients with asthma, and the other above-mentioned conditions.

SYMPTOMS AND FINDINGS (STATEMENTS 10–13)

LPR symptoms and findings are non-specific because they present as a mucosal inflammatory process in regions of the upper aerodigestive tract which may be commonly encountered in other prevalent otolaryngologic conditions.^{53,54} Patient-reported outcome questionnaires (PROM) assessing severity, frequency, and/or quality-of-life impact of symptoms are used to improve baseline and posttreatment evaluations.^{40,55} The most common LPR PROM is the reflux symptom index (RSI),⁵⁶ which includes nine symptoms. As evidenced by the worldwide use of RSI, the association between LPR and these nine symptoms seems to have been ingrained into clinical practice. Accordingly, the expert panel of the current study also agreed on these symptoms. Throat pain, an additional symptom validated by the experts despite its absence in the RSI, has been associated with LPR in some clinical studies.^{57,58} The development of the reflux symptom score (RSS),⁵⁹ and RSS-12,⁶⁰ with their subsequent validations in German,⁶¹ Chinese⁶² Korean,⁶³ Portuguese,⁶⁴ and Persian⁶⁵ include a broader list of symptoms potentially associated with LPR, for example, odynophagia, ear pain, dyspnea, or halitosis. However, the prevalence of these symptoms in the LPR population requires future controlled studies. This is especially true in regard to the association of LPR and dyspnea, specifically

regarding the impact of LPR on asthma.^{49,50} The term “respiratory reflux” may prove particularly appropriate.

To date, it is important to keep in mind that PROMs used in LPR (e.g., RSI, RSS) include non-specific symptoms, which may be found in other common laryngopharyngeal irritative conditions.^{53,54} Thus, PROMs do not diagnose LPR but rather document laryngopharyngeal symptoms, both at baseline and as they change after treatment. PROMs cannot replace available objective diagnostic tools.

FINDINGS OF LPR (STATEMENTS 14–17)

Similar to symptoms, findings are non-specific and may be found in many inflammatory diseases of the upper aerodigestive tract mucosa.⁶⁶ The non-specificity of signs supports the interest in using clinical instruments to assess LPR.⁶⁷ In 2001, Belafsky et al., developed the reflux finding score (RFS), which considers most laryngeal signs.⁶⁸ The more recent development of the reflux sign assessment (RSA) has added oral and pharyngeal signs to the list of evaluation areas, for example, coated tongue, tongue tonsil hypertrophy or oropharyngeal erythema, that are felt to be more prevalent in LPR patients as compared to healthy individuals.⁶⁸ Laryngeal findings represent one more tool that may suggest possible LPR, but laryngoscopic signs alone cannot be considered diagnostic for LPR given their suboptimal specificity.

ADDITIONAL EXAMINATIONS

Upper Gastrointestinal Endoscopy or Transnasal Esophagoscopy (Statements 18–20)

The main complications of GERD, for example, esophagitis, esophageal hemorrhage, stricture, Barrett’s esophagus and adenocarcinoma,¹⁹ are less prevalent in LPR patients compared to GERD patients.⁶⁹ Erosive esophagitis was found in 10%–30% of LPR patients with an even lower proportion of patients (<10%) demonstrating Barrett’s metaplasia.^{69–71} These observations led to the discussion regarding the place of upper endoscopy in the workup of LPR patients, especially in those without GERD symptoms. The statements supported by our panel of experts may prove more cost-effective, taking into account patient complaints, medical and/or therapeutic history, and family history for consideration of upper endoscopy.

Esophageal Manometry (Statement 21)

Some anatomic and physiologic esophageal characteristics associated with LPR patients have been identified on high-resolution manometry. Notably, esophageal sphincters tonicity and length, intrabolus pressure, proximal or distal esophageal body contractility, intra-abdominal esophagus length and complete bolus clearance decreased among LPR patients.^{8,72} Primary esophageal motility disorders per Chicago classification have also been identified in 43%–63% of patients with LPR symptoms, including ineffective esophageal motility (31%–41%), hypercontractile esophagus (4%–13%), and

disorders of esophagogastric junction outflow (8%–9%).¹⁶ Some manometric features have also correlated with severity of symptoms and reflux, such as the proportion of failed swallows.^{73,74} Thus, primary esophageal and sphincter dysfunction may contribute to both LPR severity and symptoms. However, these manometric findings are not specific to LPR, as they are also seen in GERD or other non-LPR conditions. The use of esophageal manometry is controversial in first-line management of LPR, and thus our experts only recommend high-resolution manometry specifically for the LPR population to identify a primary esophageal dysmotility disorder that may warrant treatment. It is not recommended as a primary diagnostic tool for LPR. Moreover, manometry may assist the placement of pH-impedance catheters.

Impedance/pH-monitoring (Statements 22–33)

The reflux monitoring technologies have evolved throughout the past 3 decades. Currently, most experts agree with the advantage of using HEMII-pH for detecting LPR, as recent studies found that LPR patients more frequently experience weakly acidic or nonacid pharyngeal reflux events.^{41,75} Moreover, only a portion of proximal esophageal reflux events reach the pharynx,⁷⁶ while some patients may have a high proportion of gastroesophageal reflux events reaching the proximal esophagus or pharynx despite a low distal reflux burden,⁵ therefore, making the traditional MII-pH less reliable than HEMII-pH in identifying LPR. In a recent systematic review of normative data for HEMII-pH, authors concluded that the 95th percentile thresholds were 10–73 events for proximal esophageal events, and 0–10 events for hypopharyngeal reflux events using HEMII-pH. The median number of pharyngeal reflux events among healthy individuals was 1,⁵ which supported the cutoff proposed by the current experts. Future studies are needed to confirm this cutoff on large populations of healthy individuals as some papers report hypopharyngeal cutoffs in normal patients ranging from 1 to 4.⁵ The expert panel agreed that any reflux monitoring should be performed off acid suppression therapy such as PPI, which needs to be discontinued at least 7 days prior to the test. Analysis of reflux monitoring studies for LPR should also follow the same basic principles of reflux testing for GERD, including the exclusion of reported mealtimes and the need for manual evaluation of the raw tracing data, rather than relying on automated computer analysis alone. Finally, the expert panel acknowledged the limited evidence currently in correlating reflux study findings and treatment outcomes. As proposed in statement 23, the unavailability of HEMII-pH may justify the use of an empiric therapeutic trial covering acid, weakly acid and non-acid LPR. However, this approach remains controversial: patients may experience unnecessary risk of adverse events through these empiric medication trials and increase costs are possible for both patients and the healthcare system.⁴⁰ Further prospective studies are needed to better clarify the prognostic value of objective reflux testing in predicting LPR symptom outcomes after anti-reflux therapy.

Pepsin Saliva Detection (Statements 34–37)

The sensitivity of pepsin saliva measurement ranges from 29.4% to 86.6% depending on the diagnostic pepsin concentration threshold, time of saliva collection, number of samples, and the method of measurements (e.g., ELISA, Western blot).^{25,26,77} Moreover, studies reported inconsistencies in the associations between HEMII-pH, pepsin saliva measurements, and clinical findings, which may be attributed to the potential contribution of other gastroduodenal enzymes in mucosa inflammation and the development of symptoms and signs.²⁵ All of these issues limit the establishment of salivary pepsin as a conclusive test for LPR, and the expert panel agreed that its role remains as an adjunctive, but not primary, diagnostic test.

Oropharyngeal pH-testing (Statement 38)

A systematic review investigating normative ambulatory reflux monitoring evaluated oropharyngeal pH-testing findings in healthy individuals and reported that the 95th percentiles of oropharyngeal events at thresholds of pH < 4.0, pH < 5.0, and pH < 6.0 ranged from 0 to 2.5, 0 to 107.5, and 40 to 128, respectively.⁵ The data may significantly vary from one study to another due to patient positioning and pH thresholds evaluated and some studies using simultaneous HEMII-pH and oropharyngeal pH-testing failed to demonstrate agreement in number of reflux events (oropharyngeal pH-testing missed true liquid events seen on HEMII-pH); in addition, the technological differences between HEMII-pH and lead to vastly different reported outcomes, thus the numbers obtained from oropharyngeal pH testing are very different and not transferable to HEMII-pH and MII-pH testings.⁵ Oropharyngeal pH-testing remains controversial in its ability to truly diagnose LPR.⁵

CONCLUSION

A global consensus definition for LPR is offered to improve detection and diagnosis, for otolaryngologists, gastroenterologists, surgeons, and primary care practitioners. Such statements may allow collaborative research through studies adopting common language and validated diagnostic approaches.

ACKNOWLEDGMENTS

IRIS-Recherche Grant (Foundation Roi Baudouin), Vesale Grant.

BIBLIOGRAPHY

1. Koufman JA, Aviv JE, Casiano RR, Shaw GY. Laryngopharyngeal reflux: position statement of the committee on speech, voice, and swallowing disorders of the American Academy of Otolaryngology-Head and Neck Surgery. *Otolaryngol Head Neck Surg.* 2002;127(1):32-35. <https://doi.org/10.1067/mhn.2002.125760>.
2. Koufman JA. The otolaryngologic manifestations of gastroesophageal reflux disease (GERD): a clinical investigation of 225 patients using ambulatory 24-hour pH monitoring and an experimental investigation of the role of acid and pepsin in the development of laryngeal injury. *Laryngoscope.* 1991;101(4 Pt 2 Suppl 53):1-78. <https://doi.org/10.1002/lary.1991.101.s53.1>.

3. Lien HC, Wang CC, Kao JY, et al. Distinct physiological characteristics of isolated laryngopharyngeal reflux symptoms. *Clin Gastroenterol Hepatol*. 2020;18(7):1466-1474.e4. <https://doi.org/10.1016/j.cgh.2019.08.064>.
4. Wu Y, Wang J, Huang Q, et al. The relationship between gastroesophageal reflux disease and laryngopharyngeal reflux based on pH monitoring. *Ear Nose Throat J*. 2021;100(4):249-253. <https://doi.org/10.1177/0145561320971915>.
5. Lechien JR, Chan WW, Akst LM, et al. Normative ambulatory reflux monitoring metrics for laryngopharyngeal reflux: a systematic review of 720 healthy individuals. *Otolaryngol Head Neck Surg*. 2022;166(5):802-819. <https://doi.org/10.1177/01945998211029831>.
6. Altman KW, Pruffer N, Vaezi MF. A review of clinical practice guidelines for reflux disease: toward creating a clinical protocol for the otolaryngologist. *Laryngoscope*. 2011;121(4):717-723. <https://doi.org/10.1002/lary.21429>.
7. Gaynor EB. Otolaryngologic manifestations of gastroesophageal reflux. *Am J Gastroenterol*. 1991;86(7):801-808.
8. Sikavi DR, Cai JX, Carroll TL, Chan WW. Prevalence and clinical significance of esophageal motility disorders in patients with laryngopharyngeal reflux symptoms. *J Gastroenterol Hepatol*. 2021;36(8):2076-2082. <https://doi.org/10.1111/jgh.15391>.
9. Strugala V, Dettmar PW, Bittenglova R, Fremundova L, Pešek M. Use of pepsin detection to identify airways reflux in a range of pulmonary diseases. *Clin Respir J*. 2017;11(5):666-667. <https://doi.org/10.1111/crj.12395>.
10. Jung YH, Lee DY, Kim DW, Park SS, Heo EY, Chung HS, Kim DK clinical significance of laryngopharyngeal reflux in patients with chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis*. 2015;10:1343-1351. <https://doi.org/10.2147/COPD.S84337>.
11. Thomas JP, Zubiaur FM. Over-diagnosis of laryngopharyngeal reflux as the cause of hoarseness. *Eur Arch Otorhinolaryngol*. 2013;270(3):995-999. <https://doi.org/10.1007/s00405-012-2244-8>.
12. Francis DO, Vaezi MF. Should the reflux be reflux? Throat symptoms and alternative explanations. *Clin Gastroenterol Hepatol*. 2015;13(9):1560-1566. <https://doi.org/10.1016/j.cgh.2014.08.044>.
13. Lechien JR. Do otolaryngologists over- or underestimate laryngopharyngeal reflux symptoms and findings in clinical practice? A comparison study between the true prevalence and the otolaryngologist-estimated prevalence of symptoms and findings. *J Clin Med*. 2022;11(17):5192. <https://doi.org/10.3390/jcm11175192>.
14. Bor S, Capanoglu D, Vardar R, Woodcock AD, Fisher J, Dettmar PW. Validation of Peptest™ in patients with gastro-esophageal reflux disease and laryngopharyngeal reflux undergoing impedance testing. *J Gastrointest Liver Dis*. 2019;28(4):383-387. <https://doi.org/10.15403/jgld-335>.
15. Lechien JR. Clinical update findings about pH-impedance monitoring features in laryngopharyngeal reflux patients. *J Clin Med*. 2022;11(11):3158. <https://doi.org/10.3390/jcm11113158>.
16. Van Daele DJ. Esophageal manometry, pH testing, endoscopy, and video fluoroscopy in patients with globus sensation. *Laryngoscope*. 2020;130(9):2120-2125. <https://doi.org/10.1002/lary.28289>.
17. de Bortoli N, Martinucci I, Bertani L, et al. Esophageal testing: what we have so far. *World J Gastrointest Pathophysiol*. 2016;7(1):72-85. <https://doi.org/10.4291/wjgp.v7.i1.72>.
18. Murphy MK, Black NA, Lamping DL, et al. Consensus development methods, and their use in clinical guideline development. *Health Technol Assess*. 1998;2(3):i-iv, 1-88.
19. Vakil N, van Zanten SV, Kahrilas P, Dent J, Jones R, Global Consensus Group. The Montreal definition and classification of gastroesophageal reflux disease: a global evidence-based consensus. *Am J Gastroenterol*. 2006;101(8):1900-1920; quiz 1943. <https://doi.org/10.1111/j.1572-0241.2006.00630.x>.
20. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol*. 2011;64:401-406.
21. Pauwels A, Boeckstaens V, Andrews CN, et al. How to select patients for antireflux surgery? The ICARUS guidelines (international consensus regarding preoperative examinations and clinical characteristics assessment to select adult patients for antireflux surgery). *Gut*. 2019;68(11):1928-1941. <https://doi.org/10.1136/gutjnl-2019-318260>.
22. Koufman JA, Belafsky PC, Bach KK, Daniel E, Postma GN. Prevalence of esophagitis in patients with pH-documented laryngopharyngeal reflux. *Laryngoscope*. 2002;112(9):1606-1609. <https://doi.org/10.1097/00005537-200209000-00014>.
23. Lechien JR, Bobin F, Muls V, Saussez S, Hans S. Laryngopharyngeal reflux disease is more severe in obese patients: a prospective multicenter study. *Laryngoscope*. 2021;131(11):E2742-E2748. <https://doi.org/10.1002/lary.29676>.
24. Carroll TL, Nahikian K, Asban A, Wiener D. Nissen fundoplication for laryngopharyngeal reflux after patient selection using dual pH, full column impedance testing: a pilot study. *Ann Otol Rhinol Laryngol*. 2016;125(9):722-728. <https://doi.org/10.1177/0003489416649974>.
25. Samuels TL, Johnston N. Pepsin in gastroesophageal and extraesophageal reflux: molecular pathophysiology and diagnostic utility. *Curr Opin Otolaryngol Head Neck Surg*. 2020;28(6):401-409. <https://doi.org/10.1097/MOO.0000000000000664>.
26. Klimara MJ, Johnston N, Samuels TL, et al. Correlation of salivary and nasal lavage pepsin with MII-pH testing. *Laryngoscope*. 2020;130(4):961-966. <https://doi.org/10.1002/lary.28182>.
27. Lechien JR, Saussez S, Harnegnies B, Finck C, Burns JA. Laryngopharyngeal reflux and voice disorders: a multifactorial model of etiology and pathophysiology. *J Voice*. 2017;31(6):733-752. <https://doi.org/10.1016/j.jvoice.2017.03.015>.
28. Sereg-Bahar M, Jerin A, Jansa R, Stabuc B, Hovevar-Boltezar I. Pepsin and bile acids in saliva in patients with laryngopharyngeal reflux – a prospective comparative study. *Clin Otolaryngol*. 2015;40(3):234-239. <https://doi.org/10.1111/coa.12358>.
29. Vageli DP, Doukas SG, Doukas PG, Judson BL. Bile reflux and hypopharyngeal cancer. *Oncol Rep*. 2021;46(5):244. <https://doi.org/10.3892/or.2021.8195>.
30. Aviv JE, Liu H, Parides M, Kaplan ST, Close LG. Laryngopharyngeal sensory deficits in patients with laryngopharyngeal reflux and dysphagia. *Ann Otol Rhinol Laryngol*. 2000;109(11):1000-1006. <https://doi.org/10.1177/000348940010901103>.
31. Lechien JR, Bobin F, Muls V, et al. Gastroesophageal reflux in laryngopharyngeal reflux patients: clinical features and therapeutic response. *Laryngoscope*. 2020;130(8):E479-E489. <https://doi.org/10.1002/lary.28482>.
32. Groome M, Cotton JP, Borland M, McLeod S, Johnston DA, Dillon JF. Prevalence of laryngopharyngeal reflux in a population with gastroesophageal reflux. *Laryngoscope*. 2007;117(8):1424-1428.
33. Lechien JR, Bobin F, Mouawad F, et al. Development of scores assessing the refluxogenic potential of diet of patients with laryngopharyngeal reflux. *Eur Arch Otorhinolaryngol*. 2019;276(12):3389-3404. <https://doi.org/10.1007/s00405-019-05631-1>.
34. Zalvan CH, Hu S, Greenberg B, Geliebter J. A comparison of alkaline water and mediterranean diet vs proton pump inhibition for treatment of laryngopharyngeal reflux. *JAMA Otolaryngol Head Neck Surg*. 2017;143(10):1023-1029. <https://doi.org/10.1001/jamaoto.2017.1454>.
35. Koufman JA. Low-acid diet for recalcitrant laryngopharyngeal reflux: therapeutic benefits and their implications. *Ann Otol Rhinol Laryngol*. 2011;120(5):281-287. <https://doi.org/10.1177/000348941112000501>.
36. Arivan R, Deepanjali S. Prevalence and risk factors of gastro-esophageal reflux disease among undergraduate medical students from a southern Indian medical school: a cross-sectional study. *BMC Res Notes*. 2018;11(1):448. <https://doi.org/10.1186/s13104-018-3569-1>.
37. Thomas FB, Steinbaugh JT, Fromkes JI, Mekhjian HS, Caldwell JH. Inhibitory effect of coffee on lower esophageal sphincter pressure. *Gastroenterology*. 1980;79(6):1262-1266.
38. Boekema PJ, Samsom M, Smout AJ. Effect of coffee on gastro-oesophageal reflux in patients with reflux disease and healthy controls. *Eur J Gastroenterol Hepatol*. 1999;11(11):1271-1276.
39. Cao H, Huang X, Zhi X, Han C, Li L, Li Y. Association between tea consumption and gastroesophageal reflux disease: a meta-analysis. *Medicine (Baltimore)*. 2019;98(4):e14173. <https://doi.org/10.1097/MD.00000000000014173>.
40. Lechien JR, Akst LM, Hamdan AL, et al. Evaluation and management of laryngopharyngeal reflux disease: state of the art review. *Otolaryngol Head Neck Surg*. 2019;160(5):762-782. <https://doi.org/10.1177/0194599819827488>.
41. Lechien JR, Akst LM, Saussez S, et al. Involvement of laryngopharyngeal reflux in select nonfunctional laryngeal diseases: a systematic review. *Otolaryngol Head Neck Surg*. 2021;164(1):37-48. <https://doi.org/10.1177/0194599820933209>.
42. Aldaharani A, Powell J, Ladak S, et al. The potential role of bile acids in acquired laryngotracheal stenosis. *Laryngoscope*. 2018;128(9):2029-2033. <https://doi.org/10.1002/lary.27105>.
43. Bobin F, Auregan G, Muls V, et al. Impedance-pH monitoring profile of patients with reflux and obstructive sleep apnea syndrome: a controlled study. *Clin Otolaryngol*. 2021;46(4):816-822. <https://doi.org/10.1111/coa.13745>.
44. Formánek M, Komínek P, Jančátová D, et al. Laryngopharyngeal reflux is a potential risk factor for juvenile-onset recurrent respiratory papillomatosis. *Biomed Res Int*. 2019;2019:1463896. <https://doi.org/10.1155/2019/1463896>.
45. Zelenik K, Javorská Z, Taimrová R, et al. Association between inferior turbinate hypertrophy and extraesophageal reflux. *JAMA Otolaryngol Head Neck Surg*. 2022;148(8):773-778. <https://doi.org/10.1001/jamaoto.2022.1638>.
46. Altundag A, Cayonu M, Salihoglu M, et al. Laryngopharyngeal reflux has negative effects on taste and smell functions. *Otolaryngol Head Neck Surg*. 2016;155(1):117-121. <https://doi.org/10.1177/0194599816640249>.
47. Zubčić Ž, Mendeš T, Včeva A, Mihalj H, Bogović V, Milanković SG. Presence of pepsin in laryngeal tissue and saliva in benign and malignant neoplasms. *Biosci Rep*. 2020;40(11):BSR20200216. <https://doi.org/10.1042/BSR20200216>.
48. Pan Y, Li J, Wang X, Zhang J, Zhang C, Liu Z. Prognosis of patients with laryngeal contact granuloma: development and validation of RCGSG score. *Laryngoscope Investig Otolaryngol*. 2022;7(6):1973-1978. <https://doi.org/10.1002/lary.2869>.
49. Carr TF, Peters AT. Chapter 12: asthma: principles of treatment. *Allergy Asthma Proc*. 2012;33(Suppl 1):39-43. <https://doi.org/10.2500/aap.2012.33.3545>.
50. Havemann BD, Henderson CA, El-Serag HB. The association between gastro-oesophageal reflux disease and asthma: a systematic review. *Gut*. 2007;56(12):1654-1664. <https://doi.org/10.1136/gut.2007.122465>.
51. Kang JW, Lee YC, Ko SG, Eun YG. The key timing of pharyngeal reflux in patients with laryngopharyngeal reflux. *Auris Nasus Larynx*. 2022;S0385-8146(22):00221-8. <https://doi.org/10.1016/j.anl.2022.11.002>.
52. Lechien JR, Bobin F, Dapri G, et al. Hypopharyngeal-esophageal impedance-pH monitoring profiles of laryngopharyngeal reflux patients. *Laryngoscope*. 2021;131(2):268-276. <https://doi.org/10.1002/lary.28736>.

53. Ren JJ, Zhao Y, Wang J, et al. PepsinA as a marker of laryngopharyngeal reflux detected in chronic rhinosinusitis patients. *Otolaryngol Head Neck Surg.* 2017;156(5):893-900.
54. Kayali Dinc AS, Cayonu M, Sengezer T, Sahin MM. Smoking cessation improves the symptoms and the findings of laryngeal irritation. *Ear Nose Throat J.* 2020;99(2):124-127. <https://doi.org/10.1177/0145561319881559>.
55. Francis DO, Patel DA, Sharda R, et al. Patient-reported outcome measures related to laryngopharyngeal reflux: a systematic review of instrument development and validation. *Otolaryngol Head Neck Surg.* 2016;155(6):923-935. <https://doi.org/10.1177/0194599816664330>.
56. Belafsky PC, Postma GN, Koufman JA. Validity and reliability of the reflux symptom index (RSI). *J Voice.* 2002;16(2):274-277. [https://doi.org/10.1016/S0892-1997\(02\)00097-8](https://doi.org/10.1016/S0892-1997(02)00097-8).
57. Beaver ME, Stasney CR, Weitzel E, et al. Diagnosis of laryngopharyngeal reflux disease with digital imaging. *Otolaryngol Head Neck Surg.* 2003;128(1):103-108. <https://doi.org/10.1067/mhn.2003.10>.
58. Lechien JR, Bobin F, Muls V, et al. Changes of laryngeal and extralaryngeal symptoms and findings in laryngopharyngeal reflux patients. *Laryngoscope.* 2021;131(6):1332-1342. <https://doi.org/10.1002/lary.28962>.
59. Lechien JR, Bobin F, Muls V, et al. Validity and reliability of the reflux symptom score. *Laryngoscope.* 2020;130(3):E98-E107. <https://doi.org/10.1002/lary.28017>.
60. Lechien JR, Bobin F, Rodriguez A, et al. Development and validation of the short version of the reflux symptom score: reflux symptom Score-12. *Otolaryngol Head Neck Surg.* 2021;164(1):166-174. <https://doi.org/10.1177/0194599820941003>.
61. Bruhn J, Brockmann-Bauser M, Swing T, Bohlender JE, Runggaldier D. Trans-cultural German translation of the reflux symptom score-12 questionnaire. *HNO.* 2022;70(12):886-890. <https://doi.org/10.1007/s00106-022-01233-2>.
62. Zheng XW, Chen LQ, Chen T, et al. Analysis of reliability and validity of the Chinese version of reflux symptomscore 12. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi.* 2022;57(9):1087-1094. <https://doi.org/10.3760/cma.j.cn115330-20220303-00091>.
63. Min HK, Jeon SY, Lechien JR, et al. Translation and validation of the Korean version of the reflux symptom score. *J Voice.* 2021;S0892-1997(21):00306-4. <https://doi.org/10.1016/j.jvoice.2021.08.026>.
64. Ceccon FP, Aires MM, Lechien JR, et al. Translation and cultural adaptation of the short version of the reflux symptom score: reflux symptom score-12 into Brazilian Portuguese. *Braz J Otorhinolaryngol.* 2022;S1808-8694(22):00076-3. <https://doi.org/10.1016/j.bjori.2022.05.005>.
65. Kohansal A, Khoddami SM, Ansari NN, Lechien JR, Aghazadeh K. Validity and reliability of the Persian version of reflux symptom score-12 in patients with laryngopharyngeal reflux disease. *J Voice.* 2022;S0892-1997(22):00224-7. <https://doi.org/10.1016/j.jvoice.2022.07.022>.
66. Eckley CA, Tangerina R. Using RSI and RFS scores to differentiate between reflux-related and other causes of chronic laryngitis. *Braz J Otorhinolaryngol.* 2023;89(1):54-59. <https://doi.org/10.1016/j.bjori.2021.08.003>.
67. Lechien JR, Schindler A, De Marrez LG, et al. Instruments evaluating the clinical findings of laryngopharyngeal reflux: a systematic review. *Laryngoscope.* 2019;129(3):720-736. <https://doi.org/10.1002/lary.27537>.
68. Belafsky PC, Postma GN, Koufman JA. The validity and reliability of the reflux finding score (RFS). *Laryngoscope.* 2001;111(8):1313-1317. <https://doi.org/10.1097/00005537-200108000-00001>.
69. Reichel O, Issing WJ. Should patients with pH-documented laryngopharyngeal reflux routinely undergo oesophagogastrroduodenoscopy? A retrospective analysis. *J Laryngol Otol.* 2007;121(12):1165-1169. <https://doi.org/10.1017/S0022215107000680>.
70. Hoppo T, Sanz AF, Nason KS, et al. How much pharyngeal exposure is "normal"? Normative data for laryngopharyngeal reflux events using hypopharyngeal multichannel intraluminal impedance (HMII). *J Gastrointest Surg.* 2012;16:16-24.
71. Zelenik K, Kajzrlíkova IM, Vitek P, Urban O, Hanousek M, Kominek P. There is no correlation between signs of reflux laryngitis and reflux oesophagitis in patients with gastro-oesophageal reflux disease symptoms. *Acta Otorhinolaryngol Ital.* 2017;37(5):401-405.
72. Tseng WH, Hsu WC, Hsiao TY, et al. Anatomical and physiological characteristics in patients with laryngopharyngeal reflux symptoms: a case-control study utilizing high-resolution impedance manometry. *J Formos Med Assoc.* 2022;121(6):1034-1043. <https://doi.org/10.1016/j.jfma.2021.07.025>.
73. Borges LF, Salgado S, Hathorn KE, Feldman N, Carroll TL, Chan WW. Failed swallows on high-resolution manometry independently correlates with severity of LPR symptoms. *J Voice.* 2022;36(6):832-837.
74. Borges LF, Chan WW, Carroll TL. Dual pH probes without proximal esophageal and pharyngeal impedance may be deficient in diagnosing LPR. *J Voice.* 2019;33(5):697-703.
75. Liu Z, Zhang C, Wang X, et al. Characteristics of laryngopharyngeal reflux in patients of different genders and ages. *J Voice.* 2022;S0892-1997(22):00387-3. <https://doi.org/10.1016/j.jvoice.2022.11.035>.
76. Ulualp SO, Toohill RJ, Hoffmann R, Shaker R. Pharyngeal pH monitoring in patients with posterior laryngitis. *Otolaryngol Head Neck Surg.* 1999;120(5):672-677. <https://doi.org/10.1053/hn.1999.v120.a91774>.
77. Calvo-Henríquez C, Ruano-Ravina A, Vaamonde P, Martínez-Capoccioni G, Martín-Martín C. Is pepsin a reliable marker of laryngopharyngeal reflux? A systematic review. *Otolaryngol Head Neck Surg.* 2017;157(3):385-391. <https://doi.org/10.1177/0194599817709430>.
78. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. <https://doi.org/10.1136/bmj.n71>.