

A systematic review on the impact of financial insecurity on the physical and psychological well-being for people living with terminal illness

Palliative Medicine

1–19

© The Author(s) 2024



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/02692163241257583

journals.sagepub.com/home/pmj



Ross Walker-Pow¹, Andrea Bruun^{1,2}, Nuriye Kupeli¹ , Alessandro Bosco¹
and Nicola White¹

Abstract

Background: People living with terminal illness are at higher risk of experiencing financial insecurity. The variance in definitions of financial insecurity, in addition to its impact on the well-being of this population has not yet been systematically analysed.

Aim: To understand the definition, prevalence and impact of financial insecurity on the physical and psychological well-being of people living with terminal illness.

Design: A systematic review with a narrative synthesis (prospectively registered; CRD42023404516).

Data sources: Medline, Embase, CINAHL, AMED, PsycINFO, ProQuest Central and Cochrane Central Register of Controlled Trials, from inception to May 2023. Included studies had to measure or describe the impact of financial insecurity on an aspect of participants' physical or mental well-being. Study quality was assessed using the Hawker tool.

Results: A total of 26 studies were included in the review. Financial insecurity was defined using many different definitions and terminology. Out of 4824 participants, 1126 (23%) reported experiencing high levels of financial insecurity. Nine studies reported 21 unique analyses across three domains of physical well-being. Out of those 21 analyses, 10 (48%) reported a negative result (an increase in financial insecurity was reported with a decrease in physical well-being). Twenty-one studies reported 51 unique analyses across nine domains of psychological well-being. Out of these analyses, 35 (69%) reported a negative result (an increase in financial insecurity was reported with a decrease in psychological well-being).

Conclusions: People living with terminal illness require support with their financial situation to ensure their well-being is not negatively impacted by financial insecurity.

Keywords

Economics, social sciences, palliative care, quality of life, systematic review

What is already known about the topic?

- People who are living with a terminal illness are at greater risk of experiencing poverty.
- Financial insecurity can significantly impact adherence to treatment plans and can cause undue distress.
- It is unclear how financial insecurity impacts the physical and psychological well-being of people living with terminal illness.

What this paper adds?

- Financial insecurity was defined and discussed in no consistent manner.
- Where financial insecurity was categorised into levels of insecurity, 23% of people living with terminal illness reported experiencing high levels of financial insecurity.
- An increase in financial insecurity was often reported alongside a decrease in the physical and psychological well-being of people living with terminal illness.

¹Marie Curie Palliative Care Research Department, Division of Psychiatry, University College London, London, UK

²Faculty of Health, Science, Social Care and Education, School of Nursing, Allied and Public Health, Kingston University London, London, UK

Corresponding author:

Nicola White, Marie Curie Palliative Care Research Department, Division of Psychiatry, University College London, 6th Floor, Maple House, 149 Tottenham Court Road, London W1T 7NF, UK.
Email: n.g.white@ucl.ac.uk

Implications for practice, theory or policy

- Policy makers should ensure that people living with terminal illness have the support they need to address their financial situation, including signposting to available support; so that they are not left to suffer from the ill-effects of financial insecurity, as well as their illness.
- Standardised terminology, definition and method of assessing financial insecurity would ensure clarity in the need and prevalence of financial insecurity and allow cross-comparison across studies.

Introduction

Poverty is a complex and nuanced global discussion. Terms, definitions, indicators, measurement and thresholds for identifying and assessing poverty vary between and within countries.^{1–3} The current cost-of-living crisis fuelled by rising transport, food and energy prices is having a detrimental impact on global health and well-being.^{4,5} It is estimated that 1 in 10 people in the world live in poverty.^{3,6} A recent report commissioned by Marie Curie estimated that 90,000 people die each year in poverty,⁷ and that people living with a terminal illness are more at risk of poverty; although this data is limited to the UK.

Financial insecurity refers to when an individual encounters objective difficulty or subjective distress when trying to cover necessary financial expenses.⁸ Whilst this is the definition adopted within this review, there is no universally agreed term for this concept.^{9–12} People living with terminal illness (defined as an illness that is not curable and likely to lead to death) are particularly at risk of being impacted by financial insecurity and poverty as they are often forced to give up work due to ill health.^{13,14} This decrease in income is further compounded by the increase in out-of-pocket expenditures associated with having a terminal illness, such as more travel costs for medical appointments and the energy needed to run medical equipment at home.¹⁵ Additional financial insecurity can arise if family members, who support someone with a terminal illness, need to reduce their working hours in order to fulfil their caregiving role.¹⁶ While welfare benefits may be accessed to supplement or serve as one's sole income, these payments have been found to be inadequate for people living with terminal illness.¹⁷ To the authors' knowledge, the extent to which financial insecurity affects people living with terminal illness has not been explored.

The evidence indicates that financial insecurity is a key determinant of poor healthcare experiences.^{18–21} Among people with curable cancer and survivors of cancer, experiencing financial insecurity has been found to reduce quality of life and increase rates of non-adherence to treatment plans,^{22,23} as well as depression, anxiety and psychological distress.²⁴ There is some evidence to suggest that financial insecurity impacts well-being at the end of life although this remains an underexplored area.²⁵

This review seeks to address the gaps in the current knowledge, through synthesising international evidence on financial insecurity for people living with terminal illness.

Review questions

RQ1: How is financial insecurity defined and measured for people living with terminal illness?

RQ2: What is the prevalence of financial insecurity in this population?

RQ3: What is the impact of financial insecurity on the physical and psychological well-being of people living with terminal illness?

Methods

This systematic review followed the Cochrane Handbook for Systematic Reviews²⁶ and is reported in accordance with the Preferred Reporting Items for Systematic Reviews and Analyses (PRISMA).²⁷ The review protocol was registered prospectively in PROSPERO on the 7th of March 2023. (CRD2023404516).

Eligibility criteria

Inclusion

1. Adults (18 years old and above) living with a terminal illness (synonyms: palliative, advanced, life-limiting, metastatic, non-curable, end-stage or stage 4).
2. Measured or described participant's exposure to financial insecurity.
3. Measured or described the impact of this financial insecurity on an aspect of participants' physical and/or psychological well-being.

Both qualitative and quantitative studies were included in this review, due to the multi-faceted experience of financial insecurity.^{28,29}

Exclusion

1. Abstracts (where the full text was not available), editorials, systematic/scoping reviews, research letters, theses.
2. Paediatric population (<18 years)
3. Focus solely on carers and not patients.
4. Not reported in English language.

Information sources

Ovid platform (Medline, Embase, AMED and PsycINFO), CINAHL, ProQuest Central and Cochrane Central Register of Controlled Trials were searched from inception to May 2023. Grey literature was searched for via the OpenGrey website. If only an abstract was identified (i.e. conference abstracts), authors were contacted via email to provide a full text. Additionally, authors of included papers were contacted to check if they had any further relevant published or unpublished work. Forward and backward citation searching was conducted using Google Scholar, to ensure no relevant papers were missed.

Search strategy

The key concepts, taken from the review questions, were ‘terminal illness’ (population), ‘financial insecurity’ (exposure) as well as ‘physical and psychological well-being’ (outcome). A comprehensive list of search terms, both MeSH and textual and their truncated variants were derived from existing literature on terminal illness and financial insecurity.^{13,25} The key concepts were connected using the Boolean operator ‘AND’, meanwhile search terms within each concept were connected using the Boolean operator ‘OR’ (see Supplemental Material 1 for search strategy). Terms were adapted according to each database functionality.

Selection

All references were stored using Rayyan software.³⁰ Following deduplication, two independent reviewers conducted a title and abstract screen. During this stage, the primary aim for the reviewers was to check the population under study (i.e. people living with terminal illness) and whether financial insecurity was being considered. After this initial screen, two independent reviewers conducted a full-text screen, in accordance with the eligibility criteria. During this stage, reasons for exclusion were recorded. Any discrepancies were resolved via discussion between the two reviewers. If discrepancies remained unresolved following discussion, the opinion of third independent reviewer was sought.

Data collection process and data items

Data extraction tables were developed to address the review questions.

The first table detailed the study characteristics, including: author, publication date, country, setting, design, sample size and participant characteristics (i.e. sex, age, ethnicity, employment status and primary diagnosis).

The second data extraction table detailed how each study defined and measured financial insecurity.

Two tables were developed to extract data which described the impact of financial insecurity on physical or psychological well-being. Participant quotes and author

reflections, which described the impact of financial insecurity on well-being, were extracted from qualitative studies and stored in a separate data extraction table.

The data extraction tables were completed by two independent reviewers (RWP and ABr) for the first three studies. Any discrepancies between the reviewers were resolved through discussion with a third independent reviewer (NW). Once the data extraction tables were deemed applicable, data extraction of the remaining studies were then completed by one reviewer (RWP).

Quality appraisal

The quality of included papers was assessed using the Hawker tool.³¹ This tool comprises standardised criteria suitable for appraising the quality of research using any study design.³² Two independent reviewers rated each domain as either very poor (+1), poor (+2), fair (+3) or good (+4). Total scores for each study ranged from 9 to 36. Discrepancies in the total scores awarded were resolved through discussion with a third independent reviewer. Whilst Hawker and colleagues²⁹ did not indicate how total scores could be interpreted, previous systematic reviews using this tool classified studies with a total score of 18 and below, 18–30 and 31–36 as of poor, fair or good quality, respectively.^{33,34} Studies were not excluded based on their quality, but they were reported to allow for a comprehensive understanding of the included studies and the data reported. It also helps to identify and pertinent problematic areas for future research.

Data synthesis

Relevant data from included studies were synthesised using a narrative synthesis approach.³⁵

The terminology, definitions, methods of assessment and extent of financial insecurity were tabulated. Studies were grouped by whether they measured financial insecurity with a continuous score or by grouping the participants (e.g. ‘No financial insecurity’ and ‘High financial insecurity’). A summary of the patient population was provided in relation to their sex, age, employment status and primary diagnosis.

The association between financial insecurity on the physical and psychological well-being of people living with terminal illness was analysed. Due to the substantial level of clinical heterogeneity, a meta-analysis was not completed. Therefore, the findings of quantitative studies, which assessed an impact on the same domain of well-being (e.g. psychological health), were tabulated.

All relevant unique analyses reported in the included studies were extracted, organised by the domain (physical or psychological well-being). Both correlation and predictive analyses were included and documented. To summarise the analyses, the authors extracted the main analysis findings and indicated for each analysis whether the finding was:

- (1) No difference (either correlation or association)
- (2) Negative (where higher financial insecurity was reported alongside a worsening domain)
- (3) Positive (where higher financial insecurity was reported alongside improving domain)

To interpret the correlation analysis, the recommended benchmarks were adopted,³⁶ in which studies reporting an *r* value of 0.3 or less were considered as very weak therefore no correlation was noted. Studies reporting an *r* value between 0.3 and 0.4 were noted as weak, *r* values between 0.401 and 0.699 were noted as moderate and *r* values ≥ 0.7 were interpreted as strong.

Data extracted from qualitative studies was analysed through reflexive thematic analysis.^{37,38} Modelling from previous research using this analysis,³⁹ this method enables the research (RWP) to fully explore and reflect on the data. RWP read each study and extracted all occasions in which financial insecurity was discussed. Through reading each extraction, RWP generated descriptive codes by noting similar experiences. The reflexive element of the thematic analysis was helpful to aid the researcher in getting familiar with the data, engaging and reflecting on the themes generated. The findings were discussed, shared and refined with the research team. The results were tabulated in a hierarchical structure consisting of overall themes and sub-themes, supported by quotes.

Results

Of the 2168 studies identified in the search; 26 studies were included in the final review.^{40–64} See Figure 1 for details of the screening process.

Study Characteristics

The characteristics of the 26 included studies can be found in Table 1. Whilst studies were conducted internationally, most studies were from the USA ($n = 10$).^{40,43–45,47,49,58,59,64,65} and were conducted either in a hospital or a palliative care setting (e.g. hospital ward or hospice). Twenty-one studies employed a quantitative study design,^{42–55,58–61,63–65} four studies employed a qualitative design,^{40,41,56,57} and one study employed both quantitative and qualitative methods.⁶² All studies were cross-sectional or cohort in nature, apart from five studies which were longitudinal.^{44,46,59,64,65}

Patient characteristics

In total, 6024 individuals with terminal illness were included in the review. The majority (83.5%) of individuals had a primary diagnosis of advanced cancer. Approximately 55% of these individuals were female. The mean patient age, across the studies, ranged between 38.7 to 77.8. Ten

studies reported participants' employment status.^{40,41,46,47,49,50,55,58,60,62} Of the 1208 individuals with terminal illness, whose employment status was recorded, 377 (31.2%) were employed, 357 (29.6%) were retired or of retirement age and 474 (39.2%) were unemployed.

Quality appraisal

The results of the quality appraisal, using the Hawker tool, can be found in Supplemental Material 2. Seventeen studies^{40–42,44,46,49,50,52–57,59–62} scored between 18 and 30 and were, therefore, rated as being of fair quality. The other nine studies,^{43,45,47,48,51,58,63–66} scored between 31 and 36 and were, therefore, rated as being of good quality. No studies were rated as being of poor quality.

One area consistently low across all studies, was the ethics and bias domain; in which 21/26 (81%) studies scored as 'Poor'. This was often due to very limited (if any) reflection on the relationship between the researcher and the participant.

Defining and measuring financial insecurity

The terminology and measures used to describe financial insecurity can be seen in Table 2. Studies varied in their use of terminology including: 'Financial difficulty',^{46,48,50–52,55,60,61,63,65} 'Financial distress',^{40,42,44,45,49,53,54,64} 'Financial hardship',^{43,47} 'Financial toxicity',⁵⁸ 'Financial worry',⁵⁹ 'Financial crisis',⁴¹ 'Financial problems',⁵⁶ 'Poverty',⁵⁷ and 'Economic problems'.⁶² Within studies, the definitions and measures used to define and assess financial insecurity varied.

There was little relationship between the geography of the study and the terminology used (see Table 2).

The reported level of financial insecurity

Thirteen studies collected data on financial insecurity through a continuous score.^{43,46–48,51–53,55,58,59,61,63,65} Table 3 shows the range of scores reported, the measures used for financial insecurity and the interpretation of the score. In all but two studies^{39,54} a higher score indicated a higher level of financial insecurity. Four studies all measured financial insecurity using the EORTC-QLQ-C30 tool.⁶⁶ In this tool, the question about finances has a Likert scale (1 = 'Not at all' up to 4 = 'Very much') and the score is reported out of 100 with a higher score indicating a higher level of financial insecurity. The mean scores ranged from 38.3 (SD = 34) up to 80.7 (SD = 22.42).

Püsküllüoğlu et al.⁵⁵ reported the financial scores by several characteristics, in addition to those reported in Table 3. This includes the financial impact on gender (Female = 59.49, SD = 39.74; Male = 77.04, SD = 32.43), on primary site (Breast, Prostate, Lung, GI tract, Unknown, Other) with financial scores ranging between 53.33 (SD = 50.55) for unknown primary up to 84.85 (SD = 22.92) for Other sites;




Figure 1. PRISMA 2020 flow diagram here.

on chemotherapy usage (No chemotherapy = 52.31, SD = 43.18; Chemotherapy = 75.62, SD = 31.02); surgery history (No surgery = 56.50, SD = 40.71; Surgery = 78.43, SD = 30.42); and on education level (Elementary = 82.54, SD = 29.10; Highschool = 58.62, SD = 41.09; University = 68.12, SD = 36.90).

Eight studies^{42,47,51,52,58,62,64,65} reported financial insecurity in terms of categories (i.e. no insecurity, low or mild

insecurity, moderate or some difficulties or high insecurity or difficulties). In two studies,^{51,52} participants were asked about their perceived financial difficulties regarding three domains: the cover of the cost of treatment, taking care of daily needs and buy little 'extras'. They were asked to rate each one as either 'very well' (reported as No insecurity in Table 4), 'fairly well' (reported as Some difficulties in Table 4) or 'poorly' (reported as High/present difficulties on

Table 1. Study and patient characteristics.

Author/s and year	Country	Setting	Sample size	Participant sex (M:F)	Mean participant age (SD)	Ethnicity (n)	Employment status (n)	Primary diagnosis (n)
Adler et al. ⁴⁰	USA	Community and hospital	63	0:63	53.0 (9.0)	Asian (9); Black (18); White (20); Other (16)	Employed (10); Retired/retirement age (6); Unemployed (5); Unemployable due to illness (42)	Advanced cancer (63)
Akter et al. ⁴¹	Bangladesh	Hospital	10	5:5	40.0 and 85.0 (range)	NR	Employed (3); Retired/retirement age (6); Unemployable due to illness (1)	Advanced cancer (2) Other advanced illness (8)
Barbarete et al. ⁴²	France	Hospital	143	73:70	53.8 (16.7) – with financial distress 62.0 (10.5) without financial distress	NR	NR	Advanced cancer (143)
Belcher et al. (2023) ⁴³	USA	Hospital	78	33:34	56.6 (12.2)	Black (36) White (39)	NR	Advanced cancer (78)
Delgado-Guay et al. ⁴⁴	USA	Palliative care clinic	292	156:136	61.0 (NR)	Asian (15); Black (47); Hispanic (36); White (191); Other (3)	NR	Advanced cancer (292)
Delgado-Guay et al. ⁴⁵	Chile, Guatemala and USA	Palliative care clinic	325	132:192	58.0 (NR)	NR	NR	Advanced cancer (325)
Detsyk et al. ⁴⁶	Ukraine	Palliative care clinic	219	103:116	66.2 (12.9)	NR	Employed (45); Retired/retirement age (152); Unemployable due to illness (15)	Advanced cancer (183) Other (36)
Gallups et al. ⁴⁷	USA	Hospital	100	33:67	63.4 (13.05)	Asian (1); Black (26); White (73)	Employed (21); Retired/retirement age (39); Unemployable due to illness (34)	Advanced cancer (100)
Gupta et al. ⁶⁵	USA	Hospital	954	375:579	55.7 (NR)	NR	NR	Advanced cancer (974)
Jacob et al. ⁴⁸	India	Hospital	210	100:110	49.1 (11.9)	NR	NR	Advanced cancer (210)
Kroll et al. ⁴⁹	USA	Hospital	76	37:39	58.1 (12.3)	Asian (1); Black (4); Hispanic (6); White (57); Other (2)	Employed (32); Retired/retirement age (33); Unemployed (6)	Advanced cancer (76)
Lam et al. ⁵⁰	Canada, Taiwan and Cyprus	Hospital	364	79:285	57.7 (13.0)	NR	Employed (42); Retired/retirement age (78); Unemployed (21)	Advanced cancer (364)
Malhotra et al. ⁵²	Singapore	Hospital	600	277:323	60.9 (10.6)	NR	NR	Advanced cancer (600)

(Continued)

Table 1. (Continued)

Author/s and year	Country	Setting	Sample size	Participant sex (M:F)	Mean participant age (SD)	Ethnicity (n)	Employment status (n)	Primary diagnosis (n)
Malhotra et al. ⁵¹	Singapore	Hospital	250	180:70	66.4 (12.0)	Chinese (164); Other (86)	NR	Advanced heart failure (250)
Mercadante et al. ⁵³	Italy	Palliative care clinic	236	114:122	63.9 (11.7)	NR	NR	Advanced cancer (236)
Pérez-Cruz et al. ⁵⁴	Chile	Palliative care clinic	208	104:104	64.0 (14.0)	NR	NR	Advanced cancer (208)
Püsküllüoğlu et al. ⁵⁵	Poland	Hospital	110	45:65	57.8 (13.8)	NR	Employed (20); Retired/retirement age (43); Unemployed (32)	Advanced cancer (110)
Rakic et al. ⁵⁶	Switzerland	Hospital	299	156:143	67.6 (NR)	NR	NR	Advanced cancer (211)
Rattani et al. ⁵⁷	Pakistan	Hospital	4	1:3	38.7 (NR)	NR	NR	Other (88)
Rosenzweig et al. ⁵⁸	USA	Hospital	145	0:145	58.1 (12.5)	Black (7); White (125); Other (8)	Employed (52); Unemployed (83)	Advanced cancer (145)
Segerstrom et al. ⁵⁹	USA	Hospital	102	75:27	60.3 (11.7)	Asian (1); Black (2); White (98); Other (1)	NR	Advanced amyotrophic lateral sclerosis (102)
Sudarisan et al. ⁶⁰	India	Palliative care unit	234	94:140	57.3 (10.8)	NR	Employed (73); Unemployed (161)	Advanced cancer (234)
Tada et al. ⁶¹	Japan	Hospice	24	16:8	62.5 (10.5)	NR	NR	Advanced cancer (24)
Tseng et al. ⁶²	Taiwan	Hospital	153	146:7	56.9 (9.4)	NR	Employed (79); Unemployed (74)	Advanced cancer (153)
Tu and Chiou ⁶³	Taiwan	Hospice	58	29:29	57.7 (15.3)	NR	NR	Advanced cancer (58)
Wang et al. ⁶⁴	USA	Community	779	355:424	77.8 (13.1)	Black (93); Hispanic (191); White (447); Other (48)	NR	Advanced cancer (270)

NR: not reported.

Table 2. Definitions and measures of financial insecurity.

Terminology	First author	Country	Definition	Measure
Financial difficulty	Malhotra et al. ⁵²	Singapore	'The extent to which patients perceive that their financial resources met their financial needs in relation to treatments, daily living expenses, and other obligations' (p. 38, p. 3783; p. 39, p. 1381)	Three questionnaire items: (i) 'How well does the amount of money you have enable you to cover the cost of your treatment?' (ii) 'How well does the amount of money you have take care of your daily needs?' (iii) 'How well does the amount of money you have enable you to buy those little 'extras', that is, those small luxuries?'
	Malhotra et al. ⁵¹	Singapore		
Financial distress	Jacob et al. ⁴⁸	India	No definition provided	European Organization for Research Treatment of Cancer-Quality of Life Questionnaire (EORTC-QLC-C30; 52)
	Detsyk et al. (2020) ⁴⁶	Ukraine	No definition provided	One questionnaire item: 'During the past week, has your physical condition or medical treatment caused you financial difficulties?'
	Gupta et al. ⁶⁵	USA		One questionnaire item: Presence of financial difficulties (yes/no)
	Lam et al. ⁵⁰	Canada, Taiwan and Cyprus	No definition provided	One questionnaire item: Presence of financial difficulties (yes/no)
	Püsküllüoğlu et al. ⁵⁵	Poland		One questionnaire item: Presence of financial difficulties (yes/no)
	Tada et al. ⁶¹	Japan	No definition provided	One questionnaire item: Presence of financial difficulties (yes/no)
	Sudarisan et al. ⁶⁰	India		One questionnaire item: Presence of financial difficulties (yes/no)
	Tu and Chiou ⁶³	Taiwan	No definition provided	One questionnaire item: Presence of financial difficulties (yes/no)
	Adler et al. ⁴⁰	USA	'Stress caused by inadequate financial resources, inability to work, and the precedence of financial challenges over illness concerns' (p. 961)	'Over the last 2 days, have you been disturbed by financial deficiency resulting from your illness?' Thematic analysis of semi-structured interviews conducted with people living with terminal illness
	Barbarete et al. ⁴²	France	'A subjective experience of distress attributed by the patient to financial burden' (p. 2)	Exploratory self-report questionnaire consisting of 11 items
	Mercadante et al. ⁵³	Italy	'The subjective experience of distress attributed to financial problems' (p. 486)	Edmonton Symptom Assessment System-Financial-Spiritual (ESAS-FS; 53, 54)
	Kroll et al. ⁴⁹	USA	'The subjective psychological experience of financial concerns' (p. 4486)	One questionnaire item: 'Please circle the number that best describes your symptoms': 0 (no financial distress) – 10 (worst financial distress)
	Delgado-Guay (2016) ⁴⁴	USA	No definition provided.	
	Delgado-Guay (2016) ⁴⁵	Chile	No definition provided	One questionnaire item: 'What is your current level of financial distress on a scale of 0 to 10 with 0 being no financial distress and 10 being the worst financial distress?'
	Pérez-Cruz et al. ⁵⁴	Chile		
	Wang et al. ⁶⁴	USA	No definition provided	

(Continued)

Table 2. (Continued)

Terminology	First author	Country	Definition	Measure
Financial hardship	Gallups et al. ⁴⁷	USA	No definition provided	20-item Psychological Sense of Economic Hardship Scale (56)
Financial toxicity	Belcher et al. (2023) ⁴³	USA	'The objective financial burden, subjective distress, and behaviours used by patients to cope with costs of cancer and cancer care' (p. 4)	11-item Comprehensive Score for Financial Toxicity (COST; 57)
Financial worry	Rosenzweig et al. ⁵⁸	USA	'Adverse economic consequences resulting from medical treatment' (p. 83)	One questionnaire item: Responded to on a scale of 1 (no worry) to 10 (extremely worried) Thematic analysis of semi-structured interviews conducted with people living with terminal illness, and informal carers
Financial crisis	Segerstrom et al. ⁵⁹	USA	'How much worry (one) was currently experiencing over having enough money to meet financial needs' (p. 1184–1185)	
Financial problems	Akter et al. ⁴¹	Bangladesh	No definition provided	
Poverty	Rakic et al. ⁵⁶	Switzerland	No definition provided	Latent content analysis of medical records of people living with terminal illness, written by healthcare professionals
Economic problems	Rattani et al. ⁵⁷	Pakistan	No definition provided	Analysed semi-structured interviews conducted with people living with terminal illness, informal carers and healthcare providers
	Tseng et al. ⁶²	Taiwan	No definition provided	One questionnaire item from the List of Threatening Experiences questionnaire (58)

Table 3. Financial insecurity scores (continuous measures).

First author	Sample	Additional grouping	Measure	Mean	SD	Measure range
Belcher	75		COST tool	16.8	10.1	0–100
Rosenzweig	145		COST tool	22.6	11.5	0–100
Detsyk	219		EORTC-QLQ-C30	80.7	22.42	0–100
Gupta	954		EORTC-QLQ-C30	38.3	34	0–100
Puskulluoglu	71	Not working	EORTC-QLQ-C30	61.11	37.94	0–100
	53	Over 60	EORTC-QLQ-C30	67.3	39.49	0–100
	57	Up to 60	EORTC-QLQ-C30	66.08	36.46	0–100
	21	Working	EORTC-QLQ-C30	63.33	43.12	0–100
Tada	24		EORTC-QLQ-C30	51.4	45	0–100
Gallups	100		Psychological sense of hardship scale	35.7	13.87	20–73
Mercadante	233		ESAS (item on finances)	3.55	3.1	0–10
Sergerstrom	102		Single item	4.7	2.7	0–10
Jacob	210		Three questions	7.9	1.4	3–9
Malhotra 20	600		3 questions	5.8	1.7	3–9
Malhotra 21	250		3 questions	2.6	2.1	3–9
Tu	58		Bio-psycho-social questions	1.83	1.19	1–5

Table 3). The results of financial insecurity as categorical data are displayed in Table 4. Percentage estimates of high levels of financial insecurity ranged from 9% to 62%.

Six studies presented grouped data on 'No financial insecurity',^{42,51,52,62,64,65} Out of a total sample size of 4579, 35% ($n = 1584$) said they had no experience of financial insecurity. Four studies presented group data on 'Low' or 'Mild' financial insecurity.^{47,58,64–66} Out of a total sample size of 1978, 32% ($n = 639$) reported experiencing low or mild financial insecurity. Three studies presented a moderate group.^{51,52,65} Out of a total sample size of 3504, 41% ($n = 1445$) reported moderate some difficulties. Eight studies reported financial difficulties that were high or present.^{42,47,51,52,58,62,64,65} Out of a total sample size of 4824, there were 23% ($n = 1126$) who reported experiencing high levels of financial insecurity.

The impact of financial insecurity on physical well-being

Table 5 also presents the data reported on psychological well-being. Twenty-one studies^{42,44–54,58–65} reported 51 individual analyses across nine domains of psychological well-being. These domains were: quality of life, emotional well-being, social/familial well-being, spiritual well-being, depression, anxiety, psychological distress and shame and stigma.

The impact of financial insecurity on psychological well-being

Table 5 also presents the data reported on psychological well-being. Twenty-one studies^{42,44–54,58–65} reported 51 individual analyses across nine domains of psychological

well-being. These domains were: quality of life, emotional well-being, social/familial well-being, spiritual well-being, depression, anxiety, psychological distress and shame and stigma.

From the 51 analyses completed, 15 (29%) analyses indicated no association or correlation between financial insecurity and psychological well-being. A total of 35 analyses (69%) reported a negative result (in that financial insecurity was either correlated or predictive of a domain of psychological well-being). One study did not complete an analysis but reported prevalence data, that 41% of the participants felt that financial insecurity impacted their quality of life.⁶³

Of the three studies that adopted a longitudinal design, one reported on the relationship between financial insecurity and mental and physical well-being over time (1 month). Wang et al.⁶⁴ reported that participants who were experiencing financial distress had significantly (statistically and clinically) greater reductions in symptom burden (as measured on the Edmonton Symptom Assessment Scale (ESAS), mean score difference -4.39 , 95% CI -7.91 , -1.17 , $p < .01$). This study was primarily investigating the effectiveness of a home-based palliative care team; to determine if a targeted intervention could reduce symptom burden with a group analysis on financial distress, rather than looking at the impact of financial insecurity over time.

Qualitative findings

Thematic analysis of the four international qualitative studies^{40,41,56,57} generated three themes relevant to the review aims: psychological distress, physical risk-taking and restricted access to treatment (see Supplemental Material 3).

Table 4. Financial insecurity (categorical data).

	Author	Number	Sample size	Percentage
No insecurity	Barbarett	70	143	49%
	Gupta	296	954	31%
	Malhotra 20 – treatment cost	137	600	23%
	Malhotra 20 – daily needs	173	600	29%
	Malhotra 20 – extras	114	600	19%
	Malhotra 21 – treatment costs	91	250	36%
	Malhotra 21 – daily needs	125	250	50%
	Malhotra 21 – extras	75	250	30%
	Tseng	109	153	71%
	Wang	391	779	50%
Total no insecurity		1584	4579	35%
Low insecurity/mild	Gallups	57	100	57%
	Gupta	362	954	38%
	Rosenzweig	34	145	23%
	Wang	186	779	24%
Total low/mild insecurity		639	1978	32%
Moderate insecurity/some difficulties	Gupta	155	954	16%
	Malhotra 20 – treatment cost	331	600	55%
	Malhotra 20 – daily needs	374	600	62%
	Malhotra 20 – extras	320	600	53%
	Malhotra 21 – treatment costs	92	250	37%
	Malhotra 21 – daily needs	86	250	34%
	Malhotra 21 – extras	87	250	35%
Total moderate/some insecurity		1445	3504	41%
High insecurity/presence/difficulties	Barbarett	73	143	51%
	Gallups	43	100	43%
	Gupta	141	954	15%
	Malhotra 20 – treatment cost	132	600	22%
	Malhotra 20 – daily needs	53	600	9%
	Malhotra 20 – extras	166	600	28%
	Malhotra 21 – treatment costs	64	250	26%
	Malhotra 21 – daily needs	36	250	14%
	Malhotra 21 – extras	85	250	34%
	Rosenzweig	90	145	62%
	Tseng	44	153	29%
	Wang	202	779	26%
Total moderate/some insecurity		1126	4824	23%

Psychological Distress. Psychological distress manifested itself in feelings of stress, worry, frustration and helplessness. The stress induced by financial insecurity was described by one individual living with terminal illness:

‘Last year was the worst financial year ever. We were standing in food lines and going to food banks and having to get assistance with the utility bills. It was so stressful for me.’ (40, p. 961-962, Bangladesh)

Additionally, the worry attached to financial insecurity was described by Akter et al.,⁴¹ who alluded to how people living with terminal illness worried about how the cost of their medical care may affect their family members:

‘The respondents mentioned that they felt exhausted worrying about their family members without income.’ (41, p. 6, Switzerland)

People living with terminal illness also expressed their frustration and discontent in relation to being financially dependent on others:

‘I would have to have my boyfriend pay for some of my meds, especially when I didn’t get a paycheck. I can’t even contribute! I hate depending on people.’ (40, p. 962, Bangladesh)

Physical risk-taking. Due to financial insecurity, people living with terminal illness were forced to take risks which

Table 5. Impact of Financial Insecurity on Physical and Psychological well-being.

Overall domain	Sub-domain	First author, date	Measure	Finding/s	Outcome
Physical well-being	Physical well-being	Barbarete et al. ⁴²	FACT-G	Patients with financial distress had lower physical well-being ($M = 14$, $SD = 7$) than patients with no financial distress ($M = 18$, $SD = 7$; $p = .008$)	Negative
		Jacob et al. ⁴⁸	FACT-G	There was no association between financial difficulty and physical well-being ($\beta = 0.311$, $p > 0.05$).	No association
		Malhotra et al. ⁵²	FACT-G	Experiencing higher levels of financial difficulty were associated poorer physical well-being ($\beta = -0.70$, $p < .01$ [95% CI -0.94 , -0.46])	Negative
		Malhotra et al. ⁵¹	FACT-G	There was no association between financial difficulty and physical well-being ($\beta = -0.29$, [95% CI -0.64 to 0.06], $p = \text{NR}$)	No association
		Mercadante et al. (2021) ⁵³	FACT-G	There was no correlation between financial distress and physical well-being ($r_s = 0.140$, $p = 0.033$)	No correlation
		Belcher et al. ⁴³	SF-36	There was no correlation between financial hardship and physical functioning ($r = 0.062$, $p = .599$)	No correlation
				There was no correlation between financial hardship and general health ($r = 0.025$, $p = 832$)	No correlation
		Gupta et al. ⁶⁵	QLI	Experiencing a 10-unit increase in financial difficulty predicted poorer health and physical quality of life ($\beta = -0.52$ [95% CI -0.64 to -0.41], $p < .001$)	Negative
		Gallups et al. ⁴⁷	McCorkle symptom distress	Experiencing increased economic hardship was moderately correlated with an increase in symptom distress ($r = 0.409$, $p < .01$). People with high level of economic hardship reported a higher symptom distress (29.70 ($SD = 9.97$) to people with low levels of economic hardship (22.25 ($SD = 7.44$)). ($p < .001$, 95% CI 4.01 to 10.91).	Negative
		Gallups et al. ⁴⁷	FACT-G	Experiencing increased economic hardship was weakly correlated with a decrease in quality of life ($r = -0.323$, $p < 0.01$). People with low economic hardship scored significantly higher (mean 20.21 ($SD = 6.04$) on physical well-being than those with high economic hardship (mean 13.56 ($SD = 7.63$; $p < .001$)).	Negative
	Functional well-being	Kroll et al. ⁴⁹	PROMIS-10	There was no correlation between financial distress and poorer physical quality of life ($r = -0.25$, $p = .030$)	No correlation
		Barbarete et al. ⁴²	FACT-G	There was no association between financial distress and functional well-being. Financial distress (mean = 13, $SD = 6$); No Financial distress (mean = 14, $SD = 6$), $p = 0.224$.	No association
		Jacob et al. ⁴⁸	FACT-G	Experiencing higher levels of financial difficulty were associated with worsening functional well-being (OLS = 0.438, $p < .01$)	Negative
		Malhotra et al. ⁵²	FACT-G	Experiencing higher levels of financial difficulty predicted poorer functional well-being ($\beta = -0.94$, $p < .01$ [95% CI -1.21 , -0.66])	Negative
		Malhotra et al. ⁵¹	FACT-G	Experiencing higher levels of financial difficulty were associated with poorer functional well-being ($\beta = -0.37$, $p < .05$ [95% CI -0.72 , -0.01])	Negative
		Mercadante et al. (2021) ⁵³	FACT-G	There was no correlation between financial distress and poorer functional well-being ($r_s = -0.155$, $p = .019$)	No correlation
		Belcher et al. (2023) ⁴³	SF-36	There was no correlation between financial hardship and greater role limitations due to physical health ($r = 0.282$, $p = .005$)	No correlation
		Gallups et al. ⁴⁷	FACT-G	People experiencing high economic hardship (mean 14.51 $SD = 6.52$) reported lower functional wellbeing than those with low economic hardship (mean 16.89, SD ; $p = 0.82$, [95% CI -5.07 to 0.30]).	No association

(Continued)

Table 5. (Continued)

Overall domain	Sub-domain	First author, date	Measure	Finding/s	Outcome
Psychological well-being	Physical pain	Jacob et al. ⁴⁸	BPI	There was no association between financial difficulty and pain severity (OLS = 0.109) or pain interference (OLS = 0.0831)	No association
		Malhotra et al. ⁵²	BPI	Experiencing greater financial difficulty was associated with greater pain severity ($\beta = 0.22$ [0.13 to 0.31], $p < 0.01$) as well as pain interference ($\beta = 0.29$ [0.16 to 0.42], $p < 0.01$)	Negative
		Belcher et al. (2023) ⁴³	SF-36	Experiencing greater financial hardship was weakly correlated with an increase in pain due to physical health ($r = 0.320$, $p = 0.005$)	Negative
		Detsyk et al. ⁴⁶	EORTC-QLC-C30	There was no correlation between financial difficulty and quality of life ($r_s = -0.216$, $p < .001$)	No correlation
		Lam et al. ⁵⁰	EORTC QLQ-C30	Experiencing higher levels of financial difficulty was associated with poorer quality of life ($\beta = -0.047$ (SE = 0.011), $p < 0.0001$)	Negative
		Pérez-Cruz et al. ⁵⁴	EORTC QLQ-C15-PAL	Financial distress was not associated with quality of life ($\beta = -0.65$ [-1.73 to 0.42], $p = 0.232$)	No association
		Püsküllüoğlu et al. ⁵⁵	EORTC QLQ-C30	There was no correlation between financial difficulties and quality of life domains ($r =$ not reported).	No correlation
		Tada et al. ⁶¹	EORTC QLQ-C30	Experiencing higher levels of financial difficulty was moderately correlated with poorer quality of life ($r = -0.556$, $p < .001$)	Negative
		Wang et al. ⁶⁴	PROMIS-10	People experiencing moderate to severe financial distress scored higher for physical distress (mean = 29.1, SD = 6.7) compared to those with mild financial distress (mean = 31.4, SD = 6.1) and those with no financial distress (mean = 32.3, SD = 7.0), $p < 0.001$.	Negative
				People experiencing moderate to severe financial distress scored higher for mental distress (mean = 36.9, SD = 8.5) compared to those with mild financial distress (mean = 38.9, SD = 8.2) and those with no financial distress (mean = 40.6, SD = 9.3), $p < 0.001$.	Negative
	Quality of life	Kroll et al. ⁴⁹	PROMIS-10	There was no correlation between mental quality of life and financial difficulties ($r = -0.16$, $p > 0.05$).	No correlation
		Tu and Chioy ⁶³	Original bio-psycho-social questionnaire	41.4% ($n = 24/58$) of patients reported that financial difficulty influences their quality of life	N/A
		Gupta et al. ⁶⁵	QU	Experiencing a 10-unit increase in financial difficulty predicted poorer quality of life ($\beta = -0.40$ [-0.48 to -0.32], $p < .001$)	Negative
		Gallups et al. ⁴⁷	FACT-G	Experiencing increased economic hardship was weakly correlated with poorer quality of life and spiritual wellbeing (combined FACT-G and SWB scores; $r = -0.323$, $p < .01$). People experiencing high levels of economic hardship scored lower (mean = 65.62, SD = 19.29) than those reporting low level of economic hardship (mean = 79.92, SD = 17.23) on the FACT-G scale. ($p < .001$ [95% CI -21.58 to 7.02]). On the combined FACT-G and SWB, people reporting high economic hardship scores lower (mean = 100.02, SD = 27.50) than those with low hardship (mean = 115.72, SD = 22.94), ($p = 0.002$ [95% CI -25.73 to 5.67])	Negative
		Jacob et al. ⁴⁸	FACT-G	No association between financial difficulty and general well-being (OLS = -0.641)	No association
		Rosenzweig et al. ⁵⁸	FACT-B	An increase in financial toxicity was moderately associated with worsening quality of life ($r = 0.56$, $p < .0001$)	Negative

(Continued)

Table 5. (Continued)

Overall domain	Sub-domain	First author, date	Measure	Finding/s	Outcome
Emotional well-being		Barbarett et al. ⁴²	FACT-G	Patients with financial distress had lower emotional well-being ($M = 14$, $SD = 6$) than patients without financial distress ($M = 16$, $SD = 5$, $p = .008$)	Negative
		Jacob et al. ⁴⁸	FACT-G	Experiencing higher levels of financial difficulty were associated with worsening emotional well-being (OLS = 0.627, $p < .01$)	Negative
		Malhotra et al. ⁵²	FACT-G	Experiencing higher levels of financial difficulty was associated with poorer emotional well-being ($\beta = -0.33$, $p < 0.01$ [CI -0.55, -0.12])	Negative
		Malhotra et al. ⁵¹	FACT-G	Experiencing higher levels of financial difficulty were associated with poorer emotional well-being ($\beta = -0.67$, $p < .05$ [CI -1.01, -0.33])	Negative
		Mercadante et al. ⁵³	FACT-G	There was no correlation between financial distress and emotional well-being ($r_s = 0.133$, $p = .045$)	No correlation
		Belcher et al. (2023) ⁴³	SF-36	Experiencing higher levels of financial hardship was weakly correlated with poorer emotional well-being ($r = 0.393$, $p < .001$)	Negative
		Gallups et al. ⁴⁷	FACT-G	People with high economic hardship reported a significantly lower emotional well-being (mean = 14.77, $SD = 6.06$) than those with low economic hardship (mean = 16.89, $SD = 6.84$), $p = .011$ [95% CI -4.87 to 0.65].	Negative
		Barbarett et al. ⁴²	FACT-G	People with financial distress had lower social/familial well-being ($M = 17$, $SD = 5$) than those without financial distress ($M = 19$, $SD = 4$) $p = 0.04$.	Negative
		Jacob et al. ⁴⁸	FACT-G	There was no association between financial difficulty and social/familial well-being (OLS = 0.113, $p > 0.05$)	No association
		Malhotra et al. ⁵²	FACT-G	Experiencing higher levels of financial difficulty was associated with poorer social/familial well-being ($\beta = -0.85$, $p < 0.01$ [95% CI -1.10, -0.61])	Negative
Social/familial well-being		Malhotra et al. ⁵¹	FACT-G	Experiencing higher levels of financial difficulty was associated with poorer social/familial well-being ($\beta = -0.64$, $p < .05$ [95% CI -1.01, -0.26])	Negative
		Mercadante et al. (2021) ⁵³	FACT-G	There was no correlation between financial distress and social/familial well-being ($r_s = -0.188$, $p = 0.004$)	No correlation
		Belcher et al. (2023) ⁴³	SF-36	There was no correlation between financial hardship and social functioning ($r = 0.183$, $p = .119$)	No correlation
		Gupta et al. ⁶⁵	QLU	Experiencing a 10-unit increase in financial difficulty predicted poorer family quality of life ($\beta = -0.24$ [95% CI -0.33 to -0.14], $p < .001$)	Negative
		Gallups et al. ⁴⁷	FACT-G	People with high economic hardship reported a significantly lower social/familial well-being (mean = 22.79, $SD = 6.63$) than those with low economic hardship (mean = 25.28, $SD = 3.61$), $p = .029$ [95% CI -4.73 to 0.26].	Negative
		Jacob et al. ⁴⁸	FACT-SP	Experiencing higher levels of financial difficulty was associated with poorer spiritual well-being (OLS = -1.312, $p < .01$)	Negative
		Malhotra et al. ⁵²	FACT-SP	Experiencing higher levels of financial difficulty was associated with poorer spiritual well-being ($\beta = -0.69$, $p < .01$ [95% CI -1.10, -0.29])	Negative
		Malhotra et al. ⁵¹	FACT-SP	Experiencing higher levels of financial difficulty were associated with poorer spiritual well-being ($\beta = -0.99$, $p < .05$ [95% CI -1.72, -0.26])	Negative
		Pérez-Cruz et al. ⁵⁴	ESAS-FS	Experiencing higher levels of financial difficulty was associated with higher levels of spiritual pain ($\beta = 0.27$ [95% CI 0.14 to 0.41], $p = 0.009$)	Negative
		Delgado-Guay et al. ⁴⁴	ESAS-FS	Experiencing higher levels of financial distress was moderately correlated with higher levels of spiritual pain ($r_s = 0.44$, $p < .001$)	Negative
		Delgado-Guay et al. ⁴⁵	ESAS-FS	There was no correlation between financial distress and spiritual pain ($r = 0.24682$, $p < 0.0001$)	No correlation
Spiritual well-being					

(Continued)

Table 5. (Continued)

Overall domain	Sub-domain	First author, date	Measure	Finding/s	Outcome
Depression		Gupta et al. ⁶⁵	QLI	Experiencing a 10-unit increase in financial difficulty predicted poorer psychological and spiritual quality of life ($\beta = -0.32$ [95% CI -0.44 to -0.21], $p < .001$)	Negative
		Gallups (2017)	SWB-sf	There was no difference in spiritual wellbeing between the economic groups [(high: mean = 34.40, SD = 10.01) vs (low: mean = 35.81, SD = 8.67), $p = .453$ [95% CI -5.13 to 2.30]]	No association
		Barbarett et al. ⁴²	HADS	Patients with financial distress reported more severe depression ($M = 8$, SD = 4) than patients without financial distress ($M = 6$, SD = 4, $p = .007$)	Negative
		Jacob et al. ⁴⁸	HADS	Experiencing higher levels of financial difficulty were associated with worsening depression (OLS = 0.789, $p < .01$)	Negative
		Malhotra et al. ⁵²	HADS	Experiencing higher levels of financial difficulty was associated with higher levels of depression ($B = 0.51$, $p < .01$ [95% CI 0.25, 0.76])	Negative
		Sudarisan et al. ⁶⁰	PHQ-9	Patients with financial distress had more severe depression than patients without financial distress (OR = 3.90 [95% CI 2.00 to 7.61], $p = .001$)	Negative
		Kroll et al. ⁴⁹	CES-D	There was no correlation between financial difficulty and depressive symptoms ($r = 0.29$, $p = 0.010$)	Negative
		Barbarett et al. ⁴²	HADS	Patients with financial distress reported more severe anxiety ($M = 9$, SD = 4) than patients without financial distress ($M = 7$, SD = 4), $p = .009$.	Negative
		Jacob et al. ⁴⁸	HADS	Experiencing higher levels of financial difficulty were associated with worsening anxiety (OLS = 0.396, $p < .01$)	Negative
		Malhotra et al. ⁵²	HADS	Experiencing higher levels of financial difficulty was associated more severe anxiety ($B = 0.52$, $p < .01$ [95% CI 0.25, 0.79])	Negative
Psychological distress		Mercadante et al. (2021) ⁵³	HADS	There was no correlation between financial distress and anxiety ($r_s = 0.192$, $p = 0.003$)	No correlation
		Kroll et al. ⁴⁹	PROMIS-10	There was no correlation between financial difficulty and anxiety ($r = 0.29$, $p < 0.05$).	No correlation
		Gallups et al. ⁴⁷	NCNN distress thermometer	Experiencing higher levels of financial hardship was moderately associated with higher levels of psychological distress ($r = 0.439$, $p < .01$). People experiencing high levels of economic hardship reported more distress (mean = 6.17, SD = 2.91) than those with low economic hardship (mean = 2.65, SD = 2.64), $p < .001$ [95% CI 2.41 to 4.63].	Negative
		Wang et al. ⁶⁴	NCNN Distress Thermometer	People experiencing moderate to severe financial distress scored higher in general distress (6.6 (2.5)), than those with mild financial distress (mean = 5.1, SD = 2.6) or no financial distress (mean = 0, SD = 0.0). $p < .001$.	Negative
		Segerstrom et al. ⁵⁹	Combined scores: BDI, BHS and PSS	There was no correlation between financial worry and poorer psychological health ($r = .07$, $p > 0.05$)	No correlation
		Tseng et al. ⁶²	SSS	Patients who had worry about economic problems experienced greater levels of shame and stigma (mean = 20.7, SD = 12.2) than patients who had no worry about economic problems (mean = 16.2, SD = 12.0; $t = 2.09$, $p = 0.038$)	Negative
Shame and stigma					

N/A: not applicable; EORTC-QLQ-C30: European Organization for Research Treatment of Cancer-Quality of Life Questionnaire; r_s : Spearman rank correlation; BPI: Brief Pain Inventory; PROMIS-10: Patient-Reported Outcomes Measurement Information System; QLI: Quality of Life index; FACT-G: Functional Assessment of Cancer Therapy – General; FACT-B: Functional Assessment of Cancer Therapy – Breast; SF-36: Short Form 36-item Health Survey; FACT-Sp: Functional Assessment of Chronic Illness Therapy–Spiritual-Well-being Scale; ESAS-FS: Edmonton Symptom Assessment System–Financial-Spiritual; HADS: Hospital Anxiety and Depression Scale; PHQ-9: Patient Health Questionnaire-9; NCNN: National Comprehensive Cancer Network; BHS: Beck Depression Inventory; BSS: Beck Hopelessness Scale; PSS: Perceived Stress Scale; SSS: Shame and Stigma Scale.

threatened their physical health. This phenomenon was described by one of the healthcare providers:

'They [patients] do not have any means of transportation. They come in buses. They come in on top of cars'. (67, p. 7, Pakistan)

Restricted access to treatment. Financial insecurity restricted participants' access to treatment. This was illustrated by one informal carer, who talking about her mother, described how:

'She kept her illness a secret because she was unwilling to use any of the scarce family resources to seek medical attention'. (67, p. 7, Pakistan)

Discussion

Main findings of the study

There are three key findings to highlight from this review. Firstly, there is a lack of clarity and consistency on the terminology of what it means to be 'financially insecure' at the end of life. Secondly, approximately 23% of people who have a terminal illness report experiencing high levels of financial insecurity. Finally, an increase in financial insecurity was reported alongside a decrease in physical (48%) and psychological (69%) wellbeing.

What this study adds?

The finding that there was a very high degree of variation of terminology regarding financial insecurity is hugely important. The lack of an agreed definition or terminology for financial insecurity in a palliative care setting could make initiating conversations about the topic very difficult for healthcare professionals, further exacerbating inequities to support.⁶⁷ This is particularly important when reflecting upon findings by Marie Curie, that people who have a terminal illness are more likely to experience financial insecurity.¹⁵

The Marie Curie report estimated that in the UK, every year, 90,000 people die in poverty. They also state that people who are living with terminal illness are at greater risk of experiencing poverty.⁷ This review is the first to synthesise international data to suggest that 23% of people living with terminal illness, almost 1 in 4 people, experience high levels of financial insecurity. Whilst the precision of this figure is limited by methodological and reporting limitations across the included studies, it does help to provide a global barometer at which people who are living with terminal illness might be impacted by financial insecurity. Alongside the figures reporting the global prevalence of poverty (i.e. 1 in 10 people), and the recent data from Marie Curie, these findings indicate a clear need for frequent assessment and normalising conversations, using agreed terminology, about individual finances.

The majority of studies reported an increase in financial insecurity alongside a decrease in physical and/or psychological wellbeing. These findings complement previous evidence that suggests financial insecurity is associated with worse outcomes,^{4,5,13} particularly at the end of life.^{14,15}

Strengths and Limitations

This systematic review included international evidence using both qualitative and quantitative data. A comprehensive search strategy was applied. Despite yielding no results, the grey literature search in addition to the manual forward and backward citation searching ensured no relevant papers were missed. The inclusion and integration of qualitative data, allowed for the nuances and complexities of financial insecurity to be understood in greater depth, how the different healthcare systems create or exacerbate financial difficulties. Due to time constraints and resources, studies not available in English were excluded from this review. This may have resulted in the exclusion of studies with relevant data. Most of the included studies employed a cross-sectional design. Not only does this result in there being less scope to assess the impact of financial insecurity on well-being over time, but it also raises an issue with causation. It has been well evidenced that people from lower socioeconomic groups are at greater risk of poor healthcare outcomes (e.g. higher mortality rates).⁶⁸ One cannot rule out the opposite direction of causality in which poorer well-being caused an individual with terminal illness to have higher financial insecurity. Financial insecurity was primarily measured, in the quantitative studies, using single self-report items derived from pre-existing questionnaires. It is unlikely that one questionnaire item will be able to accurately capture one's level of financial insecurity, including fluctuations over time, given the multi-faceted nature of the experience.^{28,29}

Implications for practice, policy and future research

The prevalence of financial insecurity and lack of an agreed definition or terminology for financial insecurity is a major issue for future practice, policy and research. The topics of death and dying and finances are two taboo topics that can be very difficult to discuss openly. If almost 1 in 4 people are experiencing financial insecurity, it is vital that healthcare professionals across the globe receive clear guidance and support when raising potentially both taboos in the same conversation. Agreed terms can help healthcare professionals to communicate effectively and clearly to identify the needs and necessary support for the individual. Reports such as 'Money matters at the end of life' from the Dying in the Margins project are extremely helpful to develop the building blocks to such conversations.⁶⁹

With clear terminology and definitions, research will be more readily able to make comparisons across studies to investigate the mechanisms which underlie the relationship between financial insecurity and well-being. A better understanding of the impact of financial insecurity on physical and psychological well-being in people who are living with a terminal illness, could enable clinical teams to predict negative consequences more accurately and target interventions more effectively.

This review highlights important global considerations when it comes to wealth and equity in healthcare provision.

In the UK, the National Health Service (NHS) provides health care, free of charge at the point of access. This means that people living with terminal illness are not required to cover the full cost of their care. Interestingly in this review, no studies from the UK were included. Studies were predominately from the USA, where people who are living with terminal illness but do not have appropriate health insurance may be required to pay for all or some of the medical care they receive.¹⁶ In this review, three of the four qualitative studies were conducted in countries where access to medical care relies on out-of-pocket payment (i.e. USA, Bangladesh and Pakistan). These countries are not similar in terms of wealth. Inevitably because of this wealth difference, the financial support at the end of life will vary across countries. In the UK, changes to the Social Security (Special Rules for End of Life) Bill have enabled individuals in the last 12 months of their lives to apply for additional financial support.^{14,70} In the USA, hospice access through Medicaid or Medicare (the largest USA healthcare insurance providers) is available for individuals who have a predicted survival of 6 months or less.⁷¹

Pickett and Wilkinson⁷² bring forward an important discussion about inequity and healthcare access from an international perspective. They argue that it is in unequal societies, not poor societies, where more health and social care problems exist. They note that life expectancy, for example, is lower in unequal societies. In this review, financial insecurity was experienced across countries with different economic structures and resources: suggesting that there is more to this complex phenomenon than global wealth, but rather availability and equitable access to support.

Conclusion

Financial insecurity is associated with reduced physical and psychological well-being among people living with terminal illness. That said, it continues to be poorly defined and inconsistently measured. Future research should seek to provide a uniform definition to aid healthcare professionals in initiating potentially challenging conversations about death and finances. Policy makers should ensure that people living with terminal illness have the support they need to address their financial situation,

including signposting to available support. This support would mean that they are not left to suffer from the ill-effects of financial insecurity, as well as their illness.

Acknowledgements

We would like to acknowledge Veronica Parisi (Librarian at UCL) for their support to develop robust search terms. We would like to also acknowledge Dr Victoria Vickerstaff for her input during the analysis of the quantitative data.

Author contributions

NW and NK: concept, review procedures, analysis, interpretation and write up. RWP and ABr: protocol development, review procedures, analysis, interpretation and write up. ABo: analysis, interpretation and write up. All authors approved the final draft.

Data management and sharing

All available data is reported in the manuscript and Supplemental Material.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding


The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Marie Curie Core funding (MCCC-FPO-16-U) and Marie Curie Research Grant Scheme 2022 (MCRGS-EOI-56); NK was supported by an Alzheimer's Society Junior Fellowship, Grant/Award Number: 399 AS-JF-17b-016.

Research ethics and patient consent

Ethical approval and patient consent were not required since the review only involved secondary analysis of published data.

ORCID iDs

Nuriye Kupeli  <https://orcid.org/0000-0001-6511-412X>

Nicola White  <https://orcid.org/0000-0002-7438-0072>

Supplemental material

Supplemental material for this article is available online.

References

1. Moss JL, Pinto CN, Srinivasan S, et al. Persistent poverty and cancer mortality rates: an analysis of county-level poverty designations. *Cancer Epidemiol Biomarkers Prev* 2020; 29: 1949–1954.
2. Roser M. *Global poverty in an unequal world: who is considered poor in a rich country? And what does this mean for our understanding of global poverty?* <https://ourworldindata.org/higher-poverty-global-line> (2021, accessed 24 April 2024).
3. World B. Poverty and Inequality Platform (version 20240326_2017_01_02_PROD) [data set], 2024.

4. Royal College of P. Over half of Brits say their health has worsened due to rising cost of living, <https://www.rcplondon.ac.uk/news/over-half-brits-say-their-health-has-worsened-due-to-rising-cost-of-living> (2022, accessed 24 April 2024).
5. Kilfoyle M. *How is the cost of living crisis affecting public health?* Economics Observatory, <https://www.economic-observatory.com/how-is-the-cost-of-living-crisis-affecting-public-health> (2023, accessed 24 April 2024).
6. World Bank Group. Poverty and shared prosperity: taking on inequality, <https://www.worldbank.org/en/publication/poverty-and-shared-prosperity-2016> (2016, accessed 24 April 2024).
7. Stone J and Hirsch D. *Poverty at the end of life in the UK*. Marie Curie. <https://www.mariecurie.org.uk/globalassets/media/documents/policy/dying-in-poverty/h420-poverty-at-the-end-of-life-in-the-uk-2nd-pp.pdf> (2022, accessed 24 April 2024).
8. Joseph Rowntree F. *UK Poverty 2022. The essential guide to understanding poverty in the UK*, <https://www.jrf.org.uk/uk-poverty-2022-the-essential-guide-to-understanding-poverty-in-the-uk> (2022, accessed 24 April 2024).
9. Fischer E, Martin KD, Kamakura WA, et al. Financial insecurity and deprivation life satisfaction, self determination, and consumption adequacy at the bottom of the pyramid how economic contractions and expansions affect expenditure patterns living U.S. capitalism: the normalization of credit/debt financial deprivation prompts consumers to seek scarce goods how marketplace performances produce interdependent status games and contested forms of symbolic capital. *J Consum Res* 2013; 39: vii–x.
10. Basyouni SS and El Keshky MES. Job insecurity, work-related flow, and financial anxiety in the midst of COVID-19 pandemic and economic downturn. *Front Psychol* 2021; 12: 632265.
11. Osberg L. *How should one measure economic insecurity?* 2015. DOI: 10.1787/5JS4T78Q9LQ7-EN
12. Bossert W and D'Ambrosio C. Measuring economic insecurity. *Int Econ Rev* 2013; 54: 1017–1030.
13. Brown SB and D'Angelo K. Financial insecurity. In: Harrison JA, Preeti D, Kelly MD, et al. (eds), *Social emergency medicine*. Cham: Springer International Publishing, 2021, pp.199–215.
14. Kaushal A, Yusuf M and Blodgett J. *Experience of terminal illness in working age people: a scoping review*, <https://whatworkswellbeing.org/resources/experience-of-terminal-illness-in-working-age-people-a-scoping-review/> (2022).
15. Marie Curie. *Dying in poverty: Exploring poverty at the end of life in the UK*, 2022.
16. Gardiner C, Brereton L, Frey R, et al. Exploring the financial impact of caring for family members receiving palliative and end-of-life care: a systematic review of the literature. *Palliat Med* 2014; 28: 375–390.
17. Marie Curie. *The cost of dying. The financial impact of terminal illness*, 2019.
18. Rowley J, Richards N, Carduff E, et al. The impact of poverty and deprivation at the end of life: a critical review. *Palliat Care Soc Pract* 2021; 15. DOI: 10.1177/26323524211033873
19. Fitch C, Hamilton S, Bassett P, et al. The relationship between personal debt and mental health: a systematic review. *Mental Health Rev J* 2011; 16: 153–166.
20. Turunen E and Hiilamo H. Health effects of indebtedness: a systematic review. *BMC Public Health* 2014; 14: 489–489.
21. Guan N, Guariglia A, Moore P, et al. Financial stress and depression in adults: a systematic review. *PLOS ONE* 2022; 17: e0264041.
22. Sharp L, Carsin AE and Timmons A. Associations between cancer-related financial stress and strain and psychological well-being among individuals living with cancer. *Psychooncology* 2013; 22: 745–755.
23. Bhanvadia SK, Psutka SP, Burg ML, et al. Financial toxicity among patients with prostate, bladder, and kidney cancer: a systematic review and call to action. *Eur Urol Oncol* 2021; 4: 396–404.
24. Altice CK, Banegas MP, Tucker-Seeley RD, et al. Financial hardships experienced by cancer survivors: a systematic review. *J Natl Cancer Inst* 2017; 109: djw205.
25. Hanratty B, Holland P, Jacoby A, et al. Review article: Financial stress and strain associated with terminal cancer—a review of the evidence. *Palliat Med* 2007; 21: 595–607.
26. Higgins JPT, Thomas J, Chandler J, et al. *Cochrane handbook for systematic reviews of interventions version 6.4 (updated August 2023)*. Cochrane, 2023.
27. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021; 88: 105906.
28. French D and Vigne S. The causes and consequences of household financial strain: a systematic review. *Int Rev Financ Anal* 2019; 62: 150–156.
29. Pauge S, Surmann B, Mehliis K, et al. Patient-reported financial distress in cancer: a systematic review of risk factors in universal healthcare systems. *Cancers* 2021; 13: 5015.
30. Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan—a web and mobile app for systematic reviews. *Syst Rev* 2016; 5: 1–10.
31. Hawker S, Payne S, Kerr C, et al. Appraising the evidence: reviewing disparate data systematically. *Qual Health Res* 2002; 12: 1284–1299.
32. Hong QN and Pluye P. A conceptual framework for critical appraisal in systematic mixed studies reviews. *J Mixed Methods Res* 2019; 13: 446–460.
33. den Boer K, de Veer AJE, Schoonmade LJ, et al. A systematic review of palliative care tools and interventions for people with severe mental illness. *BMC Psychiatry* 2019; 19: 106–106.
34. McInerney D, Kupeli N, Stone P, et al. Emotional disclosure in palliative care: a scoping review of intervention characteristics and implementation factors. *Palliat Med* 2021; 35: 1323–1343.
35. McKenzie JE and Brennan SE. Synthesizing | presenting findings using other methods. In: Higgins JPT, Thomas J, Chandler J, et al. (eds) *Cochrane handbook for systematic reviews of interventions version 6.4 (updated August 2023)*, 2023.
36. Akoglu H. User's guide to correlation coefficients. *Turk J Emerg Med* 2018; 18: 91–93.
37. Braun V and Clarke V. Reflecting on reflexive thematic analysis. *Qual Res Sport Exerc Health* 2019; 11: 589–597.
38. Braun V and Clarke V. Supporting best practice in reflexive thematic analysis reporting in palliative medicine: a review of published research and introduction to the Reflexive Thematic Analysis Reporting Guidelines (RTARG). *Palliat Med*. Epub ahead of print 12 March 2024. DOI: 10.1177/02692163241234800.
39. Hanna JR, Rapa E, Dalton LJ, et al. Health and social care professionals' experiences of providing end of life care

- during the COVID-19 pandemic: a qualitative study. *Palliat Med* 2021; 35: 1249–1257.
40. Adler SR, Coulter YZ, Stone K, et al. End-of-life concerns and experiences of living with advanced breast cancer among medically underserved women. *J Pain Symp Manage* 2019; 58: 959–967.
 41. Akter S, Sarker M, Hossain P, et al. Solidarity and suffering: enrolled terminal patients' and their caregiver's experiences of the community-based palliative care programme in an urban slum of Bangladesh. *Palliat Care Soc Pract* 2022; 16. DOI: 10.1177/26323524221095104
 42. Barbaret C, Brosse C, Rhondali W, et al. Financial distress in patients with advanced cancer. *PLoS ONE* 2017; 12(5): e0176470. DOI: 10.1371/journal.pone.0176470
 43. Belcher SM, Lee H, Nguyen J, et al. Financial hardship and quality of life among patients with advanced cancer receiving outpatient palliative care: a pilot study. *Cancer Nurs* 2023; 46: 3–13.
 44. Delgado-Guay MO, Chisholm G, Williams J, et al. Frequency, intensity, and correlates of spiritual pain in advanced cancer patients assessed in a supportive/palliative care clinic. *Palliat Support Care* 2016; 14: 341–348.
 45. Delgado-Guay MO, Palma A, Duarte ER, et al. Spirituality, religiosity, spiritual pain, and quality of life among caregivers of Latin American patients with advanced cancer: a multicenter study. *J Clin Oncol* 2016; 34: 245–245.
 46. Detsyk OZ, Bratsyun OP and Babich PM. Quality of life of patients receiving home-based palliative care from family physicians and mobile palliative care team. *Wiad Lek* 2020; 73: 1681–1689.
 47. Gallups S, Copeland VC and Rosenzweig M. Perceived financial hardship among patients with advanced cancer. *J Community and Support Oncol* 2017; 15: e163–e169.
 48. Jacob J, Palat G, Verghese N, et al. Health-related quality of life and its socio-economic and cultural predictors among advanced cancer patients: evidence from the APPROACH cross-sectional survey in Hyderabad-India. *BMC Palliat Care* 2019; 18(1): 94. DOI: 10.1186/s12904-019-0465-y
 49. Kroll JL, Kim S, Cho D, et al. Financial distress and its associated burden in couples coping with an advanced cancer. *Support Care Cancer* 2022; 30: 4485–4495.
 50. Lam K, Zeng L, Zhang L, et al. Predictive factors of overall well-being using the EORTC QLQ-C15-PAL extracted from the EORTC QLQ-C30. *J Palliat Med* 2013; 16: 402–408.
 51. Malhotra C, Bundoc F, Ang FJL, et al. Financial difficulties and patient-reported outcomes among patients with advanced heart failure. *Qual Life Res* 2021; 30: 1379–1387.
 52. Malhotra C, Harding R, Teo I, et al. Financial difficulties are associated with greater total pain and suffering among patients with advanced cancer: results from the COMPASS study. *Support Care Cancer* 2020; 28: 3781–3789.
 53. Mercadante S, Aielli F, Adile C, et al. Financial distress and its impact on symptom expression in advanced cancer patients. *Support Care Cancer* 2021; 29: 485–490.
 54. Pérez-Cruz PE, Langer P, Carrasco C, et al. Spiritual pain is associated with decreased quality of life in advanced cancer patients in palliative care: an exploratory study. *J Palliat Med* 2019; 22: 663–669.
 55. Püsküllüoğlu M, Zygulska AL, Tomaszewska IM, et al. Evaluation of health-related quality of life and its main influencing factors in a Polish population of patients with bone metastases. *Curr Prob Cancer* 2016; 40: 183–197.
 56. Rakic M, Escher M, Elger BS, et al. Feelings of burden in palliative care: a qualitative analysis of medical records. *J Palliat Care* 2018; 33: 32–38.
 57. Rattani SA, Dahlke S and Cameron B. Cancer care in Pakistan: a descriptive case study. *Glob Qual Nurs Res* 2022; 9: 23333936221080988. DOI: 10.1177/23333936221080988
 58. Rosenzweig M, West M, Matthews J, et al. Financial toxicity among women with metastatic breast cancer. *Oncol Nurs Forum* 2019; 46: 83–91.
 59. Segerstrom SC, Kasarskis EJ, Fardo DW, et al. Socioemotional selectivity and psychological health in amyotrophic lateral sclerosis patients and caregivers: a longitudinal, dyadic analysis. *Psychol Health* 2019; 34: 1179–1195.
 60. Sudarisan SSP, Abraham B and George C. Prevalence, correlates of depression, and its impact on quality of life of cancer patients attending a palliative care setting in South India. *Psychooncology* 2019; 28: 1308–1313.
 61. Tada T, Hashimoto F, Matsushita Y, et al. Investigation of QOL of hospice patients by using EORTC-QLQ-C30 questionnaire. *J Med Invest* 2004; 51: 125–131.
 62. Tseng WT, Lee Y, Hung CF, et al. Stigma, depression, and anxiety among patients with head and neck cancer. *Support Care Cancer* 2022; 30: 1529–1537.
 63. Tu MS and Chiou CP. Perceptual consistency of pain and quality of life between hospice cancer patients and family caregivers: a pilot study. *Int J Clin Pract* 2007; 61: 1686–1691.
 64. Wang SE, Haupt EC, Nau C, et al. Association between financial distress with patient and caregiver outcomes in home-based palliative care: a secondary analysis of a clinical trial. *J Gen Intern Med* 2022; 37: 3029–3037.
 65. Gupta D, Lis CG and Grutsch JF. Perceived cancer-related financial difficulty: implications for patient satisfaction with quality of life in advanced cancer. *Support Care Cancer* 2007; 15: 1051–1056.
 66. Aaronson NK, Ahmedzai S, Bergman B, et al. The European organization for research and treatment of cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; 85: 365–376.
 67. Quinn S, Richards N and Gott M. Dying at home for people experiencing financial hardship and deprivation: how health and social care professionals recognise and reflect on patients' circumstances. *Palliat Care Soc Pract* 2023; 17: 26323524231164162. DOI: 10.1177/26323524231164162
 68. Marmot M, Allen J, Boyce T, et al. *All together Fairer: health equity and the social determinants of health in Cheshire and Merseyside*. London: Institute of Health Equity, 2022.
 69. University of Glasgow. *Money matters at the end of life. Having open conversations about financial hardship at the end of life*, https://www.gla.ac.uk/media/Media_996697_smx.x.pdf (2023, accessed 24 April 2024).
 70. Hobson F. *Social security (special rules for end of life) bill*, <https://commonslibrary.parliament.uk/research-briefings/cbp-9580/> (2022, accessed 24 April 2024).
 71. Department of Health and Human Services. Centers for medicare & medicaid services. <https://www.medicare.gov/Pubs/pdf/02154-medicare-hospice-benefits.pdf> (2023, accessed 15 April 2024).
 72. Pickett K and Wilkinson R. *The spirit level: why equality is better for everyone*. London: Penguin, 2010.