

SPECIAL ARTICLE

Cutaneous melanoma: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up[☆]

T. Amaral¹, M. Ottaviano², A. Arance³, C. Blank^{4,5,6}, V. Chiarion-Sileni⁷, M. Donia⁸, R. Dummer⁹, C. Garbe¹⁰, J. E. Gershenwald¹¹, H. Gogas¹², M. Guckenberger¹³, J. Haanen^{14,15,16}, O. Hamid¹⁷, A. Hauschild¹⁸, C. Höller¹⁹, C. Lebbé²⁰, R. J. Lee^{21,22}, G. V. Long^{23,24,25}, P. Lorigan^{21,22}, E. Muñoz Couselo²⁶, P. Nathan²⁷, C. Robert²⁸, E. Romano²⁹, D. Schadendorf^{30,31}, V. Sondak³², K. P. M. Suijkerbuijk³³, A. C. J. van Akkooi^{23,34}, O. Michelin^{35†} & P. A. Ascierto^{36†}, on behalf of the ESMO Guidelines Committee*

¹Skin Cancer Clinical Trials Center—University of Tuebingen, Tuebingen, Germany; ²Department of Melanoma, Cancer Immunotherapy and Development Therapeutics, Istituto Nazionale Tumori IRCCS Fondazione “G. Pascale”, Napoli, Italy; ³Department of Medical Oncology and IDIBAPS, Hospital Clinic y Provincial de Barcelona, Barcelona, Spain; ⁴Department of Medical Oncology and Division of Immunology, The Netherlands Cancer Institute Antoni van Leeuwenhoek Ziekenhuis (NKI), Amsterdam; ⁵Leiden University Medical Center (LUMC), Leiden, The Netherlands; ⁶University Clinic Regensburg, Regensburg, Germany; ⁷Department of Oncology, Melanoma Unit, Istituto Oncologico Veneto, IOV-IRCCS, Padova, Italy; ⁸National Center for Cancer Immune Therapy (CCIT-DK), Department of Oncology, Copenhagen University Hospital Herlev and Gentofte, Herlev, Denmark; ⁹Department of Dermatology, Skin Cancer Center, USZ—University Hospital Zürich, University of Zürich, Zürich, Switzerland; ¹⁰Department of Dermatology, Center for DermatoOncology, University Hospital Tuebingen, Tuebingen, Germany; ¹¹Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center and The University of Texas MD Anderson Cancer Center UTHealth Houston Graduate School of Biomedical Sciences, Houston, USA; ¹²First Department of Medicine, School of Medicine, National and Kapodistrian University of Athens—School of Medicine, Athens, Greece; ¹³Department of Radiation Oncology, University Hospital Zürich, University of Zürich, Zürich, Switzerland; ¹⁴Division of Medical Oncology, Netherlands Cancer Institute, Amsterdam; ¹⁵Department of Medical Oncology, Leiden University Medical Center (LUMC), Leiden, The Netherlands; ¹⁶Oncology Service, Melanoma Clinic, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland; ¹⁷Medical Oncology, Cutaneous Malignancies, The Angeles Clinic and Research Institute, A Cedars Sinai Affiliate, Los Angeles, USA; ¹⁸Department of Dermatology, UKSH—Universitätsklinikum Schleswig-Holstein—Campus Kiel, Kiel, Germany; ¹⁹Department of Dermatology, Medical University of Vienna, Vienna, Austria; ²⁰Université Paris Cite, AP-HP Dermato-oncology and CIC, Cancer Institute APHP, Nord Paris Cité, INSERM U976, Saint Louis Hospital, Paris, France; ²¹Department of Medical Oncology, The Christie NHS Foundation Trust, Manchester; ²²Faculty of Biology Medicine and Health, The University of Manchester, Manchester, UK; ²³Melanoma Institute Australia, The University of Sydney, Sydney; ²⁴Department Medical Oncology, Royal North Shore Hospital, Sydney; ²⁵Mater Hospital, Sydney, Australia; ²⁶Department of Medical Oncology, Vall d’Hebron University Hospital and Vall d’Hebron Institute of Oncology (VHIO), Barcelona, Spain; ²⁷Mount Vernon Cancer Centre, Northwood, UK; ²⁸Department of Oncology, Institut Gustave Roussy and Paris-Saclay University, Villejuif; ²⁹Department of Oncology, Center for Cancer Immunotherapy, Institut Curie, Paris, France; ³⁰Department of Dermatology, WTZ—Westdeutsches Tumorzentrum Essen, National Center for Tumor Diseases (NCT-West), Campus Essen, Essen; ³¹University Alliance Ruhr, Research Center One Health, University Duisburg-Essen, Essen, Germany; ³²Department of Cutaneous Oncology, Moffitt Cancer Center, Tampa, USA; ³³Department of Medical Oncology, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands; ³⁴Department of Melanoma and Surgical Oncology, Royal Prince Alfred Hospital, Sydney, Australia; ³⁵Department of Oncology, Geneva University Hospital, Geneva, Switzerland; ³⁶Melanoma, Cancer Immunotherapy and Development Therapeutics Unit, Istituto Nazionale Tumori IRCCS Fondazione Pascale, Napoli, Italy



Available online 14 November 2024

Key words: cutaneous melanoma, diagnosis, ESMO Clinical Practice Guideline, prognosis, risk assessment, treatment

INTRODUCTION

This European Society for Medical Oncology (ESMO) Clinical Practice Guideline (CPG) focuses on invasive cutaneous melanoma. The management of uveal melanoma and non-melanoma skin cancers are described in separate ESMO CPGs. Mucosal melanoma is described in the

Supplementary Material Section 1 and Table S1, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

INCIDENCE AND EPIDEMIOLOGY

Details on incidence and epidemiology of cutaneous melanoma are provided in the Supplementary Material Section 2, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

DIAGNOSIS AND PATHOLOGY/MOLECULAR BIOLOGY

Diagnosis

Details on the diagnostic work-up of cutaneous melanoma are provided in the Supplementary Material Section 3 and Tables S2 and S3, and a proposed algorithm is shown in

*Correspondence to: ESMO Guidelines Committee, ESMO Head Office, Via Geneva 4, CH-6900 Lugano, Switzerland
E-mail: clinicalguidelines@esmo.org (ESMO Guidelines Committee).

†Co-last authors.

[☆]Note: Approved by the ESMO Guidelines Committee: February 2002, last update November 2024. This publication supersedes the previously published version—*Ann Oncol*. 2019;30(12):1884-1901.

0923-7534/© 2024 European Society for Medical Oncology. Published by Elsevier Ltd. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Supplementary Figure S1, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Molecular characterisation

Testing for actionable mutations is recommended for patients with resectable or unresectable stage III or IV melanoma. Mutation testing should be considered for high-risk, clinical stage IIB-IIC melanoma, but is not routinely recommended for stage I or IIA disease. Mutation testing of *BRAF* V600 is mandatory, whereas testing for other *BRAF* mutations is optional. A full list of *BRAF* mutations by class is provided in Supplementary Table S4, available at <https://doi.org/10.1016/j.annonc.2024.11.006>. Testing can be offered for *NRAS* and *c-KIT* mutations; testing for *NTRK* alterations is recommended in the absence of *BRAF* or *RAS* mutations [see ESMO Scale for Clinical Actionability of molecular Targets (ESCAT) for further details—Supplementary Table S5, available at <https://doi.org/10.1016/j.annonc.2024.11.006>]. Mutation analysis using next-generation sequencing can be offered for unresectable melanoma. Mutation analysis must be carried out in accredited (certified) institutes that have careful quality controls and appropriate bioinformatic knowledge.

The main melanoma subtypes are associated with different mutational landscapes,¹ as shown in Supplementary Table S6, available at <https://doi.org/10.1016/j.annonc.2024.11.006>. In addition to the mutational status, reporting programmed death-ligand 1 (PD-L1) expression by immunohistochemistry is recommended for all unresectable stage III and IV melanoma, since the European Medicines Agency (EMA) has approved the administration of nivolumab—relatlimab only in patients with tumour cell PD-L1 expression <1%.² Tumour mutational burden computed on full exome sequencing or on a large full-length panel and expressed as the number of mutations per megabase can be assessed and recorded, but its clinical use is currently not warranted.¹

Signatures combining different gene panels and clinical/pathological characteristics (e.g. AMBLor,³ DecisionDx-Melanoma,⁴ Immunoprint,⁵ MelaGenix,⁴ SkylineDx⁶) have shown promising prognostic value in primary cutaneous melanoma. However, current evidence does not support the use of these tests in routine clinical practice. Prospective studies are underway to assess their predictive and prognostic value, which will provide valuable insights into their potential future clinical utility.⁷

Recommendations

- Dermoscopy by an experienced physician is recommended and increases the diagnostic accuracy [II, A].
- Diagnosis should be based on a full thickness complete excision with a minimal margin of clinically uninvolved skin [II, A]. For larger lesions where complete excision is not possible without reconstructive surgery, a biopsy can be taken [II, C].
- The histology report should include at least information on the type of melanoma, maximum vertical tumour

thickness in millimetres (Breslow, measured to the nearest 0.1 mm), presence of ulceration, microsatellites, lymphovascular invasion, neurotropism/perineural invasion, tumour-infiltrating lymphocytes (TILs), presence of regression and presence or absence of tumour at the deep and peripheral edges of the biopsy [II, A]. Mitotic rate should also be reported [III, B].

- A report on the wide excision should also be made available for complete pathological characterisation [II, A].
- Testing for actionable mutations is recommended in patients with resectable or unresectable stage III or IV melanoma [I, A] and should be considered in clinical stage IIB-IIC [V, C] but not for stage I or IIA disease [V, D].
 - *BRAF* V600 testing is mandatory [I, A; ESCAT score: I-A].

STAGING AND RISK ASSESSMENT

Details on the staging and risk assessment of cutaneous melanoma are provided in the Supplementary Material Section 4 and Table S2, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Recommendations

- Staging should be according to the eighth edition of the American Joint Committee on Cancer (AJCC) TNM (tumour—node—metastasis) staging system (AJCC8) [II, A].
- Sentinel lymph node biopsy (SLNB) is not routinely recommended for patients with a melanoma of AJCC8 stage pT1a (e.g. with a tumour thickness <0.8 mm and no ulceration) [II, E].
- SLNB is not usually recommended but can be discussed in pT1a for special cases [e.g. 3 mitoses/mm², a positive deep margin or when Breslow thickness cannot be reliably determined (pTx)] [III, D].
- SLNB should be discussed with patients with a melanoma of AJCC8 stage pT1b (i.e. with a tumour thickness of 0.8–1.0 mm or with a tumour thickness of <0.8 mm with ulceration) [III, B].
- SLNB is recommended for all patients with clinically node-negative T2a or higher tumours according to AJCC8 criteria (>1.0 mm Breslow thickness) [I, A]. Whenever possible, wide excision of the primary tumour should be carried out at the same time.
- For patients with T3b, T4a and T4b melanoma who qualify for adjuvant therapy, omitting SLNB can be discussed with the patient, but the potential benefits of SLNB in terms of staging (especially in patients with a *BRAF*-mutated, thick primary stage IIB or IIC who could be upstaged to stage III melanoma) and locoregional control should also be discussed [V, C].
- Whole-body physical examination with special attention to the primary tumour residual intact component and other suspicious pigmented cutaneous lesions, tumour satellites, in-transit metastases (ITMs) and regional lymph nodes (LNs) is recommended [IV, A]. In tumour stages IIB or higher, ultrasound (US), computed tomography (CT) and/or positron emission tomography (PET)

scans and brain magnetic resonance imaging (MRI) are recommended to ensure proper tumour assessment [III, B].

MANAGEMENT OF LOCAL/LOCOREGIONAL DISEASE

Treatment of localised melanoma

Full depth, wide local excision (WLE) of primary tumours (with safety margins of 0.5 cm for *in situ* melanomas, 1 cm for tumours with a thickness of ≤ 2 mm and 2 cm for tumours thicker than 2 mm) is recommended (see [Supplementary Table S7](#), available at <https://doi.org/10.1016/j.annonc.2024.11.006>).⁸ Modifications, with reduced safety margins, or micrographic surgery, are acceptable for preservation of function in acral and facial melanomas, including lentigo maligna melanoma.

Definitive radiotherapy (RT) to the primary tumour can only be considered in carefully selected patients for local control when excision is not possible either due to severe patient comorbidities (e.g. very old age, end-stage cardiovascular disease, etc.), when the morbidity associated with the excision is considered unacceptable or when surgery is refused by the patient. Palliative RT and palliative surgery can be offered for local control of symptomatic lesions.

Treatment of locoregional melanoma

Proposed treatment algorithms for the primary treatment of locoregional melanoma are shown in [Figures 1 and 2](#).

For patients with a positive SLNB, complete LN dissection (CLND) or irradiation of regional LNs should not be carried out.⁹⁻¹² Before undertaking additional treatments, a detailed staging investigation that includes high-resolution imaging techniques, such as CT, PET/CT and brain MRI, is necessary to exclude distant metastases.¹³

Treatment of ITMs

A proposed algorithm for the management of ITMs is provided in [Figure 3](#).

Patients with resectable disease but a short disease-free interval (< 6 months), unresectable satellite, ITMs or inoperable primary tumours of the limbs without additional metastases should be treated with systemic therapy, preferentially anti-programmed cell death protein 1 (PD-1)-based therapy, as these patients are at risk for developing distant metastasis. Treatment of these patients should be discussed at a multidisciplinary tumour board.

Adjuvant RT

Details regarding adjuvant RT are provided in [Supplementary Material Section 5](#), available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Adjuvant immunotherapy for resectable stage II-IV disease

A summary of results from key trials in the adjuvant setting is shown in [Supplementary Table S8](#), available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Stage IIB-IIC disease. Considering the 10-year melanoma-specific survival (MSS) rate in stage IIB-IIC disease, and the high risk of recurrence in these patients,^{14,15} two trials investigating adjuvant therapy, with a primary endpoint of recurrence-free survival (RFS), were conducted.

In the KEYNOTE-716 trial, 976 patients (age ≥ 12 years) with completely resected AJCC8 pathological stage IIB-IIC melanoma received intravenous pembrolizumab 200 mg (2 mg/kg in paediatric patients) or placebo every 3 weeks (q3w) for 17 cycles or until disease recurrence or unacceptable toxicity. The estimated 36-month RFS rate was 76.2% for pembrolizumab and 63.4% for placebo [hazard ratio (HR) 0.62, 95% confidence interval (CI) 0.49-0.79], and the estimated 36-month distant metastasis-free survival (DMFS) rate was 84.4% for pembrolizumab versus 74.7% for placebo (HR 0.59, 95% CI 0.44-0.79).¹⁶ In June 2022, the EMA approved the use of adjuvant pembrolizumab for patients ≥ 12 years old with stage IIB-IIC melanoma following complete resection.

In the CheckMate 76K trial, patients with AJCC8 pathological stage IIB-IIC melanoma (i.e. similar to KEYNOTE-716 eligibility criteria) were randomised to receive either adjuvant nivolumab or placebo for up to 1 year or until disease recurrence or unacceptable toxicity. At a median follow-up of ~ 23 months, adjuvant nivolumab improved RFS versus placebo (HR 0.53, 95% CI 0.40-0.71). Higher RFS rates were observed for nivolumab regardless of disease stage or T category.¹⁷ The licensed indication for nivolumab as adjuvant therapy has been expanded to include patients with stage IIB-IIC melanoma.

Currently, no overall survival (OS) data are available from the KEYNOTE-716 and CheckMate 76K trials.

Based on these findings, clinicians can offer adjuvant anti-PD-1 treatment for patients with AJCC8 stage IIB-IIC disease after a detailed discussion with the patient to weigh the pros and cons of treatment benefit versus toxicity and a careful evaluation of clinical factors, including patient age, comorbidities, performance status (PS), reproductive potential, personal/family history of autoimmune disease and compliance in case of immune-related toxicity.

Resectable stage III disease. It is worth noting that entry criteria for most adjuvant trials in this setting were based on the AJCC seventh edition (AJCC7),¹⁸ and that stage IIIA disease defined by AJCC7 includes a higher-risk group of patients than stage IIIA defined by AJCC8, which also incorporates Breslow thickness into stage III disease (5-year MSS rates for AJCC7 versus AJCC8 stage IIIA disease are 78% versus 93%, respectively).¹⁵ Moreover, patients with AJCC7 stage IIIA disease were either excluded (CheckMate 238) or had to have > 1 mm tumour metastasis from the SLNB to be included. Therefore, for patients with AJCC8 stage IIIA and < 1 mm SLNB tumour burden, in the absence of prospective validation of the benefit of adjuvant therapy in this patient population, adjuvant therapy should not be considered as the standard treatment.¹⁹

Efficacy and safety data from prospective randomised controlled trials evaluating adjuvant treatment with




Figure 1. Proposed algorithm for the management of patients with pT1b-pT4b cN0 cM0 melanoma.

Purple: algorithm title; blue: systemic anticancer therapy; orange: surgery; white: non-treatment aspects.

c, clinical; DMFS, distant metastasis-free survival; EMA, European Medicines Agency; ESCAT, ESMO Scale for Clinical Actionability of molecular Targets; ESMO, European Society for Medical Oncology; FDA, Food and Drug Administration; LM, lentigo maligna; M, metastasis; MCBS, ESMO-Magnitude of Clinical Benefit Scale; N, node; NEB, no evaluable benefit; OS, overall survival; p, pathological; PD-1, programmed cell death protein 1; R1, microscopic tumour at the margin; RFS, recurrence-free survival; RT, radiotherapy; SLNB, sentinel lymph node biopsy; T, tumour; WG, working group; WLE, wide local excision.

ªRT can be considered for local tumour control in cases of inadequate resection margins of LM [III, B] and could be discussed for patients with an R1 resection [III, C]. Adjuvant RT to the primary excision site should be considered for patients with desmoplastic or neurotropic melanoma for whom adequate (≥ 8 mm) pathological resection margins cannot be achieved [IV, C].

ªTreatment discussions with the patient should include consideration of the RFS benefit but lack of mature OS data [I, A].

ªTreatment discussions with the patient should consider the DMFS and RFS benefits but lack of mature OS data compared with placebo [I, A].

ªESMO-MCBS v1.1⁷⁸ was used to calculate scores for therapies/indications approved by the EMA or FDA. The scores have been calculated and validated by the ESMO-MCBS WG and reviewed by the authors (<https://www.esmo.org/guidelines/esmo-mcbs/esmo-mcbs-evaluation-forms>).

ªTreatment discussions with the patient should consider the DMFS and RFS benefits and potential OS benefit for patients with BRAF V600E-mutated melanoma [I, A].

ªESCAT scores apply to genomic alterations only. These scores have been defined by the guideline authors and assisted as needed by the ESMO Translational Research and Precision Medicine Working Group.⁷⁷

immune checkpoint inhibitors (ICIs; ipilimumab, nivolumab, pembrolizumab) in patients with high-risk resected stage III melanoma are summarised below. It is important to note that the eligibility criteria for all trials except CheckMate 915 included complete resection of all disease, including primary tumour excision with adequate margins and CLND in patients with nodal metastases detected by SLNB and those with clinical evidence of regional disease. Although it is unclear if the recommended adjuvant treatment options have similar efficacy in the absence of CLND following a positive SLNB, the consistent 24-month RFS rates reported for nivolumab in CheckMate 238 and CheckMate 915²⁰ suggest that CLND should not be a factor in the decision to use adjuvant therapy in patients with nodal metastases detected by SLNB.

Anti-cytotoxic T lymphocyte-associated antigen 4. In the EORTC 18071 trial,²¹ long-term adjuvant therapy with the anti-cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) agent ipilimumab resulted in improved 5-year RFS (HR 0.76, 95% CI 0.64–0.89, $P < 0.001$) and OS rates (HR 0.72, 95% CI 0.58–0.88, $P = 0.001$) compared with placebo. The benefit was also observed for patients with N1b and higher disease stages. However, the treatment schedule (10 mg/kg q3w for four doses and then every 3 months for up to 3 years) was associated with several severe and long-lasting adverse reactions and is no longer used. Given the toxicity profile of anti-CTLA-4 and the duration of therapy investigated, adjuvant therapy with either anti-PD-1 agents or dabrafenib–trametinib for patients with BRAF-mutated melanoma is preferred.




Figure 2. Proposed algorithm for the management of patients with stage III melanoma and clinically positive LNs or resectable stage IV melanoma. Purple: algorithm title; turquoise: combination of treatments and treatment modalities; white: non-treatment aspects.

DMFS, distant metastasis-free survival; EMA, European Medicines Agency; ESMO, European Society for Medical Oncology; FDA, Food and Drug Administration; LN, lymph node; MCBS, ESMO-Magnitude of Clinical Benefit Scale; OS, overall survival; PD-1, programmed cell death protein 1; RFS, recurrence-free survival; RT, radiotherapy; WG, working group.

^aRT could be discussed for patients after resection of bulky LN metastases, especially if further surgical clearance is not feasible [III, C].

^bTreatment discussions with the patient regarding adjuvant anti-PD-1 therapy should consider the DMFS and RFS benefits but lack of mature OS data compared with placebo [I, A].

^cTreatment discussions with the patient regarding adjuvant targeted therapy should consider the DMFS and RFS benefits and potential OS benefit for patients with BRAF V600E-mutated melanoma [I, A].

^dNot EMA or FDA approved as neoadjuvant therapy.

^eTreatment discussions with the patient regarding neoadjuvant therapy should consider the EFS, DMFS and RFS benefits but lack of mature OS data [I, A].

^fESMO-MCBS v1.1⁷⁸ was used to calculate scores for therapies/indications approved by the EMA or FDA. The scores have been calculated and validated by the ESMO-MCBS WG and reviewed by the authors (<https://www.esmo.org/guidelines/esmo-mcbs/esmo-mcbs-evaluation-forms>).

Anti-PD-1. In the CheckMate 238 trial, adjuvant therapy with the anti-PD-1 agent nivolumab has shown a significant RFS benefit²² but no significant OS benefit compared with ipilimumab 10 mg/kg for patients with stage IIIB, IIIC or IV (AJCC7) resected melanoma with no evidence of disease (NED) (see details in 'Resectable stage III and IV NED' section below).²³

Adjuvant treatment with the anti-PD-1 agent pembrolizumab was evaluated in patients with AJCC7 stage IIIA (SLN >1 mm), IIIB or IIIC (without ITM) melanoma in the placebo-controlled EORTC 1325/KEYNOTE-054 trial. At a median follow-up of 4.9 years, the 5-year RFS rate was longer in the pembrolizumab group than in the placebo group in the intention-to-treat population (HR 0.61, 95% CI 0.51-0.72) and in those with PD-L1-positive tumours (HR 0.62, 95% CI 0.48-0.79). Moreover, the 5-year DMFS rate was higher in the pembrolizumab group than in the placebo group (HR 0.62, 95% CI 0.52-0.75).²⁴ OS data are not yet available.

These results were validated in the phase III S1404 trial, which compared adjuvant pembrolizumab with either of the two standard-of-care (SoC) options at the time (i.e.

high-dose interferon- α or ipilimumab 10 mg/kg). RFS was significantly longer in the pembrolizumab group than in the SoC group (HR 0.76, 99.62% CI 0.59-0.99, log-rank $P = 0.002$).²⁵

Based on these data, the EMA approved nivolumab and pembrolizumab for use in the adjuvant setting in August and December 2018, respectively.

Resectable stage III and IV NED. In the CheckMate 238 trial,²³ which compared nivolumab with ipilimumab 10 mg/kg in patients with AJCC7 stage IIIB, IIIC or IV resected melanoma, the 5-year RFS rates were 50% in the nivolumab group and 39% in the ipilimumab group (HR 0.72, 95% CI 0.60-0.86). The 5-year RFS rates according to disease stage (IIIB, IIIC, IV M1a-b, IV M1c) were 58%, 43%, 47% and 55% for nivolumab and 48%, 35%, 29% and 49% for ipilimumab. The 5-year DMFS and OS rates were 58% versus 51% (HR 0.79, 95% CI 0.63-0.99) and 76% versus 72% (HR 0.86, 95% CI 0.66-1.12), respectively.²² Moreover, nivolumab treatment was associated with fewer grade 3/4 adverse events (AEs) compared with ipilimumab (14.4% versus 45.9%, respectively).²³




Figure 3. Proposed algorithm for the management of patients with ITMs.

Purple: algorithm title; turquoise: combination of treatments and treatment modalities; white: non-treatment aspects.

DFI, disease-free interval; DMFS, distant metastasis-free survival; ECT, electrochemotherapy; EMA, European Medicines Agency; ESMO, European Society for Medical Oncology; FDA, Food and Drug Administration; ILI, isolated limb infusion; ILP, isolate limb perfusion; ITM, in-transit metastasis; MCBS, ESMO-Magnitude of Clinical Benefit Scale; OS, overall survival; PD-1, programmed cell death protein 1; RFS, recurrence-free survival; RT, radiotherapy; T-VEC, talimogene laherparepvec; WG, working group.

^aFor anti-PD-1-based therapy, treatment discussions with the patient should consider the DMFS and RFS benefits but lack of mature OS data compared with placebo [I, A]. For dabrafenib–trametinib, these discussions should also consider the DMFS and RFS benefits and potential OS benefit for patients with *BRAF* V600E-mutated melanoma [I, A].

^bNot EMA or FDA approved as neoadjuvant therapy.

^cESMO-MCBS v1.1⁷⁸ was used to calculate scores for therapies/indications approved by the EMA or FDA. The scores have been calculated and validated by the ESMO-MCBS WG and reviewed by the authors (<https://www.esmo.org/guidelines/esmo-mcbs/esmo-mcbs-evaluation-forms>).

In the IMMUNED phase II trial, 167 patients with stage IV melanoma and NED after surgery or RT were randomised to receive nivolumab 1 mg/kg plus ipilimumab 3 mg/kg q3w for four doses followed by nivolumab 3 mg/kg every 2 weeks (q2w), nivolumab monotherapy (3 mg/kg q2w) or matching placebo for up to 1 year. The HR for RFS for nivolumab–ipilimumab versus placebo was 0.25 (97.5% CI 0.13–0.48, $P < 0.0001$) and for nivolumab versus placebo was 0.60 (97.5% CI 0.36–1.00, $P = 0.024$). The HR (95% CI) for RFS according to disease stage (M1a–b and M1c) for nivolumab–ipilimumab versus placebo was 0.29 (0.15–0.55) and 0.18 (0.06–0.57), and for nivolumab versus placebo was 0.50 (0.29–0.86) and 0.87 (0.40–1.90). The HR for OS was significantly in favour of nivolumab–ipilimumab versus placebo (HR 0.41, 95% CI 0.17–0.99, $P = 0.040$) but not for nivolumab versus placebo (HR 0.75, 95% CI 0.36–1.56, $P = 0.44$). Rates of grade 3/4 treatment-related AEs (TRAEs) were numerically higher with combination therapy, but types of toxicity were similar to what is already known for these agents.²⁶

CheckMate 915 was a phase III, double-blind trial in resected stage IIIB–IIID or IV melanoma. Patients were randomised to receive nivolumab 240 mg q2w plus ipilimumab 1 mg/kg once every 6 weeks or nivolumab 480 mg every 4 weeks (q4w) for ≤ 1 year. There was no significant difference between treatment groups reported for median

RFS in the all-randomly assigned patient population (HR 0.92, 95% CI 0.77–1.09, $P = 0.269$) or in patients with PD-L1 tumour expression $< 1\%$ (HR 0.91, 95% CI 0.73–1.14), or in 2-year RFS rates according to disease stage (IIIB: HR 0.91, 95% CI 0.68–1.21; IIIC: HR 0.92, 95% CI 0.75–1.13; IIID: HR 1.61, 95% CI 0.70–3.67; IV: HR 0.88, 95% CI 0.58–1.32).²⁰

Currently, adjuvant treatment with nivolumab–ipilimumab is included as a treatment option in the National Comprehensive Cancer Network guideline for cutaneous melanoma in patients with stage IV melanoma and NED,²⁷ but this adjuvant treatment combination is not approved by the EMA or Food and Drug Administration (FDA).

Other systemic therapies are currently being evaluated as adjuvant therapy for patients with resected stage IIB–IV melanoma. In an ongoing phase II clinical trial, the addition of messenger RNA-4157 to adjuvant pembrolizumab has shown a 49% risk reduction in recurrence and/or death (HR 0.510, 95% CI 0.288–0.906, two-sided nominal $P = 0.019$) versus pembrolizumab alone.²⁸ A phase III trial is also ongoing.

Adjuvant targeted therapy for resectable stage II–III disease

The BRIM8 study evaluated single-agent vemurafenib versus placebo in patients with stage IIC and stage III (AJCC7

criteria) melanoma after complete surgical resection. The study did not meet its primary endpoint of DFS.²⁹ Therefore, BRAF inhibitor (BRAFi) monotherapy cannot be recommended as adjuvant treatment for melanoma.

The phase III COMBI-AD trial³⁰ included patients with resected AJCC7 stage III (SLN >1 mm) melanoma with *BRAF* V600E or V600K mutations who were randomised to receive either 12 months of adjuvant dabrafenib–trametinib or placebo. At the final analysis (>10 years of follow-up), the median OS was not reached (NR) in either arm (HR 0.80, 95% CI 0.62–1.01, $P = 0.06$). The estimated RFS (HR 0.52, 95% CI 0.43–0.63) and DMFS (HR 0.56, 95% CI 0.44–0.71) both favoured the dabrafenib–trametinib arm.³¹ In a subgroup analysis, patients with a tumour *BRAF* V600E mutation (91%) in particular appeared to derive benefit in terms of OS (HR 0.75, 95% CI 0.58–0.96) and RFS (HR 0.52). The *BRAF* V600K mutation subgroup did not appear to derive any survival benefit (HR 1.95, 95% CI 0.84–4.50), although patient numbers in this group were small and so definitive conclusions cannot be drawn. Translational and retrospective data suggest that patients with advanced-stage melanoma and *BRAF* V600K mutations derive a greater benefit from ICI therapy than BRAF-targeted therapy. Based on these results, adjuvant dabrafenib–trametinib is an SoC adjuvant treatment option for *BRAF* V600E-mutated stage III melanoma and is approved by the EMA.

Neoadjuvant and neoadjuvant plus adjuvant systemic therapy for resectable stage III melanoma and clinically or radiologically detectable LN metastasis

Prospectively planned treatment with neoadjuvant therapy followed by surgery and adjuvant therapy may also be referred to as perioperative therapy. This approach differs from prospectively planned neoadjuvant therapy alone, where any subsequent systemic therapy may be given depending on the pathological response.

Several early clinical trials investigated neoadjuvant and neoadjuvant plus adjuvant therapy, including ICIs alone or in combination, BRAFi–MEK inhibitor (MEKi) combination therapy and intralesional therapies alone or in combination, with the principal aim of studying the association between pathological response, RFS and OS. In a pooled analysis of data from 633 (77%) clinical trial patients and 185 (23%) real-world patients treated with ICI-based therapy, BRAFi–MEKi targeted therapy or ICI plus targeted therapy, a pathological complete response (pCR) or near pCR occurred in 55% of patients: 51% with targeted therapy, 58% with ICIs and 46% with ICI plus targeted therapy. In patients who achieved a pCR or near pCR, the 3-year RFS rates were 57% with targeted therapy, 93% with ICIs and 85% with ICI plus targeted therapy. In contrast, patients who achieved a pathological partial response (pPR) or pathological non-response (pNR) had 3-year RFS rates of 15% and 13% with targeted therapy, 79% and 41% with ICIs and 88% and 48% with ICI plus targeted therapy, respectively.³²

The randomised phase II Southwest Oncology Group (SWOG) S1801 trial included patients with histologically

confirmed, measurable, clinically detectable and resectable stage IIIB–IV cutaneous, acral and mucosal melanomas without brain metastases (BMs) who were randomised 1 : 1 to receive either adjuvant therapy (upfront surgery followed by 18 doses of pembrolizumab 200 mg q3w) or neoadjuvant plus adjuvant therapy (3 doses of neoadjuvant pembrolizumab followed by surgery and 15 doses of adjuvant pembrolizumab).³³ The majority of patients [288/313 (92%)] included in this trial had stage III disease. With a median follow-up of 14.7 months, event-free survival (EFS) was significantly longer with neoadjuvant plus adjuvant versus adjuvant therapy ($P = 0.004$ log-rank test); this EFS benefit was consistent across predefined subgroups.³³ There was no significant difference in OS (HR 0.63, 95% CI 0.32–1.24, one-sided $P = 0.091$), although OS data were immature at the time of reporting.³⁴ The AE rates were similar in both groups.³³ In patients who received neoadjuvant plus adjuvant therapy, 40% achieved a pCR.³⁵

NADINA was a phase III, randomised trial of neoadjuvant nivolumab–ipilimumab (two cycles) versus adjuvant nivolumab in 423 patients with biopsy-proven, resectable stage III melanoma involving LNs \pm a maximum of three ITMs.³⁶ In the neoadjuvant group, only patients who had a pPR or pNR received subsequent adjuvant treatment with either dabrafenib–trametinib (for *BRAF*-mutated melanoma) or nivolumab. The design and dosing schedule for NADINA were based on results from the two neoadjuvant trials that evaluated two cycles of nivolumab–ipilimumab (OpACIN-neo³⁷ and PRADO³⁸). At a median follow-up of 15.4 months, the estimated 18-month EFS and 18-month DMFS rates were 80.8% in the neoadjuvant group versus 53.9% in the adjuvant group (HR 0.32, 95% CI 0.22–0.48) and 85.7% in the neoadjuvant group versus 62.4% in the adjuvant group (HR 0.37, 95% CI 0.24–0.57), respectively. In the neoadjuvant group, 60.8% of patients had a major pathological response (MPR) defined according to International Neoadjuvant Melanoma Consortium (INMC) criteria (see below). The estimated 18-month RFS and 18-month DMFS rates were 93.1% and 96.9% for patients who had an MPR, 80.5% and 80.5% for those who had a pPR and 55.1% and 60.6% for those who had a pNR, respectively. These findings suggest that for patients who achieve an MPR after two cycles of neoadjuvant nivolumab–ipilimumab, further adjuvant therapy is not needed. However, longer follow-up is required to confirm these results.

PIVOTAL is a randomised phase III trial of neoadjuvant daromun (a combination of two antibody–cytokine fusions L19IL2 and L19TNF) followed by surgery versus upfront surgery in 256 patients with resectable stage III melanoma (60% of patients had cutaneous or subcutaneous metastases and 33% had received prior systemic therapy). At a median follow-up of 21.2 months, RFS [blinded independent central review (BICR) assessment] and DMFS were both significantly longer in the neoadjuvant treatment group (HR 0.59, 95% CI 0.41–0.86, log-rank $P = 0.005$ and HR 0.60, 95% CI 0.37–0.95, $P = 0.029$, respectively). Neoadjuvant therapy also resulted in a pCR rate of 21%. Daromun-related AEs were mostly local events, with

limited, low-grade systemic AEs and no autoimmune TRAEs recorded.³⁹ Given these data, daromun may be an option for patients with resectable stage III melanoma and cutaneous metastases, but further data are required.

A randomised phase II trial evaluated neoadjuvant talimogene laherparepvec (T-VEC) followed by surgery versus surgery alone in 150 patients with resectable stage IIIB-IV M1a melanoma. At a median follow-up of 32.1 months for the neoadjuvant group and 30.9 months for the surgery group, the 2-year RFS rates were 29.5% and 16.5%, respectively (HR 0.75, 80% CI 0.58-0.96). Neoadjuvant therapy was associated with a pCR rate of 17.1%.⁴⁰

The INMC was established with the aim of developing recommendations for investigating neoadjuvant therapy in melanoma to align future trial designs and correlative analyses.⁴¹ Although neoadjuvant therapy is not currently approved, it is reimbursed in some countries. Indeed, neoadjuvant therapy may be particularly beneficial in the following clinical situations:

- Patients with resectable stage III melanoma confined to the LNs, detectable by clinical or radiological assessment.
- Patients with resectable ITMs or oligometastatic stage IV disease.

Pathological response of patients who have undergone neoadjuvant treatment should be assessed based on guidance from the INMC (see: <https://melanoma-inc.org/>). Definitions of best pathological response are provided in [Supplementary Table S9](#), available at <https://doi.org/10.1016/j.annonc.2024.11.006>.⁴²

Recommendations

Treatment of localised melanoma

- Full depth, WLE of primary tumours with safety margins of 0.5 cm for *in situ* melanomas, 1 cm for tumours with a thickness ≤ 2 mm and 2 cm for tumours > 2 mm is recommended [III, B].

Treatment of locoregional melanoma

- Patients with pT1b-T4b cN0 cM0 melanoma and a positive SLNB should undergo standard follow-up [III, A] \pm systemic therapy according to disease stage (see [Figure 1](#)) [I, A].
- CLND is not recommended for patients with a positive SLNB [I, E].
- Patients with a negative SLNB can be offered standard follow-up [III, A], clinical trial participation [V, A] or anti-PD-1 therapy for 12 months (stages IIB-IIC) [I, A].
- Enrolment into a clinical trial wherever possible is preferred [V, A].

Treatment of ITMs

- Patients with resectable ITMs should undergo complete excision with clear margins [IV, B]. These patients can also be evaluated for neoadjuvant nivolumab–ipilimumab [I, A; ESMO-Magnitude of Clinical Benefit Scale (ESMO-MCBS) v1.1 score: A; not EMA or FDA

approved] followed by adjuvant therapy based on pathological response and *BRAF* status, neoadjuvant plus adjuvant pembrolizumab [II, A; not EMA or FDA approved] or adjuvant therapy [I, A].

- Patients with resectable disease but a short disease-free interval (< 6 months), unresectable satellite, ITMs or inoperable primary tumours of the limbs without additional metastases should be treated with systemic therapy [III, B].
- Patients with unresectable satellite or ITMs may be treated with systemic therapy with anti-PD-1 based immunotherapy or BRAFi–MEKi, according to *BRAF* mutation status [I, A].
 - Local therapy with T-VEC, [I, B; ESMO-MCBS v1.1 score: 3], isolated limb infusion or isolated limb perfusion [IV, C], RT [IV, C], electrochemotherapy [IV, C] or limited palliative excision [IV, C] can also be considered (no impact on OS).

Adjuvant RT

- Adjuvant RT is not routinely recommended [III, D].
- RT can be considered for local tumour control in cases of inadequate resection margins of lentigo maligna [III, B].
- Adjuvant RT to the primary excision site should be considered for patients with desmoplastic or neurotropic melanoma for whom adequate (≥ 8 mm) pathological resection margins cannot be achieved [IV, C].
- RT could be discussed for patients with an R1 resection (resection with microscopic tumour at the margin) or after resection of bulky LN metastases, especially if further surgical clearance is not feasible [III, C].

Adjuvant systemic therapy in stage IIB-IIC melanoma

- Adjuvant therapy with either pembrolizumab [ESMO-MCBS v1.1 score: A] or nivolumab [ESMO-MCBS v1.1 score: A] for 12 months should be considered for patients with stage IIB-IIC disease; treatment discussions with the patient should include consideration of the RFS benefit but lack of mature OS data [I, A].

Adjuvant systemic therapy in resected stage III and IV NED

- Adjuvant systemic therapy options are anti-PD-1 therapy (nivolumab for resected stage IIIB-IV NED [I, A; ESMO-MCBS v1.1 score: no evaluable benefit] or pembrolizumab for resected stage III [I, A; ESMO-MCBS v1.1 score: A]) or dabrafenib–trametinib for patients with resected stage III *BRAF* V600E-mutated melanoma [I, A; ESCAT score: I-A].
 - For anti-PD-1-based therapy, treatment discussions with the patient should consider the DMFS and RFS benefits but lack of mature OS data compared with placebo [I, A].
 - For dabrafenib–trametinib, these discussions should also consider the DMFS and RFS benefits and potential OS benefit for patients with *BRAF* V600E-mutated melanoma [I, A].
 - These treatments should be given within 12 weeks of complete resection [I, A].

- Targeted therapy should not be offered to patients with *BRAF* V600K-mutated melanoma in light of the potential detrimental effect on OS reported in the COMBI-AD trial [II, D].
- For patients with AJCC8 stage IIIA and <1 mm tumour burden, adjuvant systemic treatment is generally not recommended [I, D].
- The use of adjuvant nivolumab—ipilimumab is not recommended for resected stage III melanoma [I, D; not EMA or FDA approved].
- Patients with resectable stage IV melanoma can be offered systemic therapy [V, A], clinical trial [V, A] or metastasectomy or local ablative therapy [III, B] followed by adjuvant anti-PD-1 therapy [I, A].
 - The use of adjuvant nivolumab—ipilimumab according to the dosing schedule utilised in the phase II IMMUNED trial may be an option for selected patients with resected stage IV melanoma [II, C; not EMA or FDA approved].

Neoadjuvant and neoadjuvant plus adjuvant systemic therapy in resectable stage III melanoma and clinically or radiologically detectable LN metastasis

- For patients with resectable stage III melanoma and pathologically proven, clinically or radiologically detectable LN metastasis, neoadjuvant nivolumab—ipilimumab [ESMO-MCBS v1.1 score: A; not EMA or FDA approved] followed by surgery should be offered. For patients with an MPR defined according to INMC criteria, adjuvant treatment can be omitted. For patients without an MPR, further treatment should be discussed [I, A].
- Neoadjuvant plus adjuvant pembrolizumab is also recommended for these patients [II, A; not EMA or FDA approved].
- Treatment discussions with the patient regarding neoadjuvant therapy should consider the EFS, DMFS and RFS benefits but lack of mature OS data [I, A].

MANAGEMENT OF ADVANCED/METASTATIC DISEASE

Some patients with stage IV melanoma present with resectable disease. Although the value of complete surgical resection in such a clinical setting has not been validated in phase III prospective clinical trials, data from phase II trials are available.⁴³ Surgery remains an option for selected patients, preferentially combined with adjuvant or neoadjuvant systemic therapies and in a clinical trial setting.

Treatment of unresectable stage III and IV melanoma

Despite the improvements in OS with currently available systemic treatments (ICIs and targeted therapy), many questions remain unanswered, with resistance still a challenge; therefore, inclusion in clinical trials is a priority in all settings whenever possible.

Proposed algorithms for the management of unresectable stage III and IV melanoma are provided in [Figures 3 and 4](#).

First-line treatment

First-line treatment selection depends on the strategy used in the neoadjuvant and/or adjuvant setting as well as the

BRAF mutational status of the disease. The current first-line SoC treatment options for unresectable stage III/IV melanoma are PD-1 blockade (nivolumab, pembrolizumab), PD-1 blockade combined with CTLA-4 blockade (nivolumab—ipilimumab), PD-1 blockade combined with lymphocyte activation gene-3 (LAG-3) blockade (nivolumab—relatlimab) and, for *BRAF* V600-mutated melanoma, BRAFi (vemurafenib, dabrafenib, encorafenib) combined with MEKi (cobimetinib, trametinib, binimetinib). For unresectable stage IIIB-IIIC or IV M1a disease (AJCC7 criteria), T-VEC is also an option (see 'Treatment of ITMs' section above). However, combining T-VEC with PD-1 blockade does not provide any additional clinical benefit. It is worth noting that data for treatments in the unresectable stage III and IV disease setting were generated before these therapies became available in the adjuvant setting for patients with stage II/III melanoma. Therefore, the benefits seen for patients who have received adjuvant therapy may be different from the data reported here. Prospective data regarding the optimal treatment strategy for patients with unresectable stage III or IV melanoma who have received prior adjuvant therapy are required.

Immunotherapy. The superiority of nivolumab over dacarbazine (DTIC) chemotherapy (ChT) for the first-line treatment of patients with *BRAF*-wild-type (WT) melanoma was demonstrated in the prospective randomised CheckMate 066 trial, with an HR for death of 0.42 (99.79% CI 0.25-0.73, $P < 0.001$) and an HR for death or progression of disease of 0.43 (95% CI 0.34-0.56, $P < 0.001$).⁴⁴ Superiority of PD-1 blockade (nivolumab, pembrolizumab) over ipilimumab was demonstrated in two prospective randomised trials, CheckMate 067 and KEYNOTE-006.^{45,46} After a minimum follow-up of 10 years, CheckMate 067 had an HR for death for nivolumab versus ipilimumab of 0.63 (95% CI 0.52-0.76)⁴⁷ and KEYNOTE-006 (patients included in the KEYNOTE-587 extension study only) had an HR for death for pembrolizumab (both dose arms combined) versus ipilimumab of 0.71 (95% CI 0.60-0.85).⁴⁸ Based on these trials, PD-1 blockade is now an SoC option for all patients, regardless of tumour *BRAF* status, in the first-line setting.

The benefit of adding ipilimumab to nivolumab was also assessed in the CheckMate 067 trial. Treatment with nivolumab—ipilimumab according to this trial's dosing regimen resulted in numerically higher response rates (RRs) and longer response durations, time to subsequent therapies, patients alive after stopping therapy, progression-free survival (PFS) and OS.⁴⁵ By study design, the two nivolumab-containing arms could not be compared. At a minimum follow-up of 10 years, the median OS was 71.9, 36.9 and 19.9 months in the nivolumab—ipilimumab, nivolumab-only and ipilimumab-only groups, respectively. Median MSS was NR, 49.4 and 21.9 months, respectively, and median duration of response (DoR) was NR (>120 months), 103.2 months and 19.2 months, respectively.⁴⁷ Grade 3-4 AEs were reported in 55.0%, 16.3% and 27.3% of patients in the nivolumab—ipilimumab, nivolumab-only and ipilimumab-only groups,




Figure 4. Proposed algorithm for the management of patients with stage III/IV melanoma according to prior adjuvant treatment received.

Purple: algorithm title; blue: systemic anticancer therapy; turquoise: combination of treatments and treatment modalities; white: non-treatment aspects.

BM, brain metastasis; BRAFi, BRAF inhibitor; ChT, chemotherapy; ctDNA, circulating tumour DNA; EMA, European Medicines Agency; ESCAT, ESMO Scale for Clinical Actionability of molecular Targets; ESMO, European Society for Medical Oncology; FDA, Food and Drug Administration; ICI, immune checkpoint inhibitor; LDH, lactate dehydrogenase; M, metastasis; MCBS, ESMO-Magnitude of Clinical Benefit Scale; MEKi, MEK inhibitor; NGS, next-generation sequencing; PD-1, programmed cell death protein 1; PD-L1, programmed death-ligand 1; PS, performance status; RT, radiotherapy; TIL, tumour-infiltrating lymphocyte; T-VEC, talimogene laherparepvec; WG, working group; WT, wild type.

^aPatients with metastatic melanoma should have metastases (preferably) or the primary tumour screened for the detection of *BRAF* V600 mutation [IV, A; ESCAT score: I-A]. If no tumour tissue is available, ctDNA may be an alternative [III, C].

^bEnrolment into a clinical trial is preferred wherever possible [V, A].

^cAdditional treatment options include palliative resection [IV, C], RT [IV, B] and/or T-VEC [I, C] for patients with symptomatic extracranial disease; and best supportive and palliative care for all patients [V, A]. Local therapies should also be considered for all patients throughout the disease course, including for resectable recurrence after (neo)adjuvant therapy and, where needed, to achieve local control, with access to tissue for NGS analysis providing the potential for personalised therapy.

^dImmediate contraindications to ICI include rapid progression, elevated LDH levels, comorbidities and any symptoms that preclude ICI use. In these situations, ICI therapy should be reconsidered as soon as the contraindications are resolved and ICI becomes a viable therapy option.

^eAbsolute contraindications to ICI should be based on a multidisciplinary assessment.

^fESMO-MCBS v1.1⁷⁸ was used to calculate scores for therapies/indications approved by the EMA or FDA. The scores have been calculated and validated by the ESMO-MCBS WG and reviewed by the authors (<https://www.esmo.org/guidelines/esmo-mcbs/esmo-mcbs-evaluation-forms>).

^gEMA approved for PD-L1 expression <1%, FDA approval is regardless of PD-L1 expression.

^hFor patients in whom the decision to treat with targeted therapy has been made, those who cannot receive a MEKi (e.g. due to cardiovascular comorbidities, a recent BM bleeding event, history of retinal detachment or other ophthalmological contraindications) can be offered encorafenib as monotherapy [II, B; not FDA or EMA approved].

ⁱESCAT scores apply to genomic alterations only. These scores have been defined by the guideline authors and assisted as needed by the ESMO Translational Research and Precision Medicine Working Group.⁷⁷

^jInduction targeted therapy followed by anti-PD-1 therapy is not EMA or FDA approved. The optimal duration of induction targeted therapy is currently unknown.

^kFor patients who do not require a rapid tumour response to therapy due to aggressive disease.

^lNot EMA or FDA approved for second-line use.

^mAn option for selected young, fit patients with stage IV M1a-c melanoma, PS 0, normal LDH, 1-3 prior treatments and who are able to tolerate TIL-related side-effects.

ⁿNot EMA or FDA approved.

respectively,⁴⁹ with no new safety signals observed in subsequent trial reports.⁵⁰

A phase IIIb/IV trial showed that an alternative dosing schedule of nivolumab-*ipilimumab* (nivolumab 3 mg/kg plus *ipilimumab* 1 mg/kg) was associated with reduced

toxicity.⁵¹ Therefore, this dosing schedule could be discussed for some frail patients. However, as this trial was designed to evaluate the safety (and not the efficacy) profile of this alternative dosing schedule, its general use cannot be recommended.

More recently, an improvement in PFS has been reported for relatlimab–nivolumab. In the phase II/III, multicentre, double-blind, randomised RELATIVITY-047 trial, relatlimab–nivolumab was administered as a fixed-dose q4w to patients with previously untreated unresectable or metastatic melanoma. A median PFS of 10.1 months was reached with relatlimab–nivolumab versus 4.6 months with nivolumab (HR 0.75, 95% CI 0.62–0.92, $P = 0.006$ log-rank test). The 12-month PFS rate was 47.7% with relatlimab–nivolumab versus 36% with nivolumab. PFS across key subgroups also favoured relatlimab–nivolumab over nivolumab. Grade 3/4 TRAEs occurred in 18.9% of patients in the relatlimab–nivolumab group versus 9.7% in the nivolumab group.⁵² In an updated analysis at a median follow-up of 19.3 months, the HR for PFS by BICR in patients with a PD-L1 expression of $<1\%$ ($n = 209$) and $\geq 1\%$ ($n = 147$) was 0.68 (95% CI 0.53–0.86) and 0.96 (95% CI 0.70–1.31), respectively. The OS HRs in these subsets were 0.78 (95% CI 0.59–1.04) and 0.84 (95% CI 0.57–1.24), respectively.⁵³ Based on these data, relatlimab–nivolumab can be considered as a first-line treatment option. For patients who need to discontinue relatlimab–nivolumab due to toxicity, continuation of anti-PD-1 monotherapy can be discussed. In July 2022, the EMA approved the use of relatlimab–nivolumab for the first-line treatment of adults and adolescents (≥ 12 years of age) with advanced melanoma and a PD-L1 tumour expression of $<1\%$.

Pembrolizumab–lenvatinib as first-line treatment for patients with unresectable or metastatic melanoma was evaluated in the phase III LEAP-003 trial. Despite a significant improvement in PFS observed for the combination in an early interim analysis, findings from a subsequent analysis showed no OS benefit and a significant increase in toxicity and the trial was subsequently discontinued.⁵⁴

Given this collective evidence, treatment decisions should be tailored based on several parameters, including prior (neo)adjuvant therapy received, timing of recurrence on/after adjuvant therapy, resectability status, suitability to receive ICI therapy and PD-L1 status. For patients eligible to receive ICI therapy, the treatment choice of single-agent PD-1 blockade versus nivolumab–ipilimumab or relatlimab–nivolumab should be individualised to each patient.

Targeted therapy. In case of *BRAF*-mutated melanoma, additional first-line options are provided by BRAFis and MEKis. Combined BRAFi–MEKi is superior to single-agent BRAFi in terms of RRs, PFS and OS.^{55–58} However, findings from a *post hoc* analysis suggest that encorafenib as monotherapy provides a similar OS benefit to combined BRAFi–MEKi (HR 0.93, 95% CI 0.73–1.18) and so could be an option for patients with contraindications to MEKis, although encorafenib is not EMA or FDA approved as monotherapy.⁵⁹ In addition to improved efficacy, skin-related side-effects and the incidence of squamous-cell carcinomas are reduced with the combination, although MEKis add specific toxicities (e.g. muscle, heart, eyes). Single-agent BRAFis should only be used in case of an absolute contraindication for MEKis.

Treatment selection. First-line treatment decisions between targeted therapies and immunotherapies have been evaluated in several prospective trials, with the aim of defining the best sequencing approach. The phase III DREAMSEQ trial included 265 patients with treatment-naïve *BRAF* V600-mutated metastatic melanoma, stratified by Eastern Cooperative Oncology Group PS 0 or 1 and lactate dehydrogenase (LDH) level; patients with untreated melanoma BMs (MBMs) were excluded. Patients were randomised 1 : 1 to receive either nivolumab–ipilimumab (arm A) or dabrafenib–trametinib (arm B), and at disease progression, patients in arm A received dabrafenib–trametinib (arm C) and patients in arm B received nivolumab–ipilimumab (arm D). The median DoR was significantly longer for arm A than for arm B (NR versus 12.7 months, $P < 0.001$). The PFS showed a trend in favour of arm A (log-rank $P = 0.054$). The 2-year OS rate for those starting in arm A was 71.8% and was 51.5% for those starting with arm B (log-rank $P = 0.010$). It is worth noting that only ~50% of patients who had disease progression in arm A or B were enrolled into arm C or D, respectively, since most died within 6 months of their initial disease progression, many due to MBMs. Also, given the significant difference in terms of 2-year OS rates between arms A and B, the data safety monitoring committee recommended the study be closed to accrual and for patients in arm B to be given the option to switch to arm D without disease progression. Given this, the initial question regarding the best therapeutic sequence was not completely answered. Overall, grade ≥ 3 toxicity was 60% in arm A and 52% in arm B. Grade 5 TRAEs included two patients in arm A and one in arm C.⁶⁰

In the randomised, three-arm, non-comparative phase II SECOMBIT trial, patients with previously untreated, metastatic *BRAF* V600-mutated melanoma were randomly assigned to arm A ($n = 69$; encorafenib–binimetinib until disease progression and then ipilimumab–nivolumab), arm B ($n = 71$; ipilimumab–nivolumab until disease progression and then encorafenib–binimetinib) or arm C ($n = 69$; encorafenib–binimetinib for 8 weeks followed by ipilimumab–nivolumab until disease progression and then encorafenib–binimetinib). At a median follow-up of 32.2 months, median OS was NR in any arm and >30 patients were alive in all arms. No new safety signals emerged.⁶¹ The 5-year OS rates were 45% in arm A, 52% in arm B and 57% in arm C.⁶²

The randomised, phase II EBIN trial evaluated a total of 2 years of therapy, either with immunotherapy (nivolumab–ipilimumab for four cycles followed by nivolumab; arm A) or encorafenib–binimetinib for 3 months followed by immunotherapy, as per arm A (arm B) in 271 patients with *BRAF*-V600E/K unresectable stage III/IV melanoma. At a median follow-up of 21 months, there was no PFS benefit associated with the addition of induction targeted therapy (HR 0.87, 90% CI 0.67–1.12, $P = 0.36$). However, findings from a prespecified subgroup analysis suggested a PFS benefit for induction targeted therapy among patients with LDH $>2\times$ the upper limit of normal (ULN; HR 0.46, 95% CI 0.21–1.03).

In a *post hoc* analysis, patients with liver metastases also benefited from the sequential design (HR 0.48, 95% CI 0.28-0.80).⁶³

Based on these results, first-line nivolumab—ipilimumab is the preferred treatment when this can be safely delivered for the first few months (i.e. when a rapid response is not required due to aggressive/symptomatic disease), with targeted therapies reserved for subsequent treatment lines. The optimal duration of induction targeted therapy and the best targeted therapy combination is currently unknown.

Oligometastatic disease. Oligometastatic disease is difficult to define. It is dependent on the number and localisation of metastatic sites and can be resectable or unresectable, but there is currently no consensus regarding its definition. Treatment options mimic those for patients with resectable stage IV melanoma and include systemic therapy, surgery, local treatment (stereotactic RT or ablative therapy) or a clinical trial (see Figure 4). However, based on results from trials evaluating neoadjuvant systemic therapy, the use of first-line systemic therapy instead of ablative therapy, even in patients with resectable oligometastatic disease, seems preferable and should be discussed. Data from clinical trials investigating this question are required.

Second-line treatment

Second-line treatment selection depends on the strategy used in the neoadjuvant, adjuvant and first-line metastatic setting as well as the *BRAF* mutational status of the disease, as illustrated in Figure 4. Clinical trials should always be the first choice, when available, based on an appropriate backbone regimen and comparator arm if randomised. At all time points for patients with oligometastatic disease, ablative therapy can be considered.

***BRAF*-WT melanoma.** For *BRAF*-WT melanoma, approved second-line options are very limited. Thus, consideration for clinical trials and/or personalised approaches is appropriate. If the first-line treatment was anti-PD-1 monotherapy or if patients had primary refractory disease following anti-PD-1 therapy, ipilimumab and ipilimumab—nivolumab are options based on results from the phase II SWOG S1616 trial.⁶⁴⁻⁶⁶ In this trial, treatment with ipilimumab—nivolumab was associated with a statistically significant improvement in PFS compared with ipilimumab alone (HR 0.63, 90% CI 0.41-0.97, one-sided $P = 0.04$).⁶⁶ Nivolumab—relatlimab might also represent an option after failure of single-agent anti-PD-1 therapy; in the phase I/IIa RELATIVITY-020 trial, objective response rates (ORRs) of 12.0% and 9.2% were reported for patients who had progressed after one or more than one prior anti-PD-1-containing regimen, respectively. Corresponding median PFS values were 2.1 and 3.2 months, respectively.⁶⁷

In some cases, such as (i) toxicity to anti-PD-1-based immunotherapy which precludes the use of second-line anti-PD-1-based therapy, (ii) rapidly progressing disease or

(iii) high tumour volume with symptomatic disease, clinical trials including bispecifics, T-cell engagers, etc. should be a preferred option. ChT with DTIC, carboplatin—paclitaxel, temozolomide or fotemustine can be discussed. However, none of these ChT regimens provide an OS advantage⁶⁸ and they are associated with a low RR, short PFS and increased toxicity in patients who have progressed on ICI therapy.⁶⁹

TIL therapy is another option which can be manufactured using different techniques. However, one of the limitations of this therapy is the time needed for manufacturing, which is currently 3-6 weeks, making it an option only for a selected group of patients. TIL therapy is currently not EMA approved for use in this setting.

In a phase II trial of TIL therapy, lifileucel (an autologous, centrally manufactured TIL product) demonstrated durable responses in patients with previously treated metastatic melanoma and limited treatment options. In this trial, patients received a non-myeloablative lymphodepletion regimen, a single infusion of lifileucel and up to six doses of high-dose interleukin-2 (IL-2). The ORR was 36%, with two complete responses (CRs) and 22 partial responses. The disease control rate (DCR) was 80% and the median DoR was NR after a median follow-up of 18.7 months. In the subset of patients with primary refractory disease following prior anti-PD-1 therapy, the ORR and DCR were 41% and 81%, respectively. The safety profile was consistent with AEs associated with non-myeloablative lymphodepletion and IL-2.⁷⁰

In an open-label phase III trial, 168 patients (86% with anti-PD-1-refractory disease) with unresectable stage IIIC-IV melanoma (AJCC7 criteria) were randomised 1 : 1 to receive either TILs (manufactured at each trial centre) or ipilimumab. Infusion of TILs was preceded by non-myeloablative, lymphodepleting ChT followed by high-dose IL-2. After a median follow-up of 33.0 months, median PFS was 7.2 months for the TIL group versus 3.1 months for the ipilimumab group (HR 0.50, 95% CI 0.35-0.72, $P < 0.001$). The ORR was 49% for the TIL group and 21% for the ipilimumab group, with 20% and 7% achieving a CR, respectively. Median OS was 25.8 months for the TIL group and 18.9 months for the ipilimumab group (HR 0.83, 95% CI 0.54-1.27, $P = 0.39$). Grade ≥ 3 TRAEs occurred in all TIL-treated patients and 57% of ipilimumab-treated patients.⁷¹ Currently, TIL therapy remains a highly toxic treatment option for selected patients who can tolerate its side-effects, administered within regional reference centres.⁷² Current clinical trial evidence suggests that those who derive most benefit are young patients with stage IV M1a-c melanoma, PS 0, normal LDH and one to three prior treatments.⁷³

In the phase II, LEAP-004 trial, pembrolizumab—lenvatinib was evaluated in patients progressing within 12 weeks of the last dose of an anti-PD-1 inhibitor, given as monotherapy or with other therapies, including CTLA-4 inhibitors. In the overall population, after a median follow-up of 15.3 months, the ORR was 21.4%, median DoR was 8.3 months, median PFS was 4.2 months and median OS

was 14.0 months.⁷⁴ Pembrolizumab—lenvatinib is not EMA or FDA approved for use in this setting.

BRAF-mutated melanoma. For *BRAF*-mutated melanoma, all the options available for *BRAF*-WT melanoma are still valid, with the addition of combined BRAFi—MEKi therapy if not already used as the immediate prior treatment. BRAFi—MEKi therapy after disease progression with first-line immunotherapy should be offered.

NRAS-mutated melanoma. For *NRAS*-mutated melanoma, due to the limited efficacy of MEKis, first-line immunotherapy options are identical to those for *NRAS*-WT melanoma. Binimetinib as a single agent, however, can be considered for patients who do not benefit from prior anti-PD-1 therapy, in accordance with findings from the NEMO trial, but there is no OS benefit and its use as monotherapy is not EMA approved.⁷⁵

Subsequent lines

Subsequent lines of therapy are not currently evidence based. Clinical trials or rechallenge, either with targeted therapy (for patients with *BRAF*-mutated melanoma) or immunotherapies, can be an option⁷⁶ (see Figure 4).

MBMs

A proposed algorithm for the management of patients with MBMs is provided in Figure 5. Details of management are provided in Supplementary Material Section 6, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Predictive and prognostic biomarkers

Information regarding potential predictive and prognostic markers in melanoma is provided in Supplementary Material Section 6, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Recommendations

General recommendations

- Patients with metastatic melanoma should have metastases (preferably) or the primary tumour screened for the detection of *BRAF* V600 mutation [IV, A; ESCAT score: I-A].
 - If no tumour tissue is available, circulating tumour DNA may be an alternative [III, C].
- Enrolment into a clinical trial wherever possible is preferred [V, A].
- In addition to the treatment options outlined below, palliative resection [IV, C] and/or RT [IV, B] and/or T-VEC [I, C] can be considered for symptomatic extracranial disease.

First-line treatment

- First-line treatment decisions must take into consideration prior neoadjuvant and/or adjuvant therapy received, timing of recurrence on/after adjuvant therapy,

resectability status and suitability to receive ICI therapy, as outlined in Figure 4 [V, A]. Primary or secondary resistance must be considered as this is also an eligibility criterion for clinical trials in pretreated patients [V, A].

- Patients with treatment-naïve resectable disease can be offered the following:
 - Stage III:
 - Wide excision of the primary tumour [III, B].
 - Neoadjuvant nivolumab—ipilimumab [I, A; ESMO-MCBS v1.1 score: A; not EMA or FDA approved] followed by adjuvant therapy based on pathological response and *BRAF* status.
 - Neoadjuvant plus adjuvant pembrolizumab [II, A; not EMA or FDA approved].
 - Adjuvant anti-PD-1 therapy [I, A] or dabrafenib—trametinib for *BRAF* V600E-mutated tumours [I, A].
 - Stage IV:
 - Clinical trial [V, A].
 - Metastasectomy or local ablative therapy [III, B] followed by adjuvant anti-PD-1 therapy [I, A].
 - Anti-PD-1 therapy alone [V, A].
- For patients with unresectable disease, first-line ipilimumab—nivolumab [ESMO-MCBS v1.1 score: A/4] is a preferred option for all patients regardless of *BRAF* status when this can be safely delivered for the first few months (i.e. when a rapid response is not required due to aggressive/symptomatic disease) [I, A].
- First-line nivolumab [I, A; ESMO-MCBS v1.1 score: A/4] or pembrolizumab [I, A; ESMO-MCBS v1.1 score: A/4] is also recommended.
 - Nivolumab—relatlimab can be offered as first-line treatment but EMA approval is only for patients with tumour cell PD-L1 expression <1% [I, B; ESMO-MCBS v1.1 score: 3; EMA approved for PD-L1 expression <1%, FDA approval is regardless of PD-L1 expression].
- If anti-PD-1-based therapy is not available or patients are considered ineligible for its use, BRAFi—MEKi combination therapy (dabrafenib—trametinib [ESMO-MCBS v1.1 score: 5]; vemurafenib—cobimetinib [ESMO-MCBS v1.1 score: A/5]; binimetinib—encorafenib [ESMO-MCBS v1.1 score: A/5]) is also an option in the first line for patients with *BRAF*-mutated melanoma [I; A; ESCAT score: I-A].
 - BRAFi—MEKi for 8-12 weeks followed by ipilimumab—nivolumab (as per SECOMBIT arm C or EBIN arm B) is also an option, especially for patients with high LDH levels and/or liver metastases [II, C; ESCAT score: I-A; induction targeted therapy is not EMA or FDA approved].
 - For patients in whom the decision to treat with targeted therapy has been made, those who cannot receive a MEKi (e.g. due to cardiovascular comorbidities, a recent BM bleeding event, history of retinal detachment or other ophthalmological contraindications) can be offered encorafenib as monotherapy [II, B; not FDA or EMA approved].




Figure 5. Proposed algorithm for the management of patients with MBMs.

Purple: algorithm title; blue: systemic anticancer therapy; turquoise: combination of treatments and treatment modalities; dark green, RT; white: non-treatment aspects.

BRAFi, BRAF inhibitor; ChT, chemotherapy; EMA, European Medicines Agency; ESCAT, ESMO Scale for Clinical Actionability of Molecular Targets; ESMO, European Society for Medical Oncology; FDA, Food and Drug Administration; LMD, leptomeningeal disease; MBM, melanoma brain metastasis; MEKi, MEK inhibitor; MRI, magnetic resonance imaging; RT, radiotherapy; SRS, stereotactic radiosurgery; WT, wild-type.

^aEnrolment into a clinical trial wherever possible is preferred [V, A].

^bNone of the systemic treatment options listed are EMA or FDA approved to treat MBMs.

^cIn patients where local treatment has been discounted due to the number and/or volume of MBMs, evaluate for the possibility of resection of dominant lesion(s).

^dEarly concurrent SRS may be preferred over late SRS as salvage treatment [IV, C]. Close monitoring with MRI is recommended so that SRS can be added when indicated [IV, B].

^eESCAT scores apply to genomic alterations only. These scores have been defined by the guideline authors and assisted as needed by the ESMO Translational Research and Precision Medicine Working Group.⁷⁷

- Patients with *BRAF*-mutated melanoma who have relapsed on or within 6 months of adjuvant BRAFi-MEKi therapy and who have an immediate or absolute contraindication to ICI can be offered the following first-line treatments:
 - BRAFi/MEKi (if >3 months after stopping adjuvant BRAFi/MEKi) [V, A].
 - ChT [II, C] (no OS benefit).
- First-line immunotherapy options for patients with *NRAS*-mutated melanoma are identical to those for patients with *NRAS*-WT disease [I, A].

Oligometastatic disease

- Treatment options for patients with oligometastatic disease mimic those for patients with resectable stage IV

melanoma and include systemic therapy [I, A], surgery [V, C], local treatment (stereotactic RT or ablative therapy [V, C]) or a clinical trial [V, A].

Second-line treatment

- Treatment options for the second-line setting depend on the therapy used in the first line and include ipilimumab-nivolumab [II, B], pembrolizumab [I, A; ESMO-MCBS v1.1 score: A/4], nivolumab [II, B], ipilimumab [II, B; ESMO-MCBS v1.1 score: 4] and BRAFi-MEKi combination therapy for patients with *BRAF*-mutated melanoma [II, B; ESCAT score: I-A]. Only pembrolizumab and ipilimumab monotherapy are EMA or FDA approved for second-line use.

- Nivolumab—relatlimab might also represent an option after failure of single-agent anti-PD-1 therapy [III, B; not EMA or FDA approved as second-line therapy].
- TIL therapy is an aggressive treatment option for selected patients (young, stage IV M1a-c melanoma, PS ≤ 1 , LDH $< 2 \times$ ULN and 1-3 prior treatments) who can tolerate its side-effects [II, B; not EMA or FDA approved].

Subsequent lines of treatment

- Third-line treatment rechallenge with the drug class (BRAFi—MEKi [IV, C] or ICI [IV, B]) not used in the immediate previous line can be considered, if feasible.
- If clinical trials, ICIs or BRAFis/MEKis are not available, ChT may be administered as later-line therapy [IV, C], with modest activity and no impact on OS.
- For patients with *NRAS*-mutated melanoma, binimetinib as a single agent can be offered to patients who do not benefit from prior anti-PD-1 therapy [III, C].

MBMs

- There are currently no systemic treatment options specifically approved for use in treating MBMs; enrolment into a clinical trial wherever possible is preferred [V, A].
- Patients with MBMs should be evaluated for stereotactic radiosurgery (SRS) [III, B]. Early concurrent SRS may be preferred over late SRS as salvage treatment [IV, C]. Since multiple sessions of SRS can be carried out at different time points of the disease course, close monitoring using MRI is recommended so that SRS can be added when indicated [IV, B].
- Patients with asymptomatic MBMs should preferably be treated upfront with nivolumab—ipilimumab [II, A].
 - If unsuitable for immunotherapy, patients with asymptomatic MBMs and *BRAF* V600-mutated melanoma can be offered BRAFi—MEKi [III, B; ESCAT I-A].
- For patients with symptomatic MBMs requiring steroids (< 10 mg/day prednisolone or equivalent):
 - BRAFi—MEKi if *BRAF* V600-mutated can be offered [III, A; ESCAT I-A].
 - Nivolumab—ipilimumab (*BRAF*-mutated or *BRAF*-WT) can be offered [III, A].
 - Neurosurgery should be discussed, especially if an accessible, resectable tumour is causing symptoms as this may render the patient asymptomatic and provide a bridging strategy to nivolumab—ipilimumab [IV, C].
- Patients with MBMs and neurological symptoms requiring steroids (> 10 mg prednisolone/day or equivalent) for whom local therapy is not an option can be considered for the following:
 - Clinical trial [V, A].
 - BRAFi—MEKi (if *BRAF* V600-mutated) [III, A; ESCAT I-A].
 - Ipilimumab—nivolumab [III, B].
 - ChT (if *BRAF*-WT) [IV, C].

- Patients with leptomeningeal disease can also receive the above therapies. Local treatment (RT [III, B] or intrathecal nivolumab [III, C]) can also be considered.
- Best supportive and palliative care should be discussed and activated for all patients with MBMs [V, A].

SELF-EXAMINATION, RISK ASSESSMENT AND FOLLOW-UP

A proposed algorithm for the follow-up of patients with melanoma is provided in Figure 6. Details of follow-up are provided in Supplementary Material Section 7, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.

Recommendations

- Patients with melanoma should be advised to avoid sunburn or unprotected solar exposure or any artificial UV exposure; lifelong regular self-examinations of the skin and peripheral LNs is also recommended [III, A].
- Patients must be made aware that family members have an increased melanoma risk [III, B].
- Follow-up should comprise a multidisciplinary approach, including oncologists, dermatologists and other specialties, as required by each individual patient's prior therapy and needs [V, A].
- During melanoma follow-up, patients should be clinically monitored with whole-body examinations, preferentially by a dermatologist, to detect relapse and to recognise additional skin tumours, especially secondary melanomas, as early as possible [III, B].
- There is no consensus on the optimal follow-up schedule or the utility of imaging and blood tests for patients with resected melanoma; respective national guidelines should be consulted, with adjustment as required, considering available resources, particularly after 3 years of follow-up [IV, B].
- The follow-up schedule should be tailored to each individual patient, considering the disease stage, individual risk and personal needs of the patient, and may include clinical—dermatological examination, LN US, laboratory examinations and imaging, as outlined in Figure 6 [V, B].

METHODOLOGY

This CPG was developed in accordance with the ESMO standard operating procedures for CPG development (<https://www.esmo.org/Guidelines/ESMO-Guidelines-Methodology>). All recommendations provided are based on current scientific evidence and the authors' collective expert opinion. Where recommendations for multiple different treatment options exist, prioritisation is illustrated by ordering these options according to: level of evidence (LoE) and grade of recommendation (GoR); where equal, by ESMO-MSBC score; where equal, by alphabetical order. The relevant literature has been selected by the expert authors. A table of ESCAT scores is included in Supplementary Table S5, available at <https://doi.org/10.1016/j.annonc.2024.11.006>.




Figure 6. Proposed algorithm for the follow-up of patients with melanoma.

Purple: algorithm title; white: non-treatment aspects.

CR, complete response; CT, computed tomography; ICI, immune checkpoint inhibitor; LN, lymph node; MDT, multidisciplinary team; mo, month; MRI, magnetic resonance imaging; NA, not applicable; NED, no evidence of disease; PET, positron emission tomography; Q, every; US, ultrasound; UV, ultraviolet; yr, year.

^aThe follow-up schedule should be tailored to each individual patient, considering the disease stage, individual risk and personal needs of the patient, and may include clinical–dermatological examination, LN US, laboratory examinations and imaging [V, B].

2024.11.006. ESCAT scores have been defined by the authors, assisted if needed by the ESMO Translational Research and Precision Medicine Working Group (WG).⁷⁷ A table of ESMO-MCBS scores is included in [Supplementary Table S10](#), available at <https://doi.org/10.1016/j.annonc.2024.11.006>. **2024.11.006.** ESMO-MCBS v1.1⁷⁸ was used to calculate scores for new therapies/indications approved by the EMA or FDA (<https://www.esmo.org/Guidelines/ESMO-MCBS>). The scores have been calculated and validated by the ESMO-MCBS WG and reviewed by the authors. The FDA/EMA or other regulatory body approval status of new therapies/indications is reported at the time of writing this CPG. LoEs and GoRs have been applied using the system shown in [Supplementary Table S11](#), available at <https://doi.org/10.1016/j.annonc.2024.11.006>.⁷⁹ Statements without grading were considered justified standard clinical practice by the authors. For future updates to this CPG, including eUpdates and Living Guidelines, please see the ESMO Guidelines website: <https://www.esmo.org/guidelines/guidelines-by-topic/melanoma>.

ACKNOWLEDGEMENTS

Manuscript editing support was provided by Claire Bramley and Ioanna Ntai (ESMO Guidelines staff) and Angela Corstorphine of Kstorfin Medical Communications Ltd (KMC); this support was funded by ESMO. Nathan Cherny, member of the ESMO-MCBS WG, and Urania Dafni, Giota Zygoura, Georgia Dimopoulou and Tereza Dellaporta, of Frontier Science Foundation Hellas, provided review and validation of the ESMO-MCBS scores. Nicola Latino and Francesca Chiovaro (ESMO Scientific Affairs staff) provided coordination and support of the ESMO-MCBS scores and Angela Corstorphine and Sian-Marie Lucas of KMC provided medical writing and editing support in the preparation of the ESMO-MCBS table; this support was funded by ESMO. Dr Joaquin Mateo (ESMO Translational Research and Precision Medicine Working Group) provided validation support for ESCAT scores. Dr Svetlana Jezdic (ESMO Medical Affairs staff) provided coordination and support of the ESCAT scoring.

FUNDING

No external funding has been received for the preparation of these guidelines. Production costs have been covered by ESMO from central funds.

DISCLOSURE

TA reports personal fees for advisory board membership from Delcath and Philogen; personal fees as an invited speaker from Bristol Myers Squibb (BMS), NeraCare, Novartis and Pierre Fabre; personal fees for a writing engagement from CeCaVa and Medtrix; institutional fees as local principal investigator (PI) from Agenus Inc., AstraZeneca, BioNTech, BMS, HUYA Bioscience, Immunocore, IO Biotech, Merck Sharp & Dohme (MSD), Pfizer, Philogen, Regeneron, Roche and University Hospital Essen; institutional fees as coordinating PI from Unicancer; institutional research grants from iFIT and Novartis; institutional funding from MNI—Naturwissenschaftliches und Medizinisches Institut, NeraCare, Novartis, Pascoe, Sanofi and Skyline-Dx; nonremunerated membership of the American Society of Clinical Oncology (ASCO) and the Portuguese Society for Medical Oncology; and role as clinical expert in the area of medical oncology for Infarmed. MO reports personal fees as an invited speaker from Amgen, BMS, MSD and Novartis; and nonremunerated membership of AIOM Associazione Italiana di Oncologia Medica, European Neuroendocrine Tumor Society (ENETS) and the IGG Italian Germ Cell Cancer Group. AA reports personal fees for advisory board membership from Almirall, BioNTech, BMS, MSD, Novartis, Pierre Fabre and Roche; institutional research funding from Amgen, BioNTech, BMS, MSD, Novartis, Replimune and Roche; and travel grants from BMS, MSD, Novartis and Pierre Fabre. CB reports institutional fees for advisory board membership from AstraZeneca, BMS, GenMab, GSK, Lilly, MSD, Novartis, Pfizer, Pierre Fabre and Roche; institutional funding as coordinating PI from 4SC, BMS, NanoString and Novartis; personal fees for expert testimony from Third Rock Ventures; personal stocks and shares from Flindr Therapeutics and Signature Oncology (intention to develop IFN signature algorithm); and a patent pending for WO 2021/177822 A1. VCS reports personal fees as an invited speaker for Immunocore and Pierre Fabre; and personal travel grants from Pierre Fabre and Sanofi. MD reports personal fees for an advisory role from Achilles Therapeutics; personal fees for an advisory role (not for pharmaceutical companies) from Alphasights and Guidepoint Global LLC; and a nonremunerated role as the Chairman of the Melanoma and Non-Melanoma Skin Cancer Scientific Committee of the Danish Medicines Council. RD reports personal fees for a consulting and/or an advisory role for Amgen, BMS, Catalym, MaviVAX SA, MSD, Novartis, Pfizer, Pierre Fabre, Regeneron, Roche, Sanofi, Second Genome, Simcere, Sun Pharma, T3 Pharma and Takeda. CG reports personal fees for advisory board membership from MSD and Philogen; personal fees for an advisory role from CeCaVa and NeraCare; personal fees for expert testimony from NeraCare; and a nonremunerated role as President of

the European Association of Dermato-Oncology (EADO). JEG reports personal fees for consultancy and advisory board/scientific advisory board/scientific advisory committee membership from Merck; royalties from intellectual property from UpToDate; nonremunerated membership of the executive committee of the American Cancer Surgery Standards Program, American College of Surgeons; nonremunerated leadership roles as Chair of the executive committee and overall Vice Chair of the American Joint Committee on Cancer (AJCC); and nonremunerated member of the medical advisory panel and Chair of the Melanoma Research Alliance Grant Review Committee. HG reports personal fees for advisory board membership from BMS, MSD, Pierre Fabre and Sanofi; personal fees as an invited speaker from BMS, MSD, Novartis, Pierre Fabre and Sanofi; institutional funding as local PI from Amgen, Bayer, BMS, Iovance, MSD and Replimmune; institutional research funding from AstraZeneca, BMS, Lilly, Pfizer and Pierre Fabre; and nonremunerated steering committee membership from Amgen and Replimmune. MG reports personal fees for advisory board membership from AstraZeneca; institutional funding as coordinating PI from Varian and ViewRay; institutional funding as trial Chair from AstraZeneca; nonremunerated membership of the Board of Directors for European Organisation for Research and Treatment of Cancer (EORTC) and the Swiss Academy of Multidisciplinary Oncology (SAMO); nonremunerated leadership role for the European Society for Radiotherapy and Oncology (ESTRO); and nonremunerated membership of the Radiation Oncology Science Council (ROSC). JH reports personal fees for advisory board membership from CureVac, Imcyse, Neogene Therapeutics, Sastra Cell Therapy, Scenic and Third Rock Venture; personal fees for stocks and shares from Neogene Therapeutics; institutional fees for advisory board membership from Achilles Therapeutics, AstraZeneca, BioNTech, BMS, Gadeta, Immunocore, Instil Bio, Iovance Biotherapeutics, Ipsen, Merck Serono, MSD, Molecular Partners, Novartis, Pfizer, PokeAcel, Roche, Sanofi and T-Knife; institutional research funding from Amgen, Asher Bio, BioNTech US, BMS, MSD, Novartis and Sastra Cell Therapy; nonremunerated member of the American Association for Cancer Research (AACR), ASCO and the Society for Immunotherapy of Cancer (SITC); a nonremunerated role as Editor-in-Chief for *Immuno-Oncology and Technology (IOTECH)*; and nonremunerated membership of the Editorial Boards for *ESMO Open* and *Kidney Cancer*. OH reports personal fees for advisory board membership from Alkermes, Amgen, Bactonix, BeiGene, Bioatla, BMS, Eisai, Georgiamune, GigaGen, Grit Bio, GSK, Idera, Immunocore, Incyte, Instil Bio, IO Biotech, Iovance, Janssen, KSQ Therapeutics, Merck, Novartis, Obsidian, Pfizer, Roche Genentech, Sanofi/Regeneron, Seagen, Tempus, Vial and Zelluna; personal fees as an invited speaker from BMS, Immunocore, Novartis, Pfizer and Sanofi/Regeneron; stocks and shares from Bactonix; institutional funding as local PI from Aduro, Akeso, Amgen, Arcus, Bioatla, BMS, CytomX, Exelixis, GSK, Idera, Immunocore, Incyte, Iovance, Merck, Merck Serono, Moderna, Nextcure, Novartis, Pfizer, Roche Genentech,

Sanofi/Regeneron, Seagen, Torque and Zelluna. AH reports personal fees for advisory board membership from Agenus, BMS, Dermagnostix, Eisai, Highlight Therapeutics, Immunocore, Incyte, IO Biotech, Merck Pfizer, MSD, NeraCare, Novartis, Philogen, Pierre Fabre, Regeneron, Replimune, Roche, Sanofi, Seagen, Xenthera; institutional funding as local PI from Agenus, Amgen, BMS, MSD, PierreFabre and Roche; and institutional funding as coordinating PI from Novartis and Regeneron. CH reports personal fees for advisory board membership from Ammirall, BMS, Immunocore, MSD, Novartis, Pierre Fabre, Sanofi and Ultimovacs; personal fees for invited speaker at advisory board from Amgen, BMS, MSD, Novartis, Pierre Fabre and Sanofi; institutional funding as coordinating PI from Amgen, BMS, Pierre Fabre and Roche; institutional funding as local PI from Immunocore, MSD, Nektar Therapeutics, Novartis, Pfizer, Pierre Fabre and Regeneron; and nonremunerated membership of EORTC and of the board of EADO. CL reports personal fees for advisory board membership from Amgen, BMS, InflaRx, Merck Serono, MSD, Novartis, Pierre Fabre, Roche and Sanofi; and travel grants from MSD and Pierre Fabre. RJL reports personal fees for advisory board membership from Pierre Fabre; personal fees as an invited speaker from AstraZeneca and Pierre Fabre; institutional research funding from AstraZeneca, BMS and Pierre Fabre; and institutional funding for an advisory role from Lumisphere Technology. GVL reports personal fees for an advisory role from Agenus Inc., Amgen Inc., Array Biopharma Inc., AstraZeneca, Bayer Healthcare Pharmaceuticals, BioNTech SE, Boehringer Ingelheim International GmbH, BMS, Evaxion Biotech A/S, GI Innovation, Hexal AG, Highlight Therapeutics S.L., Immunocore Ireland Limited, Innovent Biologics USA Inc., IOBiotech, MSD, Novartis Pharma AG, PHMR Limited, Pierre Fabre, Regeneron Pharmaceuticals Inc., Scancell Limited and SkylineDX B.V.; and a nonremunerated role as PI on over 30 clinical trials. PL reports personal fees for advisory board membership from MLA Dx, MSD and Ultimovacs; personal fees as an invited speaker from BMS and Pierre Fabre; institutional research funding from BMS and Pierre Fabre; travel expenses from MSD; and nonremunerated roles for Cancer Research UK (clinical research committee), EORTC Melanoma Group (Chairman); Melanoma Focus charity (invited speaker) and Society for Melanoma Research (SMR) Edinburgh 2022 (invited session chair). EMC reports personal fees for advisory board membership from BMS, Immunocore, MSD, Novartis, Pierre Fabre, Regeneron, Sanofi and Sun Pharma; personal fees as an invited speaker from BMS, MSD, Novartis, Pierre Fabre, Roche and Sanofi; and nonremunerated membership of the Board of Directors for the Spanish Group of Melanoma (GEM) and the Society of Spanish Oncology Group (SEOM). PN reports personal fees for advisory board membership from 4SC, BMS, IDEAYA, Immunocore, Merck, Novartis and Pfizer; personal fees as an invited speaker from Immunocore and Novartis; personal fees for steering committee membership from 4SC; institutional research funding from Immunocore; and institutional funding as coordinating PI from BMS, Immunocore, Ipsen, Merck, Novartis and Pfizer.

CR reports personal consultancy fees for advisory board membership from BMS, Egle Therapeutics, MaaT Pharma, Merck, MSD, Novartis, Pfizer, Philogen, Pierre Fabre, Regeneron, Roche, Sanofi, Sun Pharma and Ultimovacs; personal fees as an invited speaker from BMS, MSD, Novartis, Pierre Fabre and Sanofi; personal fees for steering committee membership from IO Biotech, Novartis, Pfizer and Regeneron; and personal fees as member of the Independent Data Monitoring Committee for Ultimovacs. ER reports personal fees for advisory board membership from Roche; institutional research fees from Amgen, AstraZeneca, BMS, Janssen and Replimune; a nonremunerated role as coordinating PI for Light Chain Biosciences; and a nonremunerated role as PI for AstraZeneca, BMS, Dragonfly Therapeutics, ImCheck Therapeutics, MSD/Merck, Pfizer and Roche. DS reports personal fees for advisory board membership from Anaveon, BioNTech, BMS, CureVac, Erasca, Immatics, Immunocore, MSD, NeraCare, Novartis, NoviGenix, Pfizer, Philogen, Pierre Fabre, Replimune, Sanofi/Regeneron, Sun Pharma and Ultimovacs; personal fees for expert testimony from AstraZeneca, BioAlta, Daiichi Sankyo, Formycon, InFlarX, PamGene and Seagen; personal fees as an invited speaker from BMS, Merck Serono, MSD, NeraCare, Sanofi and Sun Pharma; personal fees for steering committee membership from BMS and MSD; institutional research funding from BMS and MSD; institutional funding as coordinating PI from BMS, MSD, Novartis and Pierre Fabre; institutional funding as local PI from Philogen and Sanofi; nonremunerated steering committee membership for Novartis; nonremunerated membership of the Board of Directors for EORTC-MG; and nonremunerated leadership roles as a founding member and steering committee chair for the European Melanoma Registry (EuMelaReg). VS reports personal fees for advisory board membership, Independent Data Safety Monitoring Committee membership or independent medical monitoring from BMS, Genesis Drug Discovery & Development, Iovance, Merck, Mural Oncology and Novartis; institutional research funding as PI for Neogene Therapeutics (local) and Skyline DX (coordinating); and an institutional research grant from Turnstone Biologics. KPMS reports institutional fees for advisory board membership from AbbVie, BMS and Pierre Fabre; an institutional research grant from BMS, Genmab, Novartis, Philips and Tigris; and institutional honoraria for Safety Review Committee from Sairopa. ACJVA reports personal fees for advisory board membership from NeraCare and SkylineDx; institutional fees for advisory board membership from 4SC, Amgen, BMS, Merck-Pfizer, MSD, Novartis, Pierre Fabre, Provectus, Sanofi, Sirius Medical; and institutional research grants from Amgen (NIVEC study) and Merck-Pfizer. OM reports personal fees for an advisory role from Amgen, BMS, GSL, MSD, Novartis, Pierre Fabre and Roche; personal fees for membership of the Scientific Board of Cellula Therapeutics; personal fees from stocks and shares from Cellula Therapeutics; institutional fees for advisory board membership from Amgen, BMS, MSD, Novartis, Pierre Fabre and Roche; institutional fees as an invited speaker from Amgen, BMS, Pierre Fabre and Roche;

institutional funding from Amgen, BMS, MSD and Pierre Fabre; institutional research grants from Amgen, BMS, Merck, MSD and Pierre Fabre; institutional funding for a writing engagement from BMS; and a nonremunerated role as PI for Amgen, BMS, MSD, Novartis, Pierre Fabre and Roche. PAA reports personal fees for advisory board membership from Anaveon, Bayer and BioNTech; personal fees for a consultancy role from Bio-AI Health, BMS, Boehringer Ingelheim, Erasca, Immunocore, Italfarmaco, Lunaphore, Medicenna, Merck Serono, MSD, Nouscom, Novartis, Pfizer, Pfizer/Array, Philogen, Pierre Fabre, Regeneron, Replimmune, Roche Genentech, Sandoz, Sanofi, Sun Pharma and ValoTx; personal fees for an advisory role from BMS, Boehringer Ingelheim, Erasca, Immunocore, Merck Serono, MSD, Novartis, Pfizer/Array, Pierre Fabre, Regeneron, Replimmune, Roche, Sandoz, Sanofi and Sun Pharma; institutional funding for clinical trials from BMS, Pfizer/Array, Roche Genentech and Sanofi; travel grants from Bio-AI Health, MSD, Pfizer/Array and Replimmune; and nonremunerated roles as President of the Campania Society of ImmunoTherapy of Cancer (SCITO) Italy, President of the Fondazione Melanoma Onlus Italy, member of the steering committee for the Society for Melanoma Research (SMR), member of AIOM, ASCO, EORTC Melanoma Cooperative Group, SITC and SMR and member of the Board of Directors for the SITC (November 2017-December 2021).

REFERENCES

- Hayward NK, Wilmott JS, Waddell N, et al. Whole-genome landscapes of major melanoma subtypes. *Nature*. 2017;545(7653):1-18.
- European Medicines Agency. Opdualag summary of product characteristics. Published 2022. Available at https://www.ema.europa.eu/en/documents/product-information/opdualag-epar-product-information_en.pdf. Accessed June 20, 2023.
- Ewen T, Husain A, Stefanos N, et al. Validation of epidermal AMBRA1 and loricrin (AMBLor) as a prognostic biomarker for non-ulcerated AJCC stage I/II cutaneous melanoma. *Br J Dermatol*. 2024;190(4):549-558.
- Marchetti MA, Coit DG, Dusza SW, et al. Performance of gene expression profile tests for prognosis in patients with localized cutaneous melanoma: a systematic review and meta-analysis. *JAMA Dermatol*. 2020;156(9):953-962.
- Meyer S, Fuchs TJ, Bosserhoff AK, et al. A seven-marker signature and clinical outcome in malignant melanoma: a large-scale tissue-microarray study with two independent patient cohorts. *PLoS One*. 2012;7(6):e38222.
- Amaral T, Sinnberg T, Chatzioannou E, et al. Identification of stage I/II melanoma patients at high risk for recurrence using a model combining clinicopathologic factors with gene expression profiling (CP-GEP). *Eur J Cancer*. 2023;182:155-162.
- Kashani-Sabet M, Leachman SA, Stein JA, et al. Early detection and prognostic assessment of cutaneous melanoma: consensus on optimal practice and the role of gene expression profile testing. *JAMA Dermatol*. 2023;159(5):545-553.
- Long GV, Swetter SM, Menzies AM, et al. Cutaneous melanoma. *Lancet*. 2023;402(10400):485-502.
- Veronesi U, Adamus J, Bandiera DC, et al. Inefficacy of immediate node dissection in stage 1 melanoma of the limbs. *N Engl J Med*. 1977;297(12):627-630.
- Sim FH, Taylor WF, Ivins JC, et al. A prospective randomized study of the efficacy of routine elective lymphadenectomy in management of malignant melanoma. Preliminary results. *Cancer*. 1978;41(3):948-956.
- Balch CM, Soong SJ, Bartolucci AA, et al. Efficacy of an elective regional lymph node dissection of 1 to 4 mm thick melanomas for patients 60 years of age and younger. *Ann Surg*. 1996;224(3):255-263. discussion 263-256.
- Cascinelli N, Morabito A, Santinami M, et al. Immediate or delayed dissection of regional nodes in patients with melanoma of the trunk: a randomised trial. WHO Melanoma Programme. *Lancet*. 1998;351(9105):793-796.
- Gershenwald JE, Scolyer RA, Hess KR, et al. Melanoma of the skin. In: Amin MB, Edge S, Greene F, et al., editors. *AJCC Cancer Staging Manual*. 8th ed. New York: Springer International Publishing; 2017 p. 563-586.
- Garbe C, Keim U, Amaral T, et al. Prognosis of patients with primary melanoma stage I and II according to American Joint Committee on Cancer version 8 validated in two independent cohorts: implications for adjuvant treatment. *J Clin Oncol*. 2022;40(32):3741-3749.
- Gershenwald JE, Scolyer RA, Hess KR, et al. Melanoma staging: evidence-based changes in the American Joint Committee on Cancer eighth edition cancer staging manual. *CA Cancer J Clin*. 2017;67(6):472-492.
- Luke JJ, Ascierto PA, Khattak MA, et al. Pembrolizumab versus placebo as adjuvant therapy in resected stage IIB or IIC melanoma: final analysis of distant metastasis-free survival in the phase III KEYNOTE-716 study. *J Clin Oncol*. 2024;42(14):1619-1624.
- Kirkwood JM, Weber J, Hoeller C, et al. Adjuvant nivolumab versus placebo in resected stage IIB/C melanoma: update on CheckMate 76K. Philadelphia, Pennsylvania, USA: Paper presented at the Society for Melanoma Research 2023 International Congress; November 6-9, 2023.
- Balch CM, Gershenwald JE, Soong SJ, et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol*. 2009;27(36):6199-6206.
- Amaral T, Nanz L, Stadler R, et al. Isolated melanoma cells in sentinel lymph node in stage IIIA melanoma correlate with a favorable prognosis similar to stage IB. *Eur J Cancer*. 2024;201:113912.
- Weber JS, Schadendorf D, Del Vecchio M, et al. Adjuvant therapy of nivolumab combined with ipilimumab versus nivolumab alone in patients with resected stage IIB-D or stage IV melanoma (CheckMate 915). *J Clin Oncol*. 2023;41(3):517-527.
- Eggermont AMM, Chiarion-Sileni V, Grob J-J, et al. Prolonged survival in stage III melanoma with ipilimumab adjuvant therapy. *N Engl J Med*. 2016;375(19):1845-1855.
- Larkin J, Del Vecchio M, Mandalá M, et al. Adjuvant nivolumab versus ipilimumab in resected stage III/IV melanoma: 5-year efficacy and biomarker results from CheckMate 238. *Clin Cancer Res*. 2023;29(17):3352-3361.
- Weber J, Mandala M, Del Vecchio M, et al. Adjuvant nivolumab versus ipilimumab in resected stage III or IV melanoma. *N Engl J Med*. 2017;377(19):1824-1835.
- Eggermont AMM, Kicinski M, Blank CU, et al. Five-year analysis of adjuvant pembrolizumab or placebo in stage III melanoma. *NEJM Evid*. 2022;1(11):EVIDoA2200214.
- Grossmann KF, Othus M, Patel SP, et al. Adjuvant pembrolizumab versus IFN α 2b or ipilimumab in resected high-risk melanoma. *Cancer Discov*. 2022;12(3):644-653.
- Livingstone E, Zimmer L, Hassel JC, et al. Adjuvant nivolumab plus ipilimumab or nivolumab alone versus placebo in patients with resected stage IV melanoma with no evidence of disease (IMMUNED): final results of a randomised, double-blind, phase 2 trial. *Lancet*. 2022;400(10358):1117-1129.
- NCCN Clinical Practice Guidelines in Oncology Melanoma: Cutaneous. Version 2.2024—April 3, 2024. Available at https://www.nccn.org/professionals/physician_gls/pdf/cutaneous_melanoma.pdf. Accessed July 16, 2024.
- Weber JS, Khattak MA, Carlino MS, et al. Individualized neoantigen therapy mRNA-4157 (V940) plus pembrolizumab in resected melanoma: 3-year update from the mRNA-4157-P201 (KEYNOTE-942) trial. *J Clin Oncol*. 2024;42(suppl 17):LBA9512.
- Maio M, Lewis K, Demidov L, et al. Adjuvant vemurafenib in resected, BRAFV600 mutation-positive melanoma (BRIM8): a randomised,

- double-blind, placebo-controlled, multicentre, phase 3 trial. *Lancet Oncol.* 2018;19(4):510-520.
30. Long GV, Hauschild A, Santinami M, et al. Adjuvant dabrafenib plus trametinib in stage III BRAF-mutated melanoma. *N Engl J Med.* 2017;377(19):1813-1823.
 31. Long GV, Hauschild A, Santinami M, et al. Final results for adjuvant dabrafenib plus trametinib in stage III melanoma. *N Engl J Med.* 2024;391(18):1709-1720.
 32. Long GV, Blank CU, Amaria RN, et al. LBA41 Long-term survival with neoadjuvant therapy in melanoma: updated pooled analysis from the International Neoadjuvant Melanoma Consortium (INMC). *Ann Oncol.* 2024;35:S1232.
 33. Patel SP, Othus M, Chen Y, et al. Neoadjuvant-adjuvant or adjuvant-only pembrolizumab in advanced melanoma. *N Engl J Med.* 2023;388(9):813-823.
 34. Patel S, Othus M, Prieto V, et al. LBA6 Neoadjuvant versus adjuvant pembrolizumab for resected stage III-IV melanoma (SWOG S1801). *Ann Oncol.* 2022;33(suppl 7):S1408.
 35. Patel S, Othus M, Wright P, et al. LBA48 - Pathologic response and exploratory analyses of neoadjuvant-adjuvant versus adjuvant pembrolizumab (PEM) for resectable stage IIIb-IV melanoma from SWOG S1801. *Ann Oncol.* 2023;34(suppl 2):S1288.
 36. Blank CU, Lucas MW, Scolyer RA, et al. Neoadjuvant nivolumab and ipilimumab in resectable stage III melanoma. *N Engl J Med.* 2024;391(18):1696-1708.
 37. Rozeman EA, Menzies AM, van Akkooi ACJ, et al. Identification of the optimal combination dosing schedule of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma (OpACIN-neo): a multicentre, phase 2, randomised, controlled trial. *Lancet Oncol.* 2019;20(7):948-960.
 38. Reijers ILM, Menzies AM, van Akkooi ACJ, et al. Personalized response-directed surgery and adjuvant therapy after neoadjuvant ipilimumab and nivolumab in high-risk stage III melanoma: the PRADO trial. *Nat Med.* 2022;28(6):1178-1188.
 39. Hauschild A, Hassel JC, Ziemer M, et al. Phase 3 study (PIVOTAL) of neoadjuvant intralesional daromun vs. immediate surgery in fully resectable melanoma with regional skin and/or nodal metastases. *J Clin Oncol.* 2024;42(suppl 17):LBA9501.
 40. Dummer R, Gyorki DE, Hyngstrom J, et al. Neoadjuvant talimogene laherparepvec plus surgery versus surgery alone for resectable stage IIIB-IVM1a melanoma: a randomized, open-label, phase 2 trial. *Nat Med.* 2021;27(10):1789-1796.
 41. Amaria RN, Menzies AM, Burton EM, et al. Neoadjuvant systemic therapy in melanoma: recommendations of the International Neoadjuvant Melanoma Consortium. *Lancet Oncol.* 2019;20(7):e378-e389.
 42. Tetzlaff MT, Messina JL, Stein JE, et al. Pathological assessment of resection specimens after neoadjuvant therapy for metastatic melanoma. *Ann Oncol.* 2018;29(8):1861-1868.
 43. Sosman JA, Moon J, Tuthill RJ, et al. A phase 2 trial of complete resection for stage IV melanoma: results of Southwest Oncology Group Clinical Trial S9430. *Cancer.* 2011;117(20):4740-4806.
 44. Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med.* 2015;372(4):320-330.
 45. Wolchok JD, Chiarion-Sileni V, Gonzalez R, et al. Overall survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med.* 2017;377(14):1345-1356.
 46. Robert C, Schachter J, Long GV, et al. Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med.* 2015;372(26):2521-2532.
 47. Wolchok JD, Chiarion-Sileni V, Rutkowski P, et al. Final, 10-year outcomes with nivolumab plus ipilimumab in advanced melanoma. *N Engl J Med.* 2024. <https://doi.org/10.1056/NEJMoa2407417>.
 48. Long GV, Carlino MS, McNeil C, et al. Pembrolizumab versus ipilimumab for advanced melanoma: 10-year follow-up of the phase III KEY-NOTE-006 study. *Ann Oncol.* 2024;35(12):1191-1199.
 49. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Combined nivolumab and ipilimumab or monotherapy in untreated melanoma. *N Engl J Med.* 2015;373(1):23-34.
 50. Wolchok JD, Chiarion-Sileni V, Gonzalez R, et al. Long-term outcomes with nivolumab plus ipilimumab or nivolumab alone versus ipilimumab in patients with advanced melanoma. *J Clin Oncol.* 2022;40(2):127-137.
 51. Lebbé C, Meyer N, Mortier L, et al. Evaluation of two dosing regimens for nivolumab in combination with ipilimumab in patients with advanced melanoma: results from the phase IIb/IV CheckMate 511 trial. *J Clin Oncol.* 2019;37(11):867-875.
 52. Tawbi HA, Schadendorf D, Lipson EJ, et al. Relatlimab and nivolumab versus nivolumab in untreated advanced melanoma. *N Engl J Med.* 2022;386(1):24-34.
 53. Long GV, Hodi FS, Lipson EJ, et al. Overall survival and response with nivolumab and relatlimab in advanced melanoma. *NEJM Evid.* 2023;2(4):EVIDoA2200239.
 54. MERCK. Merck and Eisai provide update on phase 3 trials of KEYTRUDA® (pembrolizumab) plus LENVIMA® (lenvatinib) in certain patients with advanced melanoma (LEAP-003) and metastatic colorectal cancer (LEAP-017). Published April 7, 2023. Available at <https://www.merck.com/news/merck-and-eisai-provide-update-on-phase-3-trials-of-keytruda-pembrolizumab-plus-lenvima-lenvatinib-in-certain-patients-with-advanced-melanoma-leap-003-and-metastatic-colorectal-cancer/>. Accessed May 31, 2024.
 55. Long GV, Stroyakovskiy D, Gogas H, et al. Dabrafenib and trametinib versus dabrafenib and placebo for Val600 BRAF-mutant melanoma: a multicentre, double-blind, phase 3 randomised controlled trial. *Lancet.* 2015;386(9992):444-451.
 56. Robert C, Karaszewska B, Schachter J, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med.* 2015;372(1):30-39.
 57. Larkin J, Ascierto PA, Dreno B, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med.* 2014;371(20):1867-1876.
 58. Dummer R, Ascierto PA, Gogas HJ, et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol.* 2018;19(5):603-615.
 59. Schadendorf D, Dummer R, Flaherty KT, et al. COLUMBUS 7-year update: a randomized, open-label, phase III trial of encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF V600E/K-mutant melanoma. *Eur J Cancer.* 2024;204:114073.
 60. Atkins MB, Lee SJ, Chmielowski B, et al. Combination dabrafenib and trametinib versus combination nivolumab and ipilimumab for patients with advanced BRAF-mutant melanoma: the DREAMseq trial-ECOG-ACRIN EA6134. *J Clin Oncol.* 2023;41(2):186-197.
 61. Ascierto PA, Mandalà M, Ferrucci PF, et al. Sequencing of ipilimumab plus nivolumab and encorafenib plus binimetinib for untreated BRAF-mutated metastatic melanoma (SECOMBIT): a randomized, three-arm, open-label phase II trial. *J Clin Oncol.* 2023;41(2):212-221.
 62. Ascierto PA, Mandalà M, Ferrucci PF, et al. 1083MO - Brain metastases and survival evaluation in the SECOMBIT trial. *Ann Oncol.* 2023;34(suppl 2):S653.
 63. Robert C, Dutriaux C, Oppong FB, et al. Combination of encorafenib and binimetinib followed by ipilimumab and nivolumab versus ipilimumab and nivolumab in patients with advanced BRAF-V600E/K-mutated melanoma: the primary analysis of an EORTC randomized phase II study (EBIN). *J Clin Oncol.* 2024;42(suppl 17):LBA9503.
 64. Ascierto PA, Del Vecchio M, Robert C, et al. Ipilimumab 10 mg/kg versus ipilimumab 3 mg/kg in patients with unresectable or metastatic melanoma: a randomised, double-blind, multicentre, phase 3 trial. *Lancet Oncol.* 2017;18(5):611-622.
 65. Zimmer L, Apuri S, Eroglu Z, et al. Ipilimumab alone or in combination with nivolumab after progression on anti-PD-1 therapy in advanced melanoma. *Eur J Cancer.* 2017;75:47-55.
 66. VanderWalde A, Bellasea SL, Kendra KL, et al. Ipilimumab with or without nivolumab in PD-1 or PD-L1 blockade refractory metastatic melanoma: a randomized phase 2 trial. *Nat Med.* 2023;29(9):2278-2285.
 67. Ascierto PA, Lipson EJ, Dummer R, et al. Nivolumab and relatlimab in patients with advanced melanoma that had progressed on anti-programmed death-1/programmed death ligand 1 therapy: results from

- the phase I/IIa RELATIVITY-020 trial. *J Clin Oncol*. 2023;41(15):2724-2735.
68. Pham JP, Joshua AM, da Silva IP, et al. Chemotherapy in cutaneous melanoma: is there still a role? *Curr Oncol Rep*. 2023;25(6):609-621.
 69. Goldinger SM, Buder-Bakhaya K, Lo SN, et al. Chemotherapy after immune checkpoint inhibitor failure in metastatic melanoma: a retrospective multicentre analysis. *Eur J Cancer*. 2022;162:22-33.
 70. Sarnaik AA, Hamid O, Khushalani NI, et al. Lifileucel, a tumor-infiltrating lymphocyte therapy, in metastatic melanoma. *J Clin Oncol*. 2021;39(24):2656-2666.
 71. Rohaan MW, Borch TH, van den Berg JH, et al. Tumor-infiltrating lymphocyte therapy or ipilimumab in advanced melanoma. *N Engl J Med*. 2022;387(23):2113-2125.
 72. Monberg TJ, Borch TH, Svane IM, et al. TIL therapy: facts and hopes. *Clin Cancer Res*. 2023;29(17):3275-3283.
 73. Haanen J. *Overview of real-world experience in unmodified tumor-infiltrating lymphocytes therapy*. Chicago, Illinois, USA: Paper presented at the American Society of Clinical Oncology Annual Meeting; June 2, 2024.
 74. Arance A, de la Cruz-Merino L, Petrella TM, et al. Phase II LEAP-004 study of lenvatinib plus pembrolizumab for melanoma with confirmed progression on a programmed cell death protein-1 or programmed death ligand 1 inhibitor given as monotherapy or in combination. *J Clin Oncol*. 2023;41(1):75-85.
 75. Dummer R, Schadendorf D, Ascierto PA, et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol*. 2017;18(4):435-445.
 76. Valpione S, Carlino MS, Mangana J, et al. Rechallenge with BRAF-directed treatment in metastatic melanoma: a multi-institutional retrospective study. *Eur J Cancer*. 2018;91:116-124.
 77. Mateo J, Chakravarty D, Dienstmann R, et al. A framework to rank genomic alterations as targets for cancer precision medicine: the ESMO Scale for Clinical Actionability of molecular Targets (ESCAT). *Ann Oncol*. 2018;29(9):1895-1902.
 78. Cherny NI, Dafni U, Bogaerts J, et al. ESMO-Magnitude of Clinical Benefit Scale version 1.1. *Ann Oncol*. 2017;28(10):2340-2366.
 79. Dykewicz CA. Summary of the guidelines for preventing opportunistic infections among hematopoietic stem cell transplant recipients. *Clin Infect Dis*. 2001;33(2):139-144 (adapted from: Gross PA, Barrett TL, Dellinger EP, et al. Purpose of quality standards for infectious diseases. *Clin Infect Dis*. 1994;18(3):421).