



For numbered affiliations see end of article.

Correspondence to: J W Busse
bussejw@mcmaster.ca

Additional material is published online only. To view please visit the journal online

Cite this as: *BMJ* 2025;388:e079970
<http://doi.org/10.1136/bmj-2024-079970>

RAPID RECOMMENDATIONS

Commonly used interventional procedures for non-cancer chronic spine pain: a clinical practice guideline

Jason W Busse,^{1,2,3} Stéphane Genevay,⁴ Arnav Agarwal,² Christopher J Standaert,⁵ Kevin Carneiro,⁶ Jason Friedrich,⁷ Manuela Ferreira,⁸ Hilde Verbeke,⁹ Jens Ivar Brox,^{10,11} Hong Xiao,¹² Jasmeer Singh Virdee,¹³ Janet Gunderson,¹⁴ Gary Foster,¹⁵ Conrad Heegsma,¹⁵ Caroline F Samer,^{16,17} Matteo Coen,^{18,19} Gordon H Guyatt,² Xiaoqin Wang,² Behnam Sadeghirad,^{2,3} Faheem Malam,²⁰ Dena Zeraatkar,^{2,3} Per O Vandvik,²¹ Ting Zhou,²² Feng Xie,² Reed A C Siemieniuk,² Thomas Agoritsas^{2,18,23}

ABSTRACT

CLINICAL QUESTION

What is the comparative effectiveness and safety of commonly used interventional procedures (such as spinal injections and ablation procedures) for chronic axial and radicular spine pain that is not associated with cancer or inflammatory arthropathy?

CURRENT PRACTICE

Chronic spine pain is a common, potentially disabling complaint, for which clinicians often administer interventional procedures. However, clinical practice guidelines provide inconsistent recommendations for their use.

RECOMMENDATIONS

For people living with chronic axial spine pain (≥3 months), the guideline panel issued strong recommendations against: joint radiofrequency ablation with or without joint targeted injection of local anaesthetic plus steroid; epidural injection of local anaesthetic, steroids, or their combination; joint-targeted injection of local anaesthetic, steroids, or their combination; and intramuscular injection of local anaesthetic with or without steroids. For people living with chronic radicular spine pain (≥3 months), the guideline panel issued strong recommendations against: dorsal root ganglion radiofrequency with or without epidural injection of local anaesthetic or local anaesthetic plus steroids; and epidural injection of local anaesthetic, steroids, or their combination.

HOW THIS GUIDELINE WAS CREATED

An international guideline development panel including four people living with chronic spine pain, 10 clinicians with experience managing chronic spine pain, and eight methodologists, produced these recommendations in adherence with standards for trustworthy guidelines using the GRADE approach. The MAGIC Evidence Ecosystem Foundation provided methodological support. The guideline panel applied an individual patient perspective when formulating recommendations.

THE EVIDENCE

These recommendations are informed by a linked systematic review and network meta-analysis of randomised trials and a systematic review of observational studies, summarising the current body of evidence for benefits and harms of common interventional procedures for axial and radicular, chronic, non-cancer spine pain. Specifically, injection

of local anaesthetic, steroids, or their combination into the cervical or lumbar facet joint or sacroiliac joint; epidural injections of local anaesthetic, steroids, or their combination; radiofrequency of dorsal root ganglion; radiofrequency denervation of cervical or lumbar facet joints or the sacroiliac joint; and paravertebral intramuscular injections of local anaesthetic, steroids, or their combination.

UNDERSTANDING THE RECOMMENDATIONS

These recommendations apply to people living with chronic spine pain (≥3 months duration) that is not associated with cancer or inflammatory arthropathy and do not apply to the management of acute spine pain. Further research is warranted and may alter recommendations in the future: in particular, whether there are differences in treatment effects based on subtypes of chronic spine pain, establishing the effectiveness of interventional procedures currently supported by low or very low certainty evidence, and effects on poorly reported patient-important outcomes (such as opioid use, return to work, and sleep quality).

Introduction

Spine pain is defined as chronic when it persists for three months or longer and has resulted in pain on at least half of the days in the past six months.¹ Pain may be localised to the midline (axial) or referred distally (radicular) typically because of nerve root irritation (such as sciatica). Advanced imaging is often acquired for chronic spine pain, but incidental findings are common²⁻⁴ and there is low correlation between pathology and symptoms.⁵ Most chronic spine pain cannot be attributed to a specific cause, and approximately 85% of patients present with non-specific pain.⁶⁻⁹

The global prevalence of chronic low back pain has been estimated at 4% among adults aged 24-39 years and 20% among adults aged 20-59.¹⁰ The prevalence among older adults is likely higher,¹¹ and chronic low back pain is the leading cause of disability worldwide.¹² Neck pain is another common type of chronic spinal pain, estimated as the third leading cause of years lived with disability.¹³ In 2016, low back and neck pain accounted for the highest healthcare spending in the US at \$134.5 billion, of which 9.2% was patients' out-of-pocket expenses.¹⁴

Current practice

Interventional procedures—including paravertebral intramuscular injections, epidural injections, nerve blocks, and nerve ablation procedures—are increasingly used to manage chronic spine pain, particularly in North America. These procedures are hypothesised to attenuate chronic pain by interrupting pain related nerve signals through reducing local inflammation (epidural steroids), numbing nerves (nerve blocks), or targeted destruction of nerves responsible for transmitting pain (radiofrequency ablation).

Between 1994 and 2001 in the US, there was a 271% increase in lumbar epidural steroid injections (from 553 to 2055 per 100 000 patients) and a 231% increase in facet injections (from 80 to 264 per 100 000 patients) for low back pain.¹⁵ Facet joint or sacroiliac joint interventions in US Medicare recipients increased from approximately 425 000 in 2000 to 2.2 million interventions in 2013.¹⁶ From 2007 to 2016, data from a national US insurer showed a 131% increase in the use of lumbar radiofrequency procedures (from 49 to 113 per 100 000 patients).¹⁷ The number of US Medicare providers administering steroid injections along the spine increased 13% from 2012 to 2016.¹⁸

Despite rapid growth in use of interventional procedures for chronic spine pain, the supporting evidence is uncertain. An analysis of 17 review articles on epidural steroid injections for spine pain found inconsistent conclusions, and that positive results were three times more likely when the review was authored by an interventionalist (9 of 10 positive; 90%) versus a non-interventionalist (2 of 7 positive;


29%).¹⁹ The study authors suggested several explanations, including confirmation bias and secondary gain, as interventional procedures for chronic pain are often well reimbursed. For example, in 2017–18, the average billings among 106 physicians working at pain clinics providing nerve blocks in Ontario, Canada, was C\$724 183/year (£405 905/year).²⁰ Concurrently, a 2016/2017 survey of 777 Canadian physicians who performed interventional procedures for chronic spine pain found that only 37% believed their colleagues practiced in accordance with the current best evidence.²¹ Further, there is large variability among providers, with the top 10% of interventionalists in the US performing nine times more procedures per patient compared with the bottom 10%.²²


Why is the guideline needed?


A 2023 synthesis of 21 clinical practice guidelines on interventional procedures for low back pain concluded: “there was no consistency in recommendations for or against any interventional procedure, even after accounting for the quality of the [clinical practice guideline]”²³ (see table 1 for examples). One contributing factor is that several clinical practice guidelines for interventional procedures are characterised as consensus based.^{28–33} Such guidelines are more likely to produce recommendations that violate the principles of evidence based medicine than guidelines characterised as evidence based.³⁴ Moreover, guidelines for interventional procedures and chronic spine pain rarely involve patient partners, fail to consider patients’ values and preferences, and typically do not describe an explicit process to evaluate the overall certainty of evidence.³⁵

Table 1 | Examples of current guidance for interventional procedures and chronic spine pain



Recent guidelines	Recommendations
2022 American Society of Pain and Neuroscience (ASPN) Evidence-Based Clinical Guideline of Interventional Treatments for Low Back Pain ²⁴	<ul style="list-style-type: none">• Strong recommendation in favour of epidural injections (interlaminar, transforaminal, or caudal) of local anaesthetic, steroids, or their combination, for chronic low back pain due to disc disease, spinal stenosis, or post-surgical syndrome• Strong recommendation in favour of trigger point injections, irrespective of the type of medication, for chronic back pain• Strong recommendation in favour of sacroiliac joint injection for short term relief from sacroiliac joint dysfunction• Strong recommendation in favour of conventional or cooled lumbar radiofrequency ablation for low back pain• Strong recommendation in favour of sacroiliac joint denervation/ablation for sacroiliac joint dysfunction pain• Strong recommendation for basivertebral nerve ablation for chronic back pain
2021 American College of Occupational and Environmental Medicine Low Back Disorders Guideline ²⁵	<ul style="list-style-type: none">• Recommended against lumbar epidural injections for spinal stenosis (moderate confidence) or chronic low back pain in the absence of significant radicular symptoms (high confidence)• Trigger point injections of local anaesthetic may be recommended for chronic low back pain that is not resolving with exercise or NSAIDs (low confidence). Glucocorticosteroids are not recommended for use in trigger point injections (moderate confidence)• Therapeutic facet block injections are not recommended for chronic low back pain (low confidence) or any radicular pain syndrome (moderate confidence)• Radiofrequency neurotomy, neurotomy, or facet rhizotomy are not recommended for treatment of chronic low back pain, including that confirmed with diagnostic blocks (low confidence)• Dorsal root ganglia radiofrequency lesioning is not recommended for chronic sciatica (moderate certainty)
2021 Epidural interventions in the management of chronic spinal pain: American society of interventional pain physicians (ASIPP) comprehensive evidence-based guidelines ²⁶	<ul style="list-style-type: none">• Recommended in favour of fluoroscopically guided epidural injections, with or without steroids, for caudal epidural injections, lumbar interlaminar epidural injections, lumbar transforaminal epidural injections, and cervical interlaminar epidural injections for chronic spine pain associated with:<ul style="list-style-type: none">- Disc herniation (strong recommendation)- Spinal stenosis (moderate to strong recommendation)- Axial discogenic pain (moderate to strong recommendation)- Post-surgery syndrome (moderate to strong recommendation)
2020 NICE Guideline on Low back pain and sciatica in over 16s: assessment and management ²⁷	<ul style="list-style-type: none">• Do not offer spinal injections for managing low back pain• Only perform radiofrequency denervation for chronic low back pain after a positive response to a medial branch block

 Visual summary of recommendation



 **Population**
 This recommendation applies only to people with these characteristics:


 Adults with chronic spine pain (≥3 months duration)


Cervical spinal level


 Axial pain
  Radicular pain


Lumbar or sacroiliac spinal level

 Axial pain
  Radicular pain





 Does NOT apply to:

 Acute spine pain
 <3 months duration



 Chronic spine pain secondary to
 cancer inflammatory arthropathy


 **Strong recommendations AGAINST**
 All or nearly all well-informed people would likely not want such interventions. Such interventions should therefore not be offered outside of a clinical trial

Chronic axial spine pain

-  Epidural injection of local anaesthetic, steroids, or their combination (1)
-  Joint radiofrequency ablation with or without joint targeted injection of local anaesthetic and steroids (2)
-  Joint-targeted injection of local anaesthetic, steroids, or their combination (2)
-  Intramuscular injection of local anaesthetic with or without steroids (3)

Chronic radicular spine pain

-  Dorsal root ganglion radiofrequency with or without epidural injection of local anaesthetic, or local anaesthetic and steroids
-  Epidural injection of local anaesthetic, steroids, or their combination (1)

 **Additional areas of uncertainty**

Subgroup effects
 There may be differential effects of interventional procedures based on subtypes of chronic spine pain


Research needed
 Chronic axial spine pain:

- effectiveness of joint radiofrequency ablation
- intramuscular injection of local anaesthetic with or without steroids
- joint-targeted injection of local anaesthetic with or without steroids

Evidence required
 How interventions affect other patient-important outcomes that are poorly reported among existing trials:

- role functioning (including return to work)
- social functioning
- mental functioning
- sleep quality
- opioid use
- adverse events

A detailed summary of benefits and harms associated with interventional procedures is available in the linked systematic review and network meta-analysis

 <https://bit.ly/bmj-spine>

Practical issues

Cost and access

- Expense may be a barrier to accessing procedures unless patients have public or private coverage
- Patients must travel to a clinic or hospital that administers interventional procedures and, if perceived effective, return approximately every 2 weeks to 3 months for injections and approximately every 6 months for nerve ablation procedures

Adverse effects

- Interventional procedures may be associated with a small risk of moderate to serious harms, such as deep infection, and temporary altered level of consciousness
- Interventional procedures may be associated with a very small risk of catastrophic harms, such as paralysis and death following epidural steroid injection

Values and preferences

Chronic spine pain
 Most adults living with chronic spine pain place high value on small but important pain relief

Acceptable risks - interventional procedures
 Patients would be willing to accept the typical risks and burden associated with interventional procedures for an improvement in pain approximating the minimally important difference of 1.5 cm on a 10 cm visual analogue scale

Unacceptable risks
 Patients would be disinclined to receive treatment with an interventional procedure for which there is very low certainty of evidence for benefit or low certainty of evidence for no benefit, and moderate to high certainty evidence of risk of harm and/or burden

Find recommendations, evidence summaries and consultation decision aids for use in your practice

MAGIC app

© 2025 BMJ Publishing Group Ltd.

Disclaimer	Validation This infographic is not a validated clinical decision aid	Updating This information is provided without any representations, conditions, or warranties that it is accurate or up to date	Responsibility BMJ and its licensors assume no responsibility for any aspect of treatment administered with the aid of this information	Risks Any reliance placed on this information is strictly at the user's own risk
-------------------	--	--	---	--

For the full disclaimer wording see BMJ's terms and conditions: <http://www.bmj.com/company/legal-information/>

Given the lack of trustworthy guidelines in this area of high unmet clinical need, the Rapid Recommendations team identified that a careful appraisal of the full body of evidence would produce guidance that, if followed, would optimise the concordance between evidence and clinical use of interventional procedures for chronic spine pain.

The population considered for our guideline was adult patients living with chronic axial and/or radicular spine pain that was not associated with cancer, infection, or inflammatory spondylarthritis. Eligible procedures included joint-targeted injections (injection of local anaesthetic, steroids, or their combination into the cervical or lumbar facet joint, or sacroiliac joint); epidural injections of local

anaesthetic, steroids, or their combination; radiofrequency of dorsal root ganglion; radiofrequency denervation of cervical or lumbar facet joints, or the sacroiliac joint; and paravertebral intramuscular injections of local anaesthetic, steroids, or their combination. The panel's recommendations were informed by linked systematic reviews (box 1). The infographic provides the recommendations together with an overview of the absolute benefits and harms of common interventional procedures for chronic spine pain in the standard GRADE format.

Box 1: Linked articles in this *BMJ* Rapid Recommendations cluster

- Busse JW, Genevay S, Agarwal A, et al. Commonly used interventional procedures for non-cancer chronic spine pain: a clinical practice guideline. *BMJ* 2025;388:e079970, doi:10.1136/bmj-2024-079970
- Summary of results from the Rapid Recommendation process
- Wang X, Martin G, Sadeghirad B, et al. Common interventional procedures for chronic non-cancer spine pain: a systematic review and network meta-analysis of randomised trials. *BMJ* 2025;388:e079971, doi:10.1136/bmj-2024-079971
- Malam F, Asif MS, Khalid MF, et al. Adverse events associated with common interventional procedures for chronic spine pain: a systematic review and meta-analysis of non-randomized studies. *BMJ Open* (submitted)
- MAGICapp (<https://app.magicapp.org/#/guideline/nBRK8n>) multi-layered version of recommendations, rationale, and evidence summaries for use on all electronic devices

Patient and public involvement

Four people living with chronic spine pain, who were full members of the guideline panel, contributed to the selection and prioritisation of outcomes, values and preferences assessments, critical feedback to the protocol, and interpretation of findings for the *BMJ* Rapid Recommendation and the associated systematic reviews.

How the recommendations were created

Our international panel—including physiatrists (also called physical medicine and rehabilitation physicians), anaesthesiologists, rheumatologists, a physiotherapist, general internists, a clinical pharmacologist, epidemiologists, methodologists, and people living with chronic spine pain—decided the scope of recommendations and the outcomes that are most important to patients. Six of our clinical experts had experience administering interventional procedures for chronic spine pain. Our patient partners reported a range of experiences regarding interventional procedures. Three had received various interventional procedures for their chronic pain, with two reporting relief and one who did not. The fourth lived with chronic spine pain but had no personal experience with interventional procedures. After parallel teams completed a systematic review and network meta-analysis of randomised controlled trials on the benefits and harms of common interventional procedures for chronic spine pain, and a systematic review of observational studies exploring harms associated with interventional procedures, the panel met online four times to discuss the evidence and formulate recommendations. No panel member had financial conflicts of interest, and none declared any strong opinions for or against interventional procedures for chronic spine pain; intellectual and professional conflicts were minimised as per Rapid Recommendations usual methodology, by balancing them in the composition of the panel, and by having both a clinical and methods co-chair who were free of any conflicts of interest lead panel deliberations (appendix 1 on bmj.com).

The panel followed the *BMJ* Rapid Recommendations approach for creating trustworthy guidance,⁶⁷ including use of GRADE to critically appraise the evidence and create recommendations.⁶⁸ The panel considered the balance of benefits, harms, and burdens of each intervention, the

certainty of the evidence for each outcome, typical and expected variations in patient values and preferences, and practical issues related to use, acceptability, feasibility, and equity.⁶⁹

Recommendations can be strong or conditional, and for or against a course of action. Strong recommendations typically require a clear imbalance between benefits and harms or burdens supported by high or moderate certainty evidence; however, there are five paradigmatic scenarios in which a strong recommendation can be made based on low certainty evidence.⁶⁹ One such scenario is when there is low certainty of benefit (or lack of benefit) and moderate to high certainty of greater risk of important harm or burden.

We required 80% consensus among panel members for strong recommendations and a majority consensus for conditional recommendations. Two experienced guideline methodologists (JWB and TA) oversaw the consensus process. The evidence synthesis teams prepared draft summary of findings tables before the panel meetings, following GRADE guidance, from the accompanying network meta-analysis³⁶ and systematic review of observational studies.⁴⁴

The evidence

The linked systematic review and network-meta-analysis included 132 randomised trials, of which 81 trials with 7977 participants were included in meta-analyses.³⁶ These trials reported on 13 categories of interventional procedures compared with usual care or sham procedures in patients with axial or radicular chronic spine pain (box 2). Table 2 and supplementary table 1 (appendix 2) on bmj.com provide an overview of the numbers and types of patients included, study funding, subtypes of chronic spine pain, and whether a positive diagnostic block was an eligibility criterion. Our panel selected eight patient-important outcomes: (1) pain relief, (2) physical functioning, (3) emotional functioning, (4) role functioning, including return to work, (5) social functioning, (6) sleep quality, (7) opioid use, and (8) adverse events. Our selection process was guided by the Initiative on Methods, Measurement, and Pain Assessment in Clinical Trials (IMMPACT).^{39 40} The panel designated pain relief as our critical outcome. This is supported by a discrete choice experiment involving 211 adults living with chronic low back pain, which found that patients are most concerned with pain relief, followed by duration of pain relief.⁴¹

Box 2: Categories of interventional procedures administered for chronic spine pain that were considered in the *BMJ* Rapid Recommendations review

- Epidural injection of local anaesthetic
- Epidural steroid injection
- Epidural injection of local anaesthetic and steroids
- Joint-targeted injection of local anaesthetic
- Joint-targeted steroid injection
- Joint-targeted injection of local anaesthetic and steroids
- Intramuscular injection of local anaesthetic
- Intramuscular injection of local anaesthetic and steroids
- Dorsal root ganglion radiofrequency
- Dorsal root ganglion radiofrequency with epidural injection of local anaesthetic
- Dorsal root ganglion radiofrequency with epidural injection of local anaesthetic with steroids
- Joint radiofrequency nerve ablation
- Joint radiofrequency nerve ablation with joint-targeted injection of local anaesthetic and steroids

Table 2 | Characteristics from 81 eligible randomised clinical trials (7977 patients) included in a linked meta-analysis

Characteristic	Median (interquartile range) of means across trials unless stated otherwise
No of patients enrolled	64 (45-110)*
Age (mean years at baseline)	48 (44-56)
Sex (% women)	59% (50-67)†
Baseline pain (on 10 cm visual analogue scale for pain)	6.8 (5.9-7.7)‡
Presenting complaint§	<ul style="list-style-type: none"> Chronic axial spine pain (n=45): facet joint syndrome (n=28; 35%), sacroiliac joint syndrome (n=9; 11%), intervertebral disc syndrome (n=3; 4%), not reported (n=2; 2%), spondylolisthesis (n=1; 1%), chronic post-surgical pain (n=1; 1%), mixed complaints (n=1; 1%) Chronic radicular spine pain (n=34): disc herniation/bulge (n=24; 30%), spinal stenosis (n=8; 10%), chronic post-surgical pain (n=1; 1%), mixed complaints (n=1; 1%) Mixed chronic axial and radicular spine pain (n=2; 3%)
Pain duration	<ul style="list-style-type: none"> 62% of trials (n=50) reported duration of chronic pain: median duration 46 months (IQR 17-98) 25% of trials (n=20) reported that pain was "chronic" or enrolled patients with chronic conditions, without other details 13% of trials (n=11) enrolled ≥80% of patients with pain duration of ≥3 months
Positive diagnostic block required for enrolment¶	<ul style="list-style-type: none"> Yes (n=32): >50% relief (n=13; 16%), >75% relief (n=5; 6%), >80% relief (n=6; 7%), unclear threshold for relief (n=6; 7%), used a positive block as an exclusion criterion (n=2; 3%) No or not reported (n=49; 60%)
Trial location	North America (n=26; 32%), Europe (n=25; 31%), Asia (n=23; 28%), Africa (n=4; 5%), South America (n=2; 2%), Australia (n=1; 1%)
Funding	Not reported (n=33; 41%), unfunded (n=25; 31%), non-industry funding (n=22; 27%), industry funded (n=1; 1%)

Additional details are available in the network meta-analysis.³⁶

* Total number of enrolled patients using study-level data.

† Proportion among eligible trials reporting this information (n=75).

‡ 81 trials reported baseline pain of patients.

§ As reported by trial authors.

¶ Diagnostic blocks (an injection of local anaesthetic to confirm the correct targeting of the nerve) carry a false positive rate of up to 60%.^{37,38}

Values and preferences

We surveyed our guideline panel, including the four patient partners, using an established framework^{42,43} to inform the perceived values and preferences of a typical person living with chronic spine pain on potential benefits as well as harms and burdens of interventional procedures. We conducted our survey before presenting the panel with the results of our evidence syntheses informing benefits and harms of interventional procedures for chronic spine pain.^{36,44} This exercise was informed by prior systematic reviews that found people living with chronic pain typically place high value on the possibility of small but important pain relief.^{45,46} Our panel survey revealed that most people living with chronic spine pain would be inclined to receive an interventional procedure, even with moderate to high certainty evidence of harms or burden, if there was moderate or high certainty evidence of important benefit.

Alternately, our panel advised that, when presented with an interventional procedure of uncertain effectiveness (very low certainty evidence), and for which there is moderate to high certainty evidence of increased risk of harm or burden, almost all people living with chronic spine pain would be disinclined to receive treatment. Further, the panel inferred that almost all people living with chronic spine pain, when presented with an interventional procedure for which the point estimate suggests no benefit, but the evidence is low certainty, and where there is moderate to high evidence of increased risk of harm or burden, would be disinclined to receive treatment. Finally, in the presence of low certainty evidence of benefit for an interventional procedure, and moderate to high evidence of harm or burden, a minority (<50%) of people

living with chronic spine pain would be interested in receiving treatment.

Following our panel meeting to establish patients' values and preferences, a discrete choice experiment was published that involved 424 individuals with chronic low back pain.⁴⁷ The investigators assessed participants' preferences towards non-surgical treatment options for chronic back pain, including mind-body interventions, supervised physical activation, physical manipulations, self management courses, physiotherapy, and corticosteroid injections. They found corticosteroid injections were rejected by the large majority and that participants were most likely to have concerns about receiving corticosteroid injections versus other non-surgical options for their chronic low back pain.

Understanding the recommendations

The panel reviewed the evidence for benefits and harms among the 13 selected interventional procedures or combinations of procedures for chronic spine pain (box 2).^{36,44} There was no high certainty evidence of important pain relief (or benefit on any other effectiveness outcome) for any intervention for either chronic axial spine pain or chronic radicular spine pain. All interventional procedures supported by moderate or low certainty evidence showed little to no pain relief compared with sham procedures (see infographic).

We found no evidence of subgroup effects based on clinical condition.³⁶ Further, since all pooled effects in our network meta-analysis supported by low or moderate certainty evidence showed little to no effect on pain relief relative to sham procedures, then if an interventional procedure was effective in certain subtypes

of axial or radicular pain it must increase pain in other subtypes; we judged this was unlikely.

Low certainty evidence supported a 0.7% incidence of deep infection (after joint radiofrequency nerve ablation, joint targeted steroid injection, and epidural injection of local anaesthetic and steroids), a 1.4% incidence of dural puncture (after epidural steroid injection, joint radiofrequency nerve ablation, and joint-targeted injection of local anaesthetic and steroids), an 8.6% incidence of prolonged (>48 hours) pain or stiffness (after joint radiofrequency nerve ablation with or without joint targeted injection of steroids, and dorsal root ganglion radiofrequency) and a 2.1% incidence of temporary altered level of consciousness (after joint targeted steroid injection, and epidural steroid injection).⁴⁴ The panel was also aware of very rare but catastrophic complications of interventional procedures for spine pain not captured in our evidence syntheses, such as paraplegia after epidural injection.⁴⁸⁻⁵⁰ The panel had high certainty that undergoing interventional procedures for chronic spine pain was associated with important burden (such as travel, discomfort, productivity loss), which would be recurring as these interventions are typically repeated on a regular basis, and that some patients would bear substantial out-of-pocket costs.

The panel concluded that all or almost all informed patients would choose to avoid interventional procedures for axial or radicular chronic spine pain because all low and moderate certainty evidence suggests little to no benefit on pain relief compared with sham procedures, and these procedures are burdensome and may result in adverse events.⁵¹⁻⁵² The panel acknowledged that the evidence for some interventional procedures was only low or very low certainty and agreed it would be appropriate to provide them in a research setting.

To whom do the recommendations apply?

The recommendations apply to adults living with moderate to severe chronic, axial or radicular, spine pain (that is, neck, back, sacroiliac) lasting three months or longer in duration. They do not apply to the management of acute spine pain (<3 months duration), or chronic spine pain associated with cancer or inflammatory arthropathy.

Absolute benefits and harms

The infographic explains the recommendations and provides links to MAGICapp with evidence summaries of absolute benefits and harms of interventional procedures for chronic spine pain. Estimates of baseline risk for effects come from the control arms of eligible trials from the associated network meta-analysis.³⁶ Only approximately half of randomised trials eligible for our network meta-analysis reported adverse events, and this evidence, as well as the results from our systematic review of observational studies on harms from interventional procedures,⁴⁴ proved only low or very low certainty.

The clinical experts on our panel considered findings from our evidence syntheses regarding the potential harms associated with interventional procedures, as well as published reports on very rare but severe harms. The resulting consensus was that interventional procedures for chronic spine pain were costly and may be associated with a small risk of moderate harms (for example, an 8.6% risk of prolonged (>48 hours) pain or stiffness, 2.1% risk of temporary altered level of consciousness, 1.4% risk of dural puncture, 0.7% risk of deep infection),⁴⁴⁻⁵³ and a very small risk of catastrophic harms (such as infection resulting in meningitis, spinal cord injury, and paraplegia).⁵⁴⁻⁵⁸ We were unable to quantify the risks of catastrophic harms as they were reported in case studies or databases that did not specify a denominator. For example, between

1997 and 2014, a total of 90 serious adverse events that occurred within minutes to 48 hours after epidural injections of corticosteroids for management of neck and back pain were captured by the US Food and Drug Administration (FDA) Adverse Event Reporting System database. These included death, spinal cord infarction, paraplegia, quadriplegia, cortical blindness, stroke, seizures, and brain oedema.⁵⁹⁻⁶⁰

- The panel was confident of the following, relative to sham procedures:
 - Moderate certainty evidence showed that, for chronic axial spine pain, epidural injection of local anaesthetic (with or without steroids) and joint-targeted steroid injections probably have little to no effect on pain relief.³⁶
 - Moderate certainty evidence showed that, for chronic radicular spine pain, epidural injection of local anaesthetic with steroids and dorsal root ganglion radiofrequency probably have little to no effect on pain relief.³⁶
 - It is unlikely that new information will result in important changes in best estimates of effect for outcomes that are supported by moderate certainty evidence.
- The panel was less confident about:
 - For chronic axial spine pain, the effect of intramuscular injection of local anaesthetic (with or without steroids), epidural injection of steroids, and joint-targeted injection of local anaesthetic (with or without steroids) on pain relief. Although effects showed little to no difference in pain relief (except for intramuscular injection of local anaesthetic and steroids, which showed increased pain) versus sham procedures, the evidence was only low certainty. We considered that a beneficial effect of epidural injection of steroids is unlikely because there is moderate certainty evidence that an epidural injection with steroids and local anaesthetic probably has little to no effect on pain. The effect of joint radiofrequency was supported by only very low certainty evidence due to small study effects and risk of bias. Four trials with unblinded providers reported larger effects on pain relief than did seven trials with blinded providers (-1.74 cm on a 10 cm visual analogue scale for pain relief (95% confidence interval -2.73 to -0.76) for unblinded trials versus -0.23 cm (-0.60 to 0.14) for blinded trials; test of interaction P value 0.005).³⁶
 - For chronic radicular spine pain, epidural injection of local anaesthetic or steroids showed little to no difference in pain relief, but the evidence was only low certainty. However, the effect of epidural injection of either local anaesthetic or steroids in isolation is unlikely as we found moderate evidence that the combination is probably not effective.³⁶
 - Harms associated with interventional procedures for chronic spine pain, which were supported by very low to low certainty evidence.³⁶⁻⁴⁴

Practical issues and other considerations

Box 3 outlines the key practical issues for patients and clinicians regarding interventional procedures for chronic spine pain. Interventional procedures are associated with burden to patients, who must travel to a healthcare provider. They are not curative and, if they have any effect at all, intramuscular, joint-targeted, or epidural injections are typically repeated approximately every 2

weeks to 3 months. Nerve ablation procedures, if they have any effect at all, are typically repeated approximately every 6 months.

Box 3: Practical issues

Cost and access

- Expense may be a barrier to accessing interventional procedures unless patients reside in a country in which the government covers the cost, or they have private coverage
- Patients must travel to a clinic or hospital that administers the interventional procedure and, if the procedure is perceived to be effective, return approximately every 2 weeks to 3 months for injections or approximately every 6 months for nerve ablation procedures

Adverse effects

- Interventional procedures may be associated with a small risk of moderate harms, such as temporary altered level of consciousness, deep infection, and prolonged pain and stiffness
- Interventional procedures may be associated with a very small risk of catastrophic harms, including paralysis and death

In some jurisdictions, patients will bear the costs of interventional procedures, which may be substantial. In the US, the average cost for a single epidural steroid injection is more than US\$1000, and can be as high as US\$5000, and the average cost for radiofrequency ablation is approximately US\$6,000.^{18 61–63} Considering a middle-income country, such as China, the cost of interventional procedures for chronic spine pain ranges from US\$3 to US\$538. We have provided a detailed breakdown of costs in China for interventional spine procedures in supplementary tables 2 and 3 (appendix 2). Despite our finding that current evidence suggests common interventional procedures are no more effective than sham procedures for chronic spine pain, the substantial reimbursement associated with these procedures may act as a perverse incentive for their delivery as opposed to less well paying, and more time-consuming, interventions that have evidence of effectiveness (for example, cognitive functional therapy,⁶⁴ exercise therapy,⁶⁵ pain reprocessing therapy⁶⁶).

Costs and resources

When formulating recommendations, the panel focused on patients' perspectives rather than that of society. However, both availability and costs of interventional procedures for chronic spine pain are likely to influence decision making.

Uncertainties for future research

- Key research questions to inform decision makers and future guidelines include:
 - Are there systematic differences in treatment effects of interventional procedures based on subtypes of chronic spine pain? Our network meta-analysis found no credible subgroup effects, but we were unable to explore all clinical conditions due to limited evidence for some types of pain.³⁶
 - What are the effects on pain relief for interventional procedures currently informed by low or very low certainty evidence? Specifically, joint radiofrequency, intramuscular injection of local anaesthetic (with or without steroids), and joint-targeted injection of local anaesthetic (with or without steroids) for chronic axial pain.
 - What are the effects of interventional procedures for chronic spine pain on patient-important outcomes that were poorly reported among trials that informed our evidence synthesis?

Specifically, role functioning (including return to work), social functioning, mental functioning, sleep quality, opioid use, and adverse events.

How patients were involved in the creation of this article:

Four people living with chronic spine pain, including two military veterans, were full panel members. These panel members identified important outcomes, informed the discussion on values and preferences and voted on all recommendations. They participated in online meetings and email discussions and met all authorship criteria.

Education in practice

- How do you currently approach the management of people living with chronic spine pain that is not associated with cancer or inflammatory arthropathy?
- How can this article help you explain the evidence to patients considering common interventional procedures for their chronic spine pain?

AUTHOR AFFILIATIONS

- 1 Michael G. DeGroot National Pain Centre, McMaster University, Hamilton, ON, Canada
- 2 Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada
- 3 Department of Anesthesia, McMaster University, Hamilton, ON, Canada
- 4 Division of Rheumatology, Department of Medicine, Geneva University Hospitals, Geneva, Switzerland
- 5 Department of Physical Medicine and Rehabilitation, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA
- 6 Department of Neurosurgery, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
- 7 Department of Physical Medicine and Rehabilitation, University of Colorado School of Medicine, Aurora, CO, USA
- 8 The University of Sydney, Sydney Musculoskeletal Health and the Kolling Institute, School of Health Sciences, Faculty of Medicine and Health, St Leonards, NSW 2064, Australia
- 9 Leuven Center for Algology and Pain Management, University Hospitals Leuven, Leuven, Belgium
- 10 Department of Physical Medicine and Rehabilitation, Oslo University Hospital Ullevaal, Oslo, Norway
- 11 Institute of Clinical Medicine, Medical Faculty, Oslo University, Oslo, Norway
- 12 Department of Pain Management, West China Hospital, Sichuan University, Chengdu, Sichuan, China
- 13 International Trade Centre, Palais des Nations, Geneva, Switzerland
- 14 Chronic Pain Network, McMaster University, Hamilton, Ontario, Canada
- 15 The Canadian Veterans Chronic Pain Centre of Excellence, Hamilton, ON, Canada
- 16 Division of Clinical Pharmacology and Toxicology, Geneva University Hospitals
- 17 Faculty of Medicine, University of Geneva, Switzerland
- 18 Division of General Internal Medicine, Department of Medicine, University Hospitals of Geneva, University of Geneva, Switzerland

- ¹⁹ Unit of Development and Research in Medical Education, Faculty of Medicine, University of Geneva, Geneva, Switzerland
- ²⁰ Michael G. DeGroote School of Medicine, McMaster University, Hamilton, ON, Canada
- ²¹ Department of Medicine, Innlandet Hospital Trust, Gjøvik, Norway
- ²² School of International Pharmaceutical Business, China Pharmaceutical University, Nanjing, China
- ²³ The MAGIC Evidence Ecosystem Foundation, Oslo, Norway

Competing interests: The BMJ has judged that there are no disqualifying financial ties to commercial companies. The authors declare the following other interests: none. Further details of The BMJ policy on financial interests is here: <https://www.bmj.com/sites/default/files/attachments/re-sources/2016/03/16-current-bmj-education-coi-form.pdf>.

Funding: This guideline was funded by the Chronic Pain Centre of Excellence for Canadian Veterans. The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication. JWB is supported, in part, by a Canadian Institutes of Health Research Canada Research Chair in the prevention and management of chronic pain.

Transparency: JWB affirms that the manuscript is an honest, accurate, and transparent account of the recommendations being reported; that no important aspects of the recommendations have been omitted; and that any discrepancies from the recommendations as planned (and, if relevant, registered) have been explained.

This *BMJ Rapid Recommendation* article is one of a series that provides clinicians with trustworthy recommendations for potentially practice changing evidence. *BMJ Rapid Recommendations* represent a collaborative effort between the MAGIC group (www.magicvidence.org) and *The BMJ*. A summary is offered here, and the full version is on the MAGICapp (www.magicapp.org) for all devices in multilayered formats. We encourage adaptation and contextualisation of our recommendations to local or other contexts. Those considering use or adaptation of content may go to MAGICapp to link or extract its content, or contact *The BMJ* for permission to reuse content in this article.

We thank: Rachel J Couban, medical librarian, for assistance in identifying guidelines for management of chronic spine pain; Will Stahl-Timmins and colleagues at *The BMJ* for co-creation of the infographic.

Our 22 member guideline panel included four patient partners (Jasmeer Singh Virdee, Janet Gunderson, Gary Foster, and Conrad Heegsma), eight methodologists, five of whom have also general medicine expertise (Jason W Busse, Arnav Agarwal, Thomas Agoritsas, Gordon H Guyatt, Xiaoqin Wang, Behnam Sadeghirad, Per O Vandvik, and Reed A C Siemieniuk), and 10 clinical experts (Stéphane Genevay [rheumatologist], Christopher J Standaert [physiatrist], Kevin Carneiro [physiatrist], Jason Friedrich [physiatrist], Manuela Ferreira [physiotherapist], Hilde Verbeke [anaesthesiologist], Jens Ivar Brox [physiatrist], Hong Xiao [anaesthesiologist], Matteo Coen [general internist], and Caroline Samer [clinical pharmacologist]).

- 1 Deyo RA, Dworkin SF, Amtmann D, et al. Report of the NIH task force on research standards for chronic low back pain. *Spine (Phila Pa 1976)* 2014;39:43. . doi: 10.1097/BRS.0000000000000434 PMID: 24887571
- 2 Kuijper B, Tans JT, van der Kallen BF, Nollef F, Lycklama A Nijeholt GJ, de Visser M. Root compression on MRI compared with clinical findings in patients with recent onset cervical radiculopathy. *J Neurol Neurosurg Psychiatry* 2011;82:3. . doi: 10.1136/jnnp.2010.217182 PMID: 21047885
- 3 Jensen MC, Brant-Zawadzki MN, Obuchowski N, Modic MT, Malkasian D, Ross JS. Magnetic resonance imaging of the lumbar spine in people without back pain. *N Engl J Med* 1994;331:73. . doi: 10.1056/NEJM199407143310201 PMID: 8208267
- 4 Brinjikji W, Luetner PH, Comstock B, et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol* 2015;36:6. . doi: 10.3174/ajnr.A4173 PMID: 25430861
- 5 Mounce K. Back pain. *Rheumatology (Oxford)* 2002;41:5. . doi: 10.1093/rheumatology/41.1.1 PMID: 11792871
- 6 Chou R, Qaseem A, Owens DK, Shekelle PC. Clinical Guidelines Committee of the American College of Physicians. Diagnostic imaging for low back pain: advice for high-value health care from the American College of Physicians. *Ann Intern Med* 2011;154:9. . doi: 10.7326/0003-4819-154-3-201102010-00008 PMID: 21282698
- 7 Liu R, Kurihara C, Tsai HT, et al. Classification and treatment of chronic neck pain: a longitudinal cohort study. *Reg Anesth Pain Med* 2017;42:61. . doi: 10.1097/AAP.0000000000000505 PMID: 27846187
- 8 Knezevic NN, Candido KD, Vlaeyen JWS, Van Zundert J, Cohen SP. Low back pain. *Lancet* 2021;398:92. . doi: 10.1016/S0140-6736(21)00733-9 PMID: 34115979
- 9 Schubiner H, Lowry WJ, Heule M, et al. Application of a clinical approach to diagnosing primary pain: prevalence and correlates of primary back and neck pain in a community physiatry clinic. *J Pain* 2024;25:81. . doi: 10.1016/j.jpain.2023.09.019 PMID: 37777033
- 10 Meucci RD, Fassa AG, Faria NM. Prevalence of chronic low back pain: systematic review. *Rev Saude Publica* 2015;49:.. doi: 10.1590/S0034-8910.20150490005874 PMID: 26487293
- 11 Wong CK, Mak RY, Kwok TS, et al. Prevalence, incidence, and factors associated with non-specific chronic low back pain in community-dwelling older adults aged 60 years and older: a systematic review and meta-analysis. *J Pain* 2022;23:34. . doi: 10.1016/j.jpain.2021.07.012 PMID: 34450274
- 12 GBD 2021 Low Back Pain Collaborators. Global, regional, and national burden of low back pain, 1990-2020, its attributable risk factors, and projections to 2050: a systematic analysis of the Global Burden of Disease Study 2021. *Lancet Rheumatol* 2023;5:29. . doi: 10.1016/S2665-9913(23)00098-X PMID: 37273833
- 13 James SL, Abate D, Abate KH, et al. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;392:858. . doi: 10.1016/S0140-6736(18)32279-7 PMID: 30496104
- 14 Dieleman JL, Cao J, Chapin A, et al. US health care spending by payer and health condition, 1996-2016. *JAMA* 2020;323:84. doi: 10.1001/jama.2020.0734 PMID: 32125402
- 15 Friedly J, Chan L, Deyo R. Increases in lumbosacral injections in the Medicare population: 1994 to 2001. *Spine (Phila Pa 1976)* 2007;32:60. . doi: 10.1097/BRS.0b013e3180b9f96e PMID: 17632396
- 16 Manchikanti L, Hirsch JA, Pampati V, Boswell MV. Utilization of facet joint and sacroiliac joint interventions in Medicare population from 2000 to 2014: Explosive growth continues!. *Curr Pain Headache Rep* 2016;20:.. doi: 10.1007/s11916-016-0588-2 PMID: 27646014
- 17 Starr JB, Gold L, McCormick Z, Suri P, Friedly J. Trends in lumbar radiofrequency ablation utilization from 2007 to 2016. *Spine J* 2019;19:28. . doi: 10.1016/j.spinee.2019.01.001 PMID: 30639589
- 18 Kaplan S. After doctors cut their opioids, patients turn to a risky treatment for back pain. *New York Times* 2018. <https://www.nytimes.com/2018/07/31/health/opioids-spinal-injections.html>.
- 19 Cohen SP, Bicket MC, Jamison D, Wilkinson I, Rathmell JP. Epidural steroids: a comprehensive, evidence-based review. *Reg Anesth Pain Med* 2013;38:200. . doi: 10.1097/AAP.0b013e31828ea086 PMID: 23598728
- 20 Oved MC, McLean J. Chronic pain management is big business in Ontario – and these four private clinic chains dominate the industry. *Toronto Star* 2020 Sept 30. https://www.thestar.com/news/investigations/chronic-pain-management-is-big-business-in-ontario-and-these-four-private-clinic-chains-dominate/article_9e6bb927-c682-5df1-b1d2-86bf37980d60.html.
- 21 Shanthanna H, Bhatia A, Radhakrishna M, et al. Interventional pain management for chronic pain: a survey of physicians in Canada. *Can J Anaesth* 2020;67:52. . doi: 10.1007/s12630-019-01547-w PMID: 31802414
- 22 Abbott ZI, Nair KV, Allen RR, Akuthota VR. Utilization characteristics of spinal interventions. *Spine J* 2012;12:43. . doi: 10.1016/j.spinee.2011.10.005 PMID: 22138113
- 23 Olivier TJ, Konda C, Pham T, et al. Clinical practice guidelines on interventional management of low back pain: A synthesis of recommendations. *PM R* 2023;15:63. . doi: 10.1002/pmrj.12930 PMID: 36507598
- 24 Sayed D, Grider J, Strand N, et al. The American Society of Pain and Neuroscience (ASPN) evidence-based clinical guideline of interventional treatments for low back pain. *J Pain Res* 2022;15:832. . doi: 10.2147/JPR.S386879 PMID: 36510616
- 25 Hegmann KT, Travis R, Andersson GBJ, et al. Invasive treatments for low back disorders. *J Occup Environ Med* 2021;63:41. . doi: 10.1097/JOM.0000000000001983 PMID: 33769405
- 26 Manchikanti L, Knezevic NN, Navani A, et al. Epidural interventions in the management of chronic spinal pain: American Society of Interventional Pain Physicians (ASIPP) comprehensive evidence-based guidelines. *Pain Physician* 2021;24(S1):208.PMID: 33492918
- 27 National Institute for Health and Care Excellence. Low back pain and sciatica in over 16s: assessment and management (NICE guideline NG59). 2020. <https://www.nice.org.uk/guidance/ng59>.
- 28 Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. *Reg Anesth Pain Med* 2020;45:67. . doi: 10.1136/rapm-2019-101243 PMID: 32245841
- 29 Hurley RW, Adams MCB, Barad M, et al. Consensus practice guidelines on interventions for cervical spine (facet) joint pain from a multispecialty international working group. *Reg Anesth Pain Med* 2022;47:59. . doi: 10.1136/rapm-2021-103031 PMID: 34764220
- 30 Bailly F, Trouvin AP, Bercier S, et al. Clinical guidelines and care pathway for management of low back pain with or without radicular pain. *Joint Bone Spine* 2021;88:105227. . doi: 10.1016/j.jbspin.2021.105227 PMID: 34051387
- 31 Occhigrossi F, Carpenedo R, Leoni MLG, Varrassi G, Chinè E, Cascella MCompain Research Group. Delphi-based expert consensus statements for the management of percutaneous radiofrequency neurotomy in the treatment of lumbar facet joint syndrome. *Pain Ther* 2023;12:77. . doi: 10.1007/s40122-023-00512-2 PMID: 37103732
- 32 Wang Y, Wang AZ, Wu BS, et al. Chinese Association for the Study of Pain: Experts consensus on ultrasound-guided injections for the treatment of spinal pain in China (2020 edition). *World J Clin Cases* 2021;9:57. . doi: 10.12998/wjcc.v9.i9.2047 PMID: 33850924
- 33 Liu JF, Shen W, Huang D, et al. Expert consensus of Chinese Association for the Study of Pain on the radiofrequency therapy technology in the Department of Pain. *World J Clin Cases* 2021;9:35. . doi: 10.12998/wjcc.v9.i9.2123 PMID: 33850931
- 34 Yao L, Ahmed MM, Guyatt GH, et al. Discordant and inappropriate discordant recommendations in consensus and evidence based guidelines: empirical analysis. *BMJ* 2021;375:e066045. . doi: 10.1136/bmj-2021-066045 PMID: 34824101
- 35 Hoydonckx Y, Kumar P, Flamer D, et al. Quality of chronic pain interventional treatment guidelines from pain societies: Assessment with the AGREE II instrument. *Eur J Pain* 2020;24:21. . doi: 10.1002/ejp.1524 PMID: 31880843

- 36 Wang X, Martin G, Sadeghirad B, et al. Interventional procedures for chronic noncancer spine pain: a systematic review and network meta-analysis of randomised trials. *BMJ* 2025;388:e079971. doi: 10.1136/bmj-2024-079971.
- 37 Manchikanti L, Pampati V, Fellows B, Bakht CE. Prevalence of lumbar facet joint pain in chronic low back pain. *Pain Physician* 1999;2:64. doi: 10.36076/ppj.1999/2/59 pmid: 16906217
- 38 Falco FJ, Erhart S, Wargo BW, et al. Systematic review of diagnostic utility and therapeutic effectiveness of cervical facet joint interventions. *Pain Physician* 2009;12:44. doi: 10.36076/ppj.2009/12/323 pmid: 19305483
- 39 Turk DC, Dworkin RH, Allen RR, et al. Core outcome domains for chronic pain clinical trials: IMMPACT recommendations. *Pain* 2003;106:45. . doi: 10.1016/j.pain.2003.08.001 pmid: 14659516
- 40 Turk DC, Dworkin RH, Revicki D, et al. Identifying important outcome domains for chronic pain clinical trials: an IMMPACT survey of people with pain. *Pain* 2008;137:85. . doi: 10.1016/j.pain.2007.09.002 pmid: 17937976
- 41 Wilson L, Zheng P, Ionova Y, et al. CAPER: patient preferences to inform nonsurgical treatment of chronic low back pain: a discrete-choice experiment. *Pain Med* 2023;24:73. . doi: 10.1093/pm/pnad038 pmid: 36975607
- 42 Zeng L, Helsing LM, Bretthauer M, et al. A novel framework for incorporating patient values and preferences in making guideline recommendations: guideline panel surveys. *J Clin Epidemiol* 2023;161:72. . doi: 10.1016/j.jclinepi.2023.07.003 pmid: 37453455
- 43 Zeng L, Li SA, Yang M, et al. Qualitative study of guideline panelists: innovative surveys provided valuable insights regarding patient values and preferences. *J Clin Epidemiol* 2023;161:80. . doi: 10.1016/j.jclinepi.2023.07.014 pmid: 37517505
- 44 Malam F, Asif MS, Khalid MF, et al. Adverse events associated with interventional procedures for chronic spine pain: a systematic review and meta-analysis of non-randomized studies. *BMJ Open* (submitted).
- 45 Zeng L, Lytyyn L, Wang X, et al. Values and preferences towards medical cannabis among people living with chronic pain: a mixed-methods systematic review. *BMJ Open* 2021;11:e050831. . doi: 10.1136/bmjopen-2021-050831 pmid: 34493521
- 46 Goshua A, Craigie S, Guyatt GH, et al. Patient values and preferences regarding opioids for chronic noncancer pain: a systematic review. *Pain Med* 2018;19:80. . doi: 10.1093/pm/pnx274 pmid: 29618109
- 47 Morillon GF, Benkhalti M, Dagenais P, Poder TG. Preferences of patients with chronic low back pain about nonsurgical treatments: Results of a discrete choice experiment. *Health Expect* 2023;26:30. . doi: 10.1111/hex.13685 pmid: 36482802
- 48 Petrin Z, Marino RJ, Oleson CV, Simon JI, McCormick ZL. Paralysis after lumbar interlaminar epidural steroid injection in the absence of hematoma: A case of congestive myelopathy due to spinal dural arteriovenous fistula and a review of the literature. *Am J Phys Med Rehabil* 2020;99:10. . doi: 10.1097/PHM.0000000000001325 pmid: 31592878
- 49 Jeon SH, Jang W, Kim SH, Cho YH, Lee HS, Ko HC. Paraplegia after transforaminal epidural steroid injection in a patient with severe lumbar disc herniation - A case report. *Anesth Pain Med (Seoul)* 2021;16:102. . doi: 10.17085/apm.20068 pmid: 33472291
- 50 Thefenne L, Dubecq C, Zing E, et al. A rare case of paraplegia complicating a lumbar epidural infiltration. *Ann Phys Rehabil Med* 2010;53:83. . doi: 10.1016/j.rehab.2010.08.029 pmid: 20870478
- 51 Lo Bianco G, Tinnirello A, Papa A, et al. Interventional pain procedures: A narrative review focusing on safety and complications. Part 1 Injections for spinal pain. *J Pain Res* 2023;16:46. . doi: 10.2147/JPR.S402798 pmid: 37223436
- 52 Lo Bianco G, Tinnirello A, Papa A, et al. Interventional pain procedures: A narrative review focusing on safety and complications. Part 2 Interventional procedures for back pain. *J Pain Res* 2023;16:72. . doi: 10.2147/JPR.S396215 pmid: 36925622
- 53 Carr CM, Plataras CT, Pingree MJ, et al. Immediate adverse events in interventional pain procedures: A multi-institutional study. *Pain Med* 2016;17:61. . doi: 10.1093/pm/pnw051 pmid: 28025351
- 54 Cooper AB, Sharpe MD. Bacterial meningitis and cauda equina syndrome after epidural steroid injections. *Can J Anaesth* 1996;43:4. . doi: 10.1007/BF03018109 pmid: 8723854
- 55 Epstein NE. The risks of epidural and transforaminal steroid injections in the Spine: Commentary and a comprehensive review of the literature. *Surg Neurol Int* 2013;4(Suppl 2):93. . doi: 10.4103/2152-7806.109446 pmid: 23646278
- 56 Glaser SE, Falco F. Paraplegia following a thoracolumbar transforaminal epidural steroid injection. *Pain Physician* 2005;8:14. doi: 10.36076/ppj.2005/8/309 pmid: 16850088
- 57 Huntoon MA, Martin DP. Paralysis after transforaminal epidural injection and previous spinal surgery. *Reg Anesth Pain Med* 2004;29:5. . doi: 10.1016/j.rapm.2004.05.002 pmid: 15372396
- 58 Wybier M, Gaudart S, Petrover D, Houdart E, Laredo JD. Paraplegia complicating selective steroid injections of the lumbar spine. Report of five cases and review of the literature. *Eur Radiol* 2010;20:9. . doi: 10.1007/s00330-009-1539-7 pmid: 19680658
- 59 Food and Drug Administration. FDA Drug Safety Communication: FDA requires label changes to warn of rare but serious neurologic problems after epidural corticosteroid injections for pain. 2014. <https://www.fda.gov/Drugs/DrugSafety/ucm394280.htm>.
- 60 Racoosin JA, Seymour SM, Cascio L, Gill R. Serious neurologic events after epidural glucocorticoid injection--The FDA's risk assessment. *N Engl J Med* 2015;373:301. . doi: 10.1056/NEJMp1511754 pmid: 26444582
- 61 Care calculator by sidecar health. Cost of epidural steroid injection by state. 2023. <https://cost.sidecarhealth.com/ts/epidural-steroid-injection-cost-by-state>.
- 62 Bookimed. Best radiofrequency ablation clinics and costs in 2023. 2023. <https://us-uk.bookimed.com/clinics/procedure=radiofrequency-ablation/>.
- 63 MD Save. Radiofrequency ablation. How much does a radiofrequency ablation cost? 2023. <https://www.mdsave.com/procedures/radiofrequency-ablation/d482f9cb>.
- 64 Thiveos L, Kent P, Pocovic NC, O'Sullivan P, Hancock MJ. Cognitive functional therapy for chronic low back pain: A systematic review and meta-analysis. *Phys Ther* 2024;104:pzae128. . doi: 10.1093/ptj/pzae128 pmid: 39236249
- 65 Hayden JA, Ellis J, Ogilvie R, Malmivaara A, van Tulder MW. Exercise therapy for chronic low back pain. *Cochrane Database Syst Rev* 2021;9:CD009790. .pmid: 34580864
- 66 Ashar YK, Gordon A, Schubiner H, et al. Effect of pain reprocessing therapy vs placebo and usual care for patients with chronic back pain: A randomized clinical trial. *JAMA Psychiatry* 2022;79:23. . doi: 10.1001/jamapsychiatry.2021.2669 pmid: 34586357
- 67 Siemieniuk RA, Agoritsas T, Macdonald H, Guyatt GH, Brandt L, Vandvik PO. Introduction to BMJ Rapid Recommendations. *BMJ* 2016;354: . doi: 10.1136/bmj.i5191 pmid: 27680768
- 68 Guyatt GH, Oxman AD, Vist GE, et al. GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:6. . doi: 10.1136/bmj.39489.470347.AD pmid: 18436948
- 69 Andrews JC, Schünemann HJ, Oxman AD, et al. GRADE guidelines: 15. Going from evidence to recommendation--determinants of a recommendation's direction and strength. *J Clin Epidemiol* 2013;66:35. . doi: 10.1016/j.jclinepi.2013.02.003 pmid: 23570745

Appendix 1: Full list of authors' declarations of interests

Appendix 2: Supplementary tables 1-3

PDF of infographic